

Part I
Amino Acid-Derived Catalysts

1

Proline-Related Secondary Amine Catalysts and Applications

Hiyoshizo Kotsuki and Niiha Sasakura

1.1

Introduction

Since the reappearance of L-proline (**1**) at the forefront of organocatalysis, tremendous efforts have been made to devise new catalysts based on a proline core structure. In this field, the chirality of a pyrrolidine fragment plays a critical role, and the principal concept that underlies the development of new catalysts can be simply explained as the attachment of acidic sites in place of a carboxylic acid group to the side chain. Accordingly, several catalysts with various acidic functionalities have been developed [1]. In general, proline-based organocatalysts can be classified into six major categories: (A) prolinamides, (B) prolinamines, (C) proline tetrazoles, (D) prolinamine sulfonamides, (E) prolinamine thioureas, and (F) diarylprolinols (Figure 1.1). Representative pK_a values of these catalysts are listed in Figure 1.2 [2]. A subtle change in the side-chain assembly may change the pK_a value in the range 8–20, which would have a significant effect on the strength of hydrogen bonding, and thus the catalytic activity and selectivity may be affected.

In this chapter we will deal with organocatalytic asymmetric transformations using these catalysts, mainly focusing on the significant and major achievements in this area published from 2000 to 2011. However, due to space constraints, this chapter will not cover the great utility of diarylprolinol catalysts (category F); for convenience, only references are given [3].

1.2

Prolinamide and Related Catalysts

Owing to the ready availability of prolinamide derivatives through the condensation of proline with amines, prolinamide-based compounds constitute a large family of organocatalysts [4]. Figure 1.3 lists representative examples of these catalysts.




Figure 1.1 Major classes of proline-based organocatalysts.




Figure 1.2 pK_a values of representative proline-related compounds (in DMSO).

These catalysts are very useful in a wide range of asymmetric syntheses. Asymmetric aldol reactions have been investigated by several research groups; selected examples are compiled in Scheme 1.1.

In this context, prolinamide **2** [5–8] and its aryl-substituted homologs such as **3–5** have been developed [9–14]. Analogous to these examples, binaphthyldiamine-derived compounds such as **6** and **7** have been introduced for use in aqueous systems and as recoverable catalysts [15–18]. For example, Benaglia and coworkers reported that the prolinamide catalyst **7** with a lipophilic side chain showed efficient catalytic activity in water [16b]. Chiral spiro diamine-derived catalysts have also been designed, albeit in moderate enantioselectivity [19]. Owing to the increased acidity of an NH group of thioamide relative to a normal amide, proline-thioamide catalysts such as **8** have been shown to be more effective [20–23].






Figure 1.3 Representative examples of prolinamide organocatalysts.



Scheme 1.1

A successful approach in this field is the design of multifunctionalized catalysts such as **9** and **10** through the incorporation of chiral aminoalcohol and related species onto the side chain [24–35]. The high catalytic activity and enantioselectivity of catalyst **9** can be explained by considering the favorable assembly of donor and acceptor molecules via double hydrogen-bonding stabilization, as depicted in the transition state model **15** [24h]. In these examples, to gain satisfactory results, it is important to match the chirality between the proline core and the attachment.



Several other related systems containing a prolinamide or proline-thioamide core have also been reported [36–42].

In some cases, a chiral diamine assembly can serve as an effective scaffold for the design of multiply functionalized catalysts like **11** and **12** [43–49]. Proline hydrazides [50], dipeptides [51–57], or other small peptides [58–63] bearing a prolinamide core have been shown to be useful in asymmetric aldol reactions.

Catalyst **13** was introduced for use in aqueous systems in the presence of surfactant Brønsted acids as co-catalysts [64], and several other catalysts such as **14** containing a heteroaromatic system have also been reported [65–67].

There has been important progress in the use of proline sulfonamides ($pK_a = 8\text{--}11$) [2] as efficient organocatalysts, and Yang and Carter provided an excellent review [68]. Therefore, only a few important aspects are addressed here. Figure 1.4 lists representative examples of these catalysts.

Various *N*-arylsulfonyl-substituted prolinamides such as **16** have been used in asymmetric aldol reactions [69–78]. Carter and coworkers actively sought new efficient catalysts of this type, and found that **17** could serve as an efficient catalyst for asymmetric aldol reactions, even in the absence of any organic solvent, with excellent diastereo- and enantioselectivity (Scheme 1.2) [79].





Figure 1.4 Representative examples of proline sulfonamide organocatalysts.




Scheme 1.2

Recently, Ellman and coworkers have shown that chiral sulfinate **18** can catalyze asymmetric aldol reactions of acetone, whereas proline (**1**) itself gave poor results [80]. Nakamura and coworkers also explored this field, and found that **19** can promote the asymmetric cross-aldol reaction of acetone with activated ketones, to generate a quaternary carbon stereogenic center bearing an OH function [81].


With regard to aldol chemistry, Mannich or domino-Mannich–Michael reactions can also be promoted by *N*-arylsulfonyl-substituted prolinamide catalysts such as **17** with high levels of enantioselectivity [82, 83].

Importantly, prolinamide catalysts work well in Michael addition reactions using nitroolefins as acceptors [58, 64, 84–95]. For example, Nájera and coworkers used bifunctional catalyst **20** by virtue of the synergistic effect of double hydrogen-bonding activation, as depicted in the transition state model **21** (Scheme 1.3) [90]. For the same purpose, prolinamides containing a heteroaromatic system like **14** have also been reported [96].



Scheme 1.3

Analogously to these examples, proline-derived peptide catalysts can also efficiently promote Michael addition reactions [97–99]. Prolinamide or prolyl sulfonamide catalysts are also effective for intramolecular Michael addition reactions [100–102]. Recently, Yang and Carter reported a short-cut strategy to construct an all-carbon substituted quaternary carbon stereogenic center on a cyclohexenone framework via Robinson-type annulation using the **17**-type catalyst (Scheme 1.4) [103].



Scheme 1.4

While some examples of prolinamide-catalyzed enantioselective Biginelli condensation [104, 105] and other types of C–C bond formation [106, 107] are known, their synthetic utility is unclear. Finally, for convenience, with regard to asymmetric heteroatom functionalization and transfer hydrogenation using prolinamides as catalysts, only references are given [108–114].

1.3

Prolinamine and Related Catalysts

Among several organocatalysts derived from L-proline (**1**) as a chiral source, pyrrolidine–tertiary amine conjugates constitute a powerful and useful family in asymmetric synthesis [115]. In 1994, Kawara and Taguchi reported pioneering work on the use of such catalysts in asymmetric Michael addition reactions [116]. Since then, several related catalysts have been developed. Figure 1.5 lists representative examples.

In 2001, after screening several chiral diamines and protonic acid additives, Yamamoto and coworkers reported that a TfOH salt of **22** could efficiently promote asymmetric aldol reactions [117]. Thereafter, similar studies using chiral diamines such as **22–24** with Brønsted or Lewis acid additives have also been reported [118–122]. In 2006, the Mase/Takabe/Barbas groups discovered that prolinamine catalyst **25** with a lipophilic side chain showed efficient catalytic activity in water (Scheme 1.5) [123]. Thus, cyclohexanone reacts smoothly with various aldehydes in water to afford the desired aldol products in high yields with excellent diastereo-





Figure 1.5 Representative examples of prolinamine organocatalysts.

and enantioselectivity. Recently, the recyclability of analogous catalysts has been reported by others [124].




Scheme 1.5

Prolinamine catalyst **26** has been introduced for the same purpose [125].

With regard to asymmetric aldol reactions, it has been shown that prolinamine catalysts such as **22** can also work well for intramolecular aldol [126–128], Henry (nitroaldol) [129], Mannich [130, 131], and domino-Michael–aldol reactions [132] as valuable asymmetric transformations.

Similar to aldol chemistry, prolinamine-catalyzed asymmetric Michael addition reactions have attracted considerable attention from synthetic chemists, and successful examples have been developed (Scheme 1.6).




Scheme 1.6

In this field, prolinamine catalysts **22** and **24** are particularly useful for promoting asymmetric Michael addition reactions between several donor and acceptor molecules [120b, 133–136]. On a related topic, catalyst **27** and related diamine or triamine catalysts have been developed [137–139]. Interestingly, the Mase/Takabe/Barbas groups reported that diamine catalyst **25** could again serve as an efficient catalyst for asymmetric Michael addition reactions even in brine solution [140]. Similarly, several types of water-active catalysts such as **26** have been developed [141, 142].

Independently, Alexakis and coworkers reported that 2,2'-bipyrrolidine catalyst **28** showed excellent catalytic activity in several types of asymmetric Michael addition reactions [143]. It has been postulated that the isopropyl group on one of the C₂-symmetric pyrrolidine rings should block not only the back face against the approach of Michael acceptors but also shift the equilibrium towards one of the two rotamers. Since then, closely related catalysts have also been reported [144]. Furthermore, different types of catalysts such as **29** have been shown to be useful in asymmetric Michael addition reactions [145–148].

While catalyst **22/23** has been known to be valuable in other C–C bond-forming strategies, for example, 1,3-dipolar cycloaddition [149], hetero-Diels–Alder reaction [150], Friedel–Crafts-type alkylation [151], double-Michael reaction [152], [2,3]-Wittig rearrangement [153], and Claisen–Schmidt condensation [154], only references are given here.

Finally, while various reactions under the catalysis of **22** or **24**, for example, asymmetric epoxidation of α,β -unsaturated aldehydes [155], β -hydroalkoxylation of α,β -unsaturated enones [156], and stereoselective reduction of α,β -unsaturated enones (Scheme 1.7) [157], have also been reported, they have been demonstrated in only a limited number of experiments.




Scheme 1.7

1.4

Proline Tetrazole and Related Catalysts


Proline tetrazole catalysts (category C in Figure 1.1) are readily accessible from L-proline (**1**) [158]. They are remarkably useful in asymmetric synthesis [159]. As shown in Figure 1.2, the pK_a of tetrazole is very similar to that of carboxylic acid. Moreover, the advantage of tetrazole catalysts is their robust and lipophilic nature compared to L-proline (**1**) itself, which allows them to escape parasitic bicyclo-oxazolidinone formation [160].

In 2004, Yamamoto and Arvidsson independently reported the catalytic activity of the L-proline tetrazole catalyst **30** in asymmetric aldol reactions of ketones with aldehydes [161–163]. At the same time, Ley and coworkers reached similar conclusions by applying this system to asymmetric Mannich and Michael addition reactions (Scheme 1.8) [164]. Since then, the scope of this chemistry has been expanded by several research groups [165–174].



Scheme 1.8


Ley and coworkers have been quite active in this field, and have found that **30** or its homolog could efficiently promote asymmetric Michael addition reactions using various Michael acceptors and donors [164b, 175]. In these cases, the reactions require the use of a basic amine such as *trans*-4,5-dimethylpiperazine as a co-catalyst to increase the nucleophilicity of donor molecules by deprotonation. Typical examples are shown in Scheme 1.9.



Scheme 1.9

A mechanistic investigation of this chemistry using density functional theory calculations [176] and reactions in ionic liquids as solvents have also been reported [177].


Interestingly, the Michael addition reaction of bromonitromethane to cyclic or acyclic enones constitutes a convenient way of preparing cyclopropane ring compounds in moderate to good enantioselectivity (Scheme 1.10) [178].



Scheme 1.10


Very similar results have also been reported with the use of sulfur ylides as donor molecules [179]. With regard to the asymmetric Biginelli reaction [180] and multicomponent coupling reactions [181] using **30** or its analog as a catalyst, only references are given here.

Finally, we should emphasize the synthetic utility of the **30**-catalyzed α -oxidation of carbonyl compounds via an “O-nitroso aldol reaction” [182]. This method is very attractive as a metal-free oxidation system. For example, Yamamoto and coworkers found that aminoxylation reactions of ketones or aldehydes proceed with almost perfect enantioselectivities (97–99% ee) in the presence of **30** as a catalyst (Scheme 1.11) [183]. Mechanistically, nitroso compounds possess two electrophilic centers, that is, nitrogen and oxygen atoms, but the exclusive formation of O-alkylation products indicates that a hydrogen-bonding transition state like **31** seems to be satisfactorily stabilized with the more basic nitrogen atom. As an extension of this strategy, asymmetric domino-Michael–aldol reactions have also been developed by these authors, and these provide a convenient way to prepare 3-oxa-2-aza-bicycloketone derivatives in high enantioselectivity [184].



Scheme 1.11

Since then, extensive efforts have been made to apply this method to other multiple functionalizations [185–189] or to discover other possible oxidizing agents such as *N*-sulfonyloxaziridines [190]. As a related chemistry, asymmetric amination at the α -position of a carbonyl compound has also been reported with the use of azodicarboxylate esters as electrophiles [187c, 191–194]. Ley and coworkers have developed an ingenious strategy for obtaining chiral dihydropyridazine derivatives by the combination of asymmetric α -amination and Wittig olefination in a one-pot operation (Scheme 1.12) [187c, 192a].



Scheme 1.12





Figure 1.6 Representative examples of proline-based heterocyclic organocatalysts.

As a different family of heteroaromatic-substituted organocatalysts, imidazole- and triazole-based compounds have been known to be quite effective for asymmetric aldol and Michael addition reactions. Figure 1.6 lists representative examples of these catalysts.

The utility of ionic liquid conjugate catalysts such as **32** and **33** can be ascribed to their recyclability [195–199]. On the other hand, triazole-based catalysts such as **34** and **35** are readily accessible via Huisgen 1,3-dipolar cycloadditions, so-called “click reactions,” from azidomethyl-pyrrolidine and acetylenic precursors, and hence make it possible to design new immobilized catalysts [200–207].

Interestingly, it has been shown that the **33**-catalyzed asymmetric S_N1 -type α -alkylation of aldehydes or ketones proceeds well in excellent diastereoselectivity and good enantioselectivity (Scheme 1.13) [208].



Scheme 1.13

1.5 Prolinamine Sulfonamide and Related Catalysts

Prolinamine sulfonamide catalysts (category D in Figure 1.1) can be envisaged as a reversal of prolinamides (A), and constitute a fascinating group of organocatalysts. The catalytic activity of these compounds can be ascribed simply to the sufficient acidity ($pK_a = 10$) of a sulfonamide key structure (Figure 1.2). Figure 1.7 lists representative examples of these catalysts.

For example, in 2004, Wang and coworkers reported a series of asymmetric transformations, for example, α -aminoxylation, Mannich reactions, and α -sulfenylation, under the catalysis of pyrrolidine trifluoromethanesulfonamide **36**; the product yields and diastereo- and enantioselectivities are quite good in most






Figure 1.7 Further representative examples of prolinamine sulfonamide organocatalysts.

cases [209–211]. The proposed mechanism is essentially the same as that in the case of tetrazole catalyst **30** (Scheme 1.14); in contrast to the planar nature of a tetrazole in **30**, a trifluoromethanesulfonyl group in **36** is non-planar. This difference may change the stability of a hydrogen-bonding network at the transition state.



Scheme 1.14

After these reports, the same group extended the utility of this catalytic system to asymmetric Michael addition and aldol reactions [212, 213]. Sulfonamide catalysts such as **37–39** have also been developed for the same purpose [214–219]. The behavior of these catalysts, typically exemplified by enantioselective Michael addition reactions of cyclohexanone with nitroolefins, is compiled in Scheme 1.15.



Scheme 1.15

With regard to the asymmetric α -amination of carbonyl compounds using pyrrolidine sulfonamides as catalysts, only references are given here [220, 221].

1.6

Prolinamine Thiourea and Related Catalysts

Prolinamine thiourea catalysts (category E, Figure 1.1) have been introduced primarily for the purpose of designing new bifunctional catalysts by connecting a pyrrolidine base with a remote hydrogen-bonding thiourea function [222]. Figure 1.8 lists representative examples of these catalysts.






Figure 1.8 Representative examples of prolinamine thiourea conjugates and related organocatalysts.

For example, Tang and coworkers found that bifunctional thiourea catalyst **40** could efficiently promote the asymmetric Michael addition reactions of ketones or α -branched aldehydes with various Michael acceptors (Scheme 1.16) [223]. In particular, the driving force in the present system can be ascribed to the strong hydrogen bond-forming character of thiourea with nitroolefin acceptors, as depicted in the transition state model **46**.



Scheme 1.16

Since then, extensive efforts have been made to devise new powerful catalysts such as **41**–**43** through the modification of a key element of **40**, albeit in most cases with similar or less efficiency [224–231]. Guanidinyl catalysts such as **44** and **45** have been developed to realize the conjugate addition reaction of malonates or nitroalkanes to α,β -unsaturated enones in high enantioselectivity, although the number of experiments has been limited (Scheme 1.17) [232, 233]. A plausible mechanism to account for the (*S*)-configuration of the major products can be ascribed to the transition state model **47**.



Scheme 1.17

Finally, it has been shown that thiourea-type bifunctional catalysts are also useful for asymmetric aldol reactions [234, 235] and α -chlorination of aldehydes [236]. Furthermore, 4-substituted bifunctional analogs have been developed for use in *anti*-selective Mannich reactions [237].

1.7 Miscellaneous

As described so far, various pyrrolidine-based chiral organocatalysts open the door to a remarkably fruitful world of synthetic chemistry. In general, the synthetic protocol used to design new catalysts relies on the naturally occurring chiral source L-proline (**1**) as a key component. This should be a reasonable approach to achieving final success by mimicking “nature.” To characterize newly designed organocatalysts, carbonyl group functionalization, typically through Michael addition and aldol reactions, seems to be the easiest and most useful approach. These transformations are initially driven by the condensation of carbonyl compounds with a





Figure 1.9 Various types of proline-related organocatalysts.


chiral pyrrolidine secondary amine to reversibly form iminium-enamine intermediates, but relatively low enantioselectivities were observed in aldol reactions catalyzed by proline ester congeners [238], except in the case of Loh's catalysts [239]. Hence, several different types of proline-related organocatalysts have been developed. Figure 1.9 lists representative examples of these catalysts.

For example, in 2003, Melchiorre and Jørgensen reported that the enantioselective Michael addition reaction of aldehydes with vinyl ketones proceeded efficiently in the presence of **48** as a catalyst (yield up to 93%, ee up to 85%) [240]. In our research laboratory, we have also been very interested in devising new catalysts with a pyridine ring as a rigid planar base adjacent to a pyrrolidine chiral ring. Along these lines, a series of new catalysts (**49**), that is, DPYMP [49a] and PPYMP [49b], were prepared from L-prolinol, and we found that they showed excellent catalytic activity in terms of productivity, diastereoselectivity, and enantioselectivity (Scheme 1.18) [241]. The results can be explained by invoking the transition state model **55**, in which the pyridinium ring must effectively shield the *Si*-face of an enamine double bond.




Scheme 1.18

For the same purpose, various chiral pyrrolidine catalysts such as **50–54** have also been introduced [242–250]. The versatile nature of pyrrolidine catalysts has been recognized by other transformations: aldol reaction [251], Mannich-type reaction [252, 253], and oxa-Michael reaction [254]. Among these, Maruoka's work on *anti*-selective Mannich reactions is noteworthy (Scheme 1.19, compare with Scheme 1.8) [253]. In this case, the remote hydrogen-bonding form **57** derived from catalyst **56** can overcome the steric preference so that the opposite sense of stereochemistry should be observed.



Scheme 1.19

In 2003, Juhl and Jørgensen found that, after screening a series of pyrrolidine catalysts, catalyst **48** is again of great value for the inverse-electron-demand hetero-Diels–Alder reaction: after pyridinium chlorochromate (PCC) oxidation, lactone products could be obtained as a single diastereomer in excellent enantioselectivity (Scheme 1.20) [255]. The proposed transition state model **58** indicates effective shielding of the *Si*-face of the enamine double bond by the diarylmethyl substituent on the pyrrolidine ring of the catalyst.



Scheme 1.20

A closely related study has also been reported with the use of a **53**-type catalyst [256].





Figure 1.10 Miscellaneous examples of pyrrolidine-based organocatalysts.

Finally, another set of pyrrolidine-derived organocatalysts is listed in Figure 1.10.


In 2004, Jørgensen and coworkers reported that the asymmetric α -chlorination of aldehydes proceeds well in the presence of C_2 -symmetric diphenylpyrrolidine (**59**) as a catalyst using *N*-chlorosuccinimide (NCS) as a chlorinating agent [257]. Thereafter, they also explored its applicability to fluorination (**48**-type catalyst) and bromination (Scheme 1.21) [258].



Scheme 1.21


The synthetic utility of this method is clear: it provides easy manipulation of the products to give various important chiral building blocks such as chlorohydrins, epoxides, aziridines, amino acids, and amino alcohols [257–259], and is readily applicable to natural product synthesis [260].

Recently, considerable efforts have been made to discover new organocatalytic systems for asymmetric epoxidation. In 2003, Aggarwal and coworkers reported that the asymmetric epoxidation of olefins proceeded in good yields and with moderate enantioselectivities using Oxone® (Wako Chemicals, Osaka, Japan) as an oxidant in the presence of a **48**-type catalyst (Scheme 1.22) [261]. According to their proposal, the protonated ammonium salt species can act not only as a phase-transfer catalyst to carry the real oxidant species to the organic phase but also as a promoter to activate the chiral oxidant via hydrogen-bonding stabilization, as depicted in **63**.




Scheme 1.22

On the other hand, Maruoka and coworkers achieved the asymmetric α -benzoyloxylation of aldehydes using the newly designed catalyst **60** (Scheme 1.23) [262].




Scheme 1.23

Novel catalysts **61** and **62** have been invented to increase the catalytic activity by incorporation of an electronegative group (fluorine or azido) at the β -position relative to the NH group: an electrostatic interaction (or *gauche*-effect) between those groups might be favorable for stabilizing reactive intermediates [263–266]. For example, Gilmour and coworkers reported the **61**-catalyzed asymmetric epoxidation of α,β -unsaturated aldehydes (Scheme 1.24) [265].



Scheme 1.24

On the other hand, Zhong and coworkers found that the **62**-catalyzed system was effective for enantioselective [4+1]-annulation using 2-nitroacrylates and α -iodoaldehydes, to form *cis*-isoxazoline *N*-oxide derivatives in high yields and in high diastereo- and enantioselectivity (Scheme 1.25) [266].



Scheme 1.25

1.8

Conclusions

As described above, a great deal of success has been achieved in a wide variety of asymmetric transformations using a series of proline-related organocatalysts. This organocatalytic asymmetric synthesis offers several advantages over metal-catalyzed systems; for example, the ready availability of both enantiomers, ease of handling without the need for an inert atmosphere or anhydrous conditions, and inexpensive and non-toxic reagents. Unfortunately, however, significant limitations still remain to be overcome in this field, including high catalyst loading, a long reaction period, and harmful organic solvent media. We hope that this exceedingly attractive field in modern organic chemistry can lead to new, much more powerful catalysts as well as highly efficient organocatalyst-based asymmetric transformations.

Acknowledgments

The authors would like to acknowledge the past and present members of their research group for their enthusiastic contributions in the field of organocatalysis. The authors also thank Professors Y. Ichikawa and K. Nakano for fruitful discussions and encouragement. Our research project in this field was supported in part by the Yamada Science Foundation, and by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) in Japan (No. 24105523). One of the authors (N.S.) is grateful for a Sasakawa Scientific Research Grant from the Japan Science Society.

References

- Selected reviews: (a) Guillena, G., Nájera, C., and Ramón, D.J. (2007) *Tetrahedron: Asymmetry*, **18**, 2249; (b) Kotsuki, H., Ikishima, H., and Okuyama, A. (2008), *Heterocycles*, **75**, 493; (c) Kotsuki, H., Ikishima, H., and Okuyama, A. (2008) *Heterocycles*, **75**, 757; (d) Panday, S.K. (2011) *Tetrahedron: Asymmetry*, **22**, 1817.
- Bordwell, F.G. (1988) *Acc. Chem. Res.*, **21**, 456; (b) Huang, X.-Y., Wang, H.-J., and Shi, J. (2010) *J. Phys. Chem. A*, **114**, 1068.
- Selected reviews: (a) Palomo, C. and Mielgo, A. (2006) *Angew. Chem., Int. Ed.*, **45**, 7876; (b) Mielgo, A. and Palomo, C. (2008) *Chem. Asian J.*, **3**, 922; (c) Bertelsen, S. and Jørgensen, K.A.

- (2009) *Chem. Soc. Rev.*, **38**, 2178;
- (d) Lattanzi, A. (2009) *Chem. Commun.*, 1452; (e) Xu, L.-W., Li, L., and Shi, Z.-H. (2010) *Adv. Synth. Catal.*, **352**, 243;
- (f) Valero, G., Companyó, X., Bravo, N., Alba, A.-N.R., Moyano, A., and Rios, R. (2010) *Synlett*, 1883.
- 4 Reviews: (a) Liu, X., Lin, L., and Feng, X. (2009) *Chem. Commun.*, 6145; (b) Chen, X.-H., Yu, J., and Gong, L.-Z. (2010) *Chem. Commun.*, **46**, 6437.
- 5 (a) Samanta, S. and Zhao, C.-G. (2006), *Tetrahedron Lett.*, **47**, 3383; (b) Samanta, S. and Zhao, C.-G. (2006) *J. Am. Chem. Soc.*, **128**, 7442; (c) Dodda, R. and Zhao, C.-G. (2006) *Org. Lett.*, **8**, 4911.
- 6 Zhang, F., Su, N., and Gong, Y. (2006) *Synlett*, 1703.
- 7 Aratake, S., Itoh, T., Okano, T., Usui, T., Shoji, M., and Hayashi, Y. (2007) *Chem. Commun.*, 2524.
- 8 Zhang, X.-M., Wang, M., Tu, Y.-Q., Fan, C.-A., Jiang, Y.-J., Zhang, S.-Y., and Zhang, F.-M. (2008) *Synlett*, 2831.
- 9 (a) Chimni, S.S., Mahajan, D., and Suresh Babu, V.V. (2005) *Tetrahedron Lett.*, **46**, 5617; (b) Chimni, S.S. and Mahajan, D. (2006) *Tetrahedron: Asymmetry*, **17**, 2108; (c) Chimni, S.S., Singh, S., and Mahajan, D. (2008) *Tetrahedron: Asymmetry*, **19**, 2276; (d) Chimni, S.S., Singh, S., and Kumar, A. (2009) *Tetrahedron: Asymmetry*, **20**, 1722.
- 10 He, L., Tang, Z., Cun, L.-F., Mi, A.-Q., Jiang, Y.-Z., and Gong, L.-Z. (2006) *Tetrahedron*, **62**, 346.
- 11 Xiong, Y., Wang, F., Dong, S., Liu, X., and Feng, X. (2008) *Synlett*, 73.
- 12 Li, X.-J., Zhang, G.-W., Wang, L., Hua, M.-Q., and Ma, J.-A. (2008) *Synlett*, 1255.
- 13 Moorthy, J.N. and Saha, S. (2009) *Eur. J. Org. Chem.*, 739.
- 14 Xu, J., Fu, X., Wu, C., and Hu, X. (2011) *Tetrahedron: Asymmetry*, **22**, 840.
- 15 (a) Guillena, G., Hita, M. del C., and Nájera, C. (2006) *Tetrahedron: Asymmetry*, **17**, 729; (b) Guillena, G., Hita, M. del C., and Nájera, C. (2006) *Tetrahedron: Asymmetry*, **17**, 1027; (c) Guillena, G., Hita, M. del C., and Nájera, C. (2006) *Tetrahedron: Asymmetry*, **17**, 1493; (d) Guillena, G., Hita, M. del C., Nájera, C., and Vióquez, S.F. (2008) *J. Org. Chem.*, **73**, 5933.
- 16 (a) Guizzetti, S., Benaglia, M., Pignataro, L., and Puglisi, A. (2006) *Tetrahedron: Asymmetry*, **17**, 2754; (b) Guizzetti, S., Benaglia, M., Raimondi, L., and Celentano, G. (2007) *Org. Lett.*, **9**, 1247.
- 17 Gryko, D., Kowalczyk, B., and Zawadzki, L. (2006) *Synlett*, 1059.
- 18 Ma, G.-N., Zhang, Y., and Shi, M. (2007) *Synthesis*, 197.
- 19 Jiang, M., Zhu, S.-F., Yang, Y., Gong, L.-Z., Zhou, X.-G., and Zhou, Q.-L. (2006) *Tetrahedron: Asymmetry*, **17**, 384.
- 20 Review: Gryko, D., Chrominski, M., and Pielacinska, D.J. (2011) *Symmetry*, **3**, 265.
- 21 (a) Gryko, D. and Lipinski, R. (2005) *Adv. Synth. Catal.*, **347**, 1948; (b) Gryko, D. and Lipinski, R. (2006) *Eur. J. Org. Chem.*, 3864; (c) Gryko, D., Zimnicka, M., and Lipinski, R. (2007) *J. Org. Chem.*, **72**, 964; (d) Gryko, D. and Saletta, W.J. (2007) *Org. Biomol. Chem.*, **5**, 2148.
- 22 (a) Almasi, D., Alonso, D.A., and Nájera, C. (2008) *Adv. Synth. Catal.*, **350**, 2467; (b) Almasi, D., Alonso, D.A., Balaguer, A.-N., and Nájera, C. (2009) *Adv. Synth. Catal.*, **351**, 1123.
- 23 Wang, B., Chen, G., Liu, L., Chang, W., and Li, J. (2009) *Adv. Synth. Catal.*, **351**, 2441.
- 24 (a) Tang, Z., Jiang, F., Yu, L.-T., Cui, X., Gong, L.-Z., Mi, A.-Q., Jiang, Y.-Z., and Wu, Y.-D. (2003) *J. Am. Chem. Soc.*, **125**, 5262; (b) Tang, Z., Jiang, F., Cui, X., Gong, L.-Z., Mi, A.-Q., Jiang, Y.-Z., and Wu, Y.-D. (2004) *Proc. Natl. Acad. Sci. U.S.A.*, **101**, 5755; (c) Guo, H.-M., Cun, L.-F., Gong, L.-Z., Mi, A.-Q., and Jiang, Y.-Z. (2005) *Chem. Commun.*, 1450; (d) Tang, Z., Yang, Z.-H., Chen, X.-H., Cun, L.-F., Mi, A.-Q., Jiang, Y.-Z., and Gong, L.-Z. (2005) *J. Am. Chem. Soc.*, **127**, 9285; (e) Xu, X.-Y., Wang, Y.-Z., Cun, L.-F., and Gong, L.-Z. (2007) *Tetrahedron: Asymmetry*, **18**, 237; (f) He, L., Jiang, J., Tang, Z., Cui, X., Mi, A.-Q., Jiang, Y.-Z., and Gong, L.-Z. (2007) *Tetrahedron: Asymmetry*, **18**, 265; (g) Jiang, J., He, L., Luo, S.-W., Cun, L.-F., and Gong, L.-Z. (2007) *Chem. Commun.*, 736; (h) Zhao, J.-F., He, L.,

- Jiang, J., Tang, Z., Cun, L.-F., and Gong, L.-Z. (2008) *Tetrahedron Lett.*, **49**, 3372.
- 25 Tanimori, S., Naka, T., and Kirihaata, M. (2004) *Synth. Commun.*, **34**, 4043.
- 26 (a) Raj, M., Vishnumaya, M.R., Ginotra, S.K., and Singh, V.K. (2006) *Org. Lett.*, **8**, 4097; (b) Maya, V., Raj, M., and Singh, V.K. (2007) *Org. Lett.*, **9**, 2593; (c) Vishnumaya, M.R. and Singh, V.K. (2009) *J. Org. Chem.*, **74**, 4289.
- 27 (a) Song, L., Chen, X., Zhang, S., Zhang, H., Li, P., Luo, G., Liu, W., Duan, W., and Wang, W. (2008) *Org. Lett.*, **10**, 5489; (b) Zhang, H., Zhang, S., Liu, L., Luo, G., Duan, W., and Wang, W. (2010) *J. Org. Chem.*, **75**, 368.
- 28 Schwab, R.S., Galetto, F.Z., Azeredo, J.B., Braga, A.L., Lüdtke, D.S., and Paixao, M.W. (2008) *Tetrahedron Lett.*, **49**, 5094.
- 29 Rulli, G., Duangdee, N., Baer, K., Hummel, W., Berkessel, A., and Gröger, H. (2011) *Angew. Chem., Int. Ed.*, **50**, 7944.
- 30 Gruttadauria, M., Giacalone, F., Marculescu, A.M., Pia Salvo, A.M., and Noto, R. (2009) *ARKIVOC*, **viii**, 5.
- 31 Siyutkin, D.E., Kucherenko, A.S., Frolova, L.L., Kuchin, A.V., and Zlotin, S.G. (2011) *Tetrahedron: Asymmetry*, **22**, 1320.
- 32 Fu, Y.-Q., Li, Z.-C., Ding, L.-N., Tao, J.-C., Zhang, S.-H., and Tang, M.-S. (2006) *Tetrahedron: Asymmetry*, **17**, 3351.
- 33 Wang, C., Jiang, Y., Zhang, X., Huang, Y., Li, B., and Zhang, G. (2007) *Tetrahedron Lett.*, **48**, 4281.
- 34 Du, J., Li, Z., Du, D.-M., and Xu, J. (2008) *ARKIVOC*, **xvii**, 145.
- 35 (a) Pearson, A.J. and Panda, S. (2011) *Tetrahedron*, **67**, 3969; (b) Pearson, A.J. and Panda, S. (2011) *Org. Lett.*, **13**, 5548.
- 36 Tsutsui, A., Takeda, H., Kimura, M., Fujimoto, T., and Machinami, T. (2007) *Tetrahedron Lett.*, **48**, 5213.
- 37 Doherty, S., Knight, J.G., McRae, A., Harrington, R.W., and Clegg, W. (2008) *Eur. J. Org. Chem.*, 1759.
- 38 Saha, S. and Moorthy, J.N. (2010) *Tetrahedron Lett.*, **51**, 912.
- 39 Rani, R. and Peddinti, R.K. (2010) *Tetrahedron: Asymmetry*, **21**, 775.
- 40 (a) Puleo, G.L., Masi, M., and Iuliano, A. (2007) *Tetrahedron: Asymmetry*, **18**, 1364; (b) Puleo, G.L. and Iuliano, A. (2007) *Tetrahedron: Asymmetry*, **18**, 2894.
- 41 Pedrosa, R., Andrés, J.M., Manzano, R., Román, D., and Téllez, S. (2011) *Org. Biomol. Chem.*, **9**, 935.
- 42 Wang, B., Liu, X., Liu, L., Chang, W., and Li, J. (2010) *Eur. J. Org. Chem.*, 5951.
- 43 (a) Chen, J.-R., Lu, H.-H., Li, X.-Y., Cheng, L., Wan, J., and Xiao, W.-J. (2005) *Org. Lett.*, **7**, 4543; (b) Chen, J.-R., Li, X.-Y., Xing, X.-N., and Xiao, W.-J. (2006) *J. Org. Chem.*, **71**, 8198; (c) Huang, W.-P., Chen, J.-R., Li, X.-Y., Cao, Y.-J., and Xiao, W.-J. (2007) *Can. J. Chem.*, **85**, 208; (d) Chen, J.-R., Liu, X., Zhu, X.-Y., Li, L., Qiao, Y.-F., Zhang, J.-M., and Xiao, W.-J. (2007) *Tetrahedron*, **63**, 10437.
- 44 Samanta, S., Liu, J., Dodda, R., and Zhao, C.-G. (2005) *Org. Lett.*, **7**, 5321.
- 45 Gandhi, S. and Singh, V.K. (2008) *J. Org. Chem.*, **73**, 9411.
- 46 Jia, Y.-N., Wu, F.-C., Ma, X., Zhu, G.-J., and Da, C.-S. (2009) *Tetrahedron Lett.*, **50**, 3059.
- 47 Yang, Y., He, Y.-H., Guan, Z., and Huang, W.-D. (2010) *Adv. Synth. Catal.*, **352**, 2579.
- 48 Fotaras, S., Kokotos, C.G., Tsandi, E., and Kokotos, G. (2011) *Eur. J. Org. Chem.*, 1310.
- 49 Li, Y., Yang, Q., Xu, X., Zhou, Y., Bai, J., Wang, F., and Wang, L. (2011) *Can. J. Chem.*, **89**, 1312.
- 50 (a) Cheng, C., Sun, J., Wang, C., Zhang, Y., Wei, S., Jiang, F., and Wu, Y. (2006) *Chem. Commun.*, 215; (b) Cheng, C., Wei, S., and Sun, J. (2006) *Synlett*, 2419.
- 51 (a) Luppi, G., Cozzi, P.G., Monari, M., Kaptein, B., Broxterman, Q.B., and Tomasini, C. (2005) *J. Org. Chem.*, **70**, 7418; (b) Luppi, G., Monari, M., Correa, R.J., Violante, F. de A., Pinto, A.C., Kaptein, B., Broxterman, Q.B., Garden, S.J., and Tomasini, C. (2006) *Tetrahedron*, **62**, 12017; (c) Angelici, G., Correa, R.J., Garden, S.J., and Tomasini, C. (2009) *Tetrahedron Lett.*, **50**, 814.
- 52 Shi, L.-X., Sun, Q., Ge, Z.-M., Zhu, Y.-Q., Cheng, T.-M., and Li, R.-T. (2004) *Synlett*, 2215.
- 53 Erkkilä, A. and Pihko, P.M. (2006) *J. Org. Chem.*, **71**, 2538.

- 54 Dodda, R. and Zhao, C.-G. (2007) *Synlett*, 1605.
- 55 Yan, J. and Wang, L. (2008) *Synthesis*, 2065.
- 56 Alberg, D.G., Poulsen, T.B., Bertelsen, S., Christensen, K.L., Birkler, R.D., Johannsen, M., and Jørgensen, K.A. (2009) *Bioorg. Med. Chem. Lett.*, **19**, 3888.
- 57 Lei, M., Shi, L., Li, G., Chen, S., Fang, W., Ge, Z., Cheng, T., and Li, R. (2007) *Tetrahedron*, **63**, 7892.
- 58 Martin, H.J. and List, B. (2003) *Synlett*, 1901.
- 59 Kofoed, J., Nielsen, J., and Reymond, J.-L. (2003) *Bioorg. Med. Chem. Lett.*, **13**, 2445.
- 60 Szöllosi, G., London, G., Baláspiri, L., Somlai, C., and Bartók, M. (2003) *Chirality*, **15**, S90.
- 61 Krattiger, P., Kovasy, R., Revell, J.D., Ivan, S., and Wennemers, H. (2005) *Org. Lett.*, **7**, 1101.
- 62 (a) Tang, Z., Yang, Z.-H., Cun, L.-F., Gong, L.-Z., Mi, A.-Q., and Jiang, Y.-Z. (2004) *Org. Lett.*, **6**, 2285; (b) Chen, X.-H., Luo, S.-W., Tang, Z., Cun, L.-F., Mi, A.-Q., Jiang, Y.-Z., and Gong, L.-Z. (2007) *Chem. Eur. J.*, **13**, 689.
- 63 Carpenter, R.D., Fettinger, J.C., Lam, K.S., and Kurth, M.J. (2008) *Angew. Chem., Int. Ed.*, **47**, 6407.
- 64 Luo, S., Xu, H., Li, J., Zhang, L., Mi, X., Zheng, X., and Cheng, J.-P. (2007) *Tetrahedron*, **63**, 11307.
- 65 Tang, Z., Cun, L.-F., Cui, X., Mi, A.-Q., Jiang, Y.-Z., and Gong, L.-Z. (2006) *Org. Lett.*, **8**, 1263.
- 66 Kinsella, M., Duggan, P.G., and Lennon, C.M. (2011) *Tetrahedron: Asymmetry*, **22**, 1423.
- 67 Zheng, J.-F., Li, Y.-X., Zhang, S.-Q., Yang, S.-T., Wang, X.-M., Wang, Y.-Z., Bai, J., and Liu, F.-A. (2006) *Tetrahedron Lett.*, **47**, 7793.
- 68 Review: Yang, H. and Carter, R.G. (2010) *Synlett*, 2827.
- 69 (a) Berkessel, A., Koch, B., and Lex, J. (2004) *Adv. Synth. Catal.*, **346**, 1141; (b) Meciárová, M., Toma, S., Berkessel, A., and Koch, B. (2006) *Let. Org. Chem.*, **3**, 437.
- 70 Bellis, E., Vasilatou, K., and Kokotos, G. (2005) *Synthesis*, 2407.
- 71 Wu, Y., Zhang, Y., Yu, M., Zhao, G., and Wang, S. (2006) *Org. Lett.*, **8**, 4417.
- 72 Silva, F., Sawicki, M., and Gouverneur, V. (2006) *Org. Lett.*, **8**, 5417.
- 73 Wang, X.-J., Zhao, Y., and Liu, J.-T. (2007) *Org. Lett.*, **9**, 1343.
- 74 Bernard, A.M., Frongia, A., Piras, P.P., Secci, F., and Spiga, M. (2008) *Tetrahedron Lett.*, **49**, 3037.
- 75 Carboni, D., Flavin, K., Servant, A., Gouverneur, V., and Resmini, M. (2008) *Chem. Eur. J.*, **14**, 7059.
- 76 (a) Zhang, S., Fu, X., Fu, S., and Pan, J. (2009) *Catal. Commun.*, **10**, 401; (b) Fu, S., Fu, X., Zhang, S., Zou, X., and Wu, X. (2009) *Tetrahedron: Asymmetry*, **20**, 2390.
- 77 Srivastava, V. (2010) *Cent. Eur. J. Chem.*, **8**, 269.
- 78 Tang, G., Hu, X., and Altenbach, H.J. (2011) *Tetrahedron Lett.*, **52**, 7034.
- 79 (a) Yang H. and Carter, R.G. (2008) *Org. Lett.*, **10**, 4649; (b) Yang, H., Mahapatra, S., Cheong, P.H.-Y., and Carter, R.G. (2010) *J. Org. Chem.*, **75**, 7279.
- 80 Robak, M.T., Herbage, M.A., and Ellman, J.A. (2011) *Tetrahedron*, **67**, 4412.
- 81 (a) Nakamura, S., Hara, N., Nakashima, H., Kubo, K., Shibata, N., and Toru, T. (2008) *Chem. Eur. J.*, **14**, 8079; (b) Hara, N., Nakamura, S., Shibata, N., and Toru, T. (2009) *Chem. Eur. J.*, **15**, 6790; (c) Hara, N., Nakamura, S., Shibata, N., and Toru, T. (2010) *Adv. Synth. Catal.*, **352**, 1621; (d) Hara, N., Tamura, R., Funahashi, Y., and Nakamura, S. (2011) *Org. Lett.*, **13**, 1662.
- 82 (a) Yang, H. and Carter, R.G. (2009) *J. Org. Chem.*, **74**, 2246; (b) Yang, H. and Carter, R.G. (2009) *J. Org. Chem.*, **74**, 5151; (c) Yang, H. and Carter, R.G. (2010) *Tetrahedron*, **66**, 4854.
- 83 Veverková, E., Strasserová, J., Sebesta, R., and Toma, S. (2010) *Tetrahedron: Asymmetry*, **21**, 58.
- 84 Palomo, C., Vera, S., Mielgo, A., and Gómez-Bengoia, E. (2006) *Angew. Chem., Int. Ed.*, **45**, 5984.
- 85 Meciárová, M., Hubinská, K., Toma, S., Koch, B., and Berkessel, A. (2007) *Monatsh. Chem.*, **138**, 1181.

- 86 Reddy, R.J., Kuan, H.-H., Chou, T.-Y., and Chen, K. (2009) *Chem. Eur. J.*, **15**, 9294.
- 87 Rani, R. and Peddinti, R.K. (2010) *Tetrahedron: Asymmetry*, **21**, 2487.
- 88 Moorthy, J.N. and Saha, S. (2010) *Eur. J. Org. Chem.*, 6359.
- 89 Puleo, G.L. and Iuliano, A. (2008) *Tetrahedron: Asymmetry*, **19**, 2045.
- 90 (a) Almasi, D., Alonso, D.A., and Nájera, C. (2006) *Tetrahedron: Asymmetry*, **17**, 2064; (b) Almasi, D., Alonso, D.A., Gómez-Bengoia, E., Nagel, Y., and Nájera, C. (2007) *Eur. J. Org. Chem.*, 2328.
- 91 Okuyama, Y., Nakano, H., Watanabe, Y., Makabe, M., Takeshita, M., Uwai, K., Kabuto, C., and Kwon, E. (2009) *Tetrahedron Lett.*, **50**, 193.
- 92 (a) Wiesner, M., Revell, J.D., Tonazzi, S., and Wennemers, H. (2008) *J. Am. Chem. Soc.*, **130**, 5610; (b) Wiesner, M., Upert, G., Angelici, G., and Wennemers, H. (2010) *J. Am. Chem. Soc.*, **132**, 6.
- 93 Wei, S., Yalalov, D.A., Tsogoeva, S.B., and Schmatz, S. (2007) *Catal. Today*, **121**, 151.
- 94 (a) Bai, J.-F., Xu, X.-Y., Huang, Q.-C., Peng, L., and Wang, L.-X. (2010) *Tetrahedron Lett.*, **51**, 2803; (b) Wang, Q., Peng, L., Fu, J., Huang, Q., Wang, L., and Xu, X. (2010) *ARKIVOC*, ii, 340.
- 95 Wang, W.-H., Wang, X.-B., Kodama, K., Hirose, T., and Zhang, G.-Y. (2010) *Tetrahedron*, **66**, 4970.
- 96 (a) Clarke, M.L. and Fuentes, J.A. (2007) *Angew. Chem., Int. Ed.*, **46**, 930; (b) Fuentes, J.A., Lebl, T., Slawin, A.M.Z., and Clarke, M.L. (2011) *Chem. Sci.*, **2**, 1997.
- 97 Znabet, A., Ruijter, E., de Kanter, F.J.J., Köhler, V., Helliwell, M., Turner, N.J., and Orru, R.V.A. (2010) *Angew. Chem., Int. Ed.*, **49**, 5289.
- 98 Arakawa, Y., Wiesner, M., and Wennemers, H. (2011) *Adv. Synth. Catal.*, **353**, 1201.
- 99 Durini, M., Sahr, F.A., Kuhn, M., Civera, M., Gennari, C., and Piarulli, U. (2011) *Eur. J. Org. Chem.*, 5599.
- 100 Kikuchi, M., Inagaki, T., and Nishiyama, H. (2007) *Synlett*, 1075.
- 101 Bournaud, C., Marchal, E., Quintard, A., Sulzer-Mossé, S., and Alexakis, A. (2010) *Tetrahedron: Asymmetry*, **21**, 1666.
- 102 Yang, H. and Carter, R.G. (2010) *J. Org. Chem.*, **75**, 4929.
- 103 Yang, H. and Carter, R.G. (2010) *Org. Lett.*, **12**, 3108.
- 104 Xin, J., Chang, L., Hou, Z., Shang, D., Liu, X., and Feng, X. (2008) *Chem. Eur. J.*, **14**, 3177.
- 105 Saha, S. and Moorthy, J.N. (2011) *J. Org. Chem.*, **76**, 396.
- 106 Wei, Y., Ma, G.-N., and Shi, M. (2011) *Eur. J. Org. Chem.*, 5146.
- 107 Hartikka, A., Slóarsczyk, A.T., and Arvidsson, P.I. (2007) *Tetrahedron: Asymmetry*, **18**, 1403.
- 108 Sohtome, Y., Tanatani, A., Hashimoto, Y., and Nagasawa, K. (2004) *Chem. Pharm. Bull.*, **52**, 477.
- 109 (a) Fu, J.-Y., Xu, X.-Y., Li, Y.-C., Huang, Q.-C., and Wang, L.-X. (2010) *Org. Biomol. Chem.*, **8**, 4524; (b) Fu, J.-Y., Huang, Q.-C., Wang, Q.-W., Wang, L.-X., and Xu, X.-Y. (2010) *Tetrahedron Lett.*, **51**, 4870.
- 110 Sundén, H., Dahlin, N., Ibrahim, I., Adolfsson, H., and Córdova, A. (2005) *Tetrahedron Lett.*, **46**, 3385.
- 111 Guo, H.-M., Cheng, L., Cun, L.-F., Gong, L.-Z., Mi, A.-Q., and Jiang, Y.-Z. (2006) *Chem. Commun.*, 429.
- 112 (a) Wang, W., Wang, J., and Li, H. (2004) *Org. Lett.*, **6**, 2817; (b) Wang, J., Li, H., Mei, Y., Lou, B., Xu, D., Xie, D., Guo, H., and Wang, W. (2005) *J. Org. Chem.*, **70**, 5678.
- 113 Wong, M.-K., Ho, L.-M., Zheng, Y.-S., Ho, C.-Y., and Yang, D. (2001) *Org. Lett.*, **3**, 2587.
- 114 Zhang, Z., Rooshenas, P., Hausmann, H., and Schreiner, P.R. (2009) *Synthesis*, 1531.
- 115 Reviews: (a) Saito, S. and Yamamoto, H. (2004) *Acc. Chem. Res.*, **37**, 570; (b) Notz, W., Tanaka, F., and Barbas, C.F. III (2004) *Acc. Chem. Res.*, **37**, 580; (c) Chowdhury, R. (2011) *Synlett*, 1784.
- 116 Kawara, A. and Taguchi, T. (1994) *Tetrahedron Lett.*, **35**, 8805.
- 117 (a) Saito, S., Nakadai, M., and Yamamoto, H. (2001) *Synlett*, 1245; (b) Nakadai, M., Saito, S., and Yamamoto, H. (2002) *Tetrahedron*, **58**, 8167.
- 118 (a) Córdova, A., Notz, W., and Barbas, C.F. III (2002) *Chem. Commun.*, 3024;

- (b) Mase, N., Tanaka, F., and Barbas, C.F. III (2004) *Angew. Chem., Int. Ed.*, **43**, 2420.
- 119 Damburuoso, P., Massi, A., and Dondoni, A. (2005) *Org. Lett.*, **7**, 4657.
- 120 (a) Luo, S., Li, J., Xu, H., Zhang, L., and Cheng, J.-P. (2007) *Org. Lett.*, **9**, 3675; (b) Luo, S., Li, J., Zhang, L., Xu, H., and Cheng, J.-P. (2008) *Chem. Eur. J.*, **14**, 1273.
- 121 Catalán-Munoz, S., Müller, C.A., and Ley, S.V. (2010) *Eur. J. Org. Chem.*, 183.
- 122 Massi, A., Cavazzini, A., Del Zoppo, L., Pandoli, O., Costa, V., Pasti, L., and Giovannini, P.P. (2011) *Tetrahedron Lett.*, **52**, 619.
- 123 (a) Mase, N., Nakai, Y., Ohara, N., Yoda, H., Takabe, K., Tanaka, F., and Barbas, C.F. III (2006) *J. Am. Chem. Soc.*, **128**, 734; (b) Mase, N., Noshiro, N., Mokuya, A., and Takabe, K. (2009) *Adv. Synth. Catal.*, **351**, 2791.
- 124 Gao, Q., Lu, S.-M., Liu, Y., and Li, C. (2011) *Tetrahedron Lett.*, **52**, 3779.
- 125 Bisai, V. and Singh, V.K. (2011) *Synlett*, 481.
- 126 Hayashi, Y., Sekizawa, H., Yamaguchi, J., and Gotoh, H. (2007) *J. Org. Chem.*, **72**, 6493.
- 127 Chen, L., Luo, S., Li, J., Li, X., and Cheng, J.-P. (2010) *Org. Biomol. Chem.*, **8**, 2627.
- 128 Akahane, Y., Inomata, K., and Endo, Y. (2011) *Heterocycles*, **82**, 1727.
- 129 Zhou, Y., Dong, J., Zhang, F., and Gong, Y. (2011) *J. Org. Chem.*, **76**, 588.
- 130 Zhuang, W., Saaby, S., and Jørgensen, K.A. (2004) *Angew. Chem., Int. Ed.*, **43**, 4476.
- 131 Li, H., Fu, A., and Shi, H. (2009) *J. Mol. Catal. A Chem.*, **303**, 1.
- 132 (a) Rios, R., Sundén, H., Ibrahim, I., Zhao, G.-L., and Córdova, A. (2006) *Tetrahedron Lett.*, **47**, 8679; (b) Rios, R., Sundén, H., Ibrahim, I., and Córdova, A. (2007) *Tetrahedron Lett.*, **48**, 2181.
- 133 (a) Betancort, J.M., Sakthivel, K., Thayumanavan, R., and Barbas, C.F. III (2001) *Tetrahedron Lett.*, **42**, 4441; (b) Betancort, J.M. and Barbas, C.F. III (2001) *Org. Lett.*, **3**, 3737; (c) Mase, N., Thayumanavan, R., Tanaka, F., and Barbas, C.F. III (2004) *Org. Lett.*, **6**, 2527; (d) Betancort, J.M., Sakthivel, K., Thayumanavan, R., Tanaka, F., and Barbas, C.F. III (2004) *Synthesis*, 1509.
- 134 Barros, M.T. and Faisca Phillips, A.M. (2008) *Eur. J. Org. Chem.*, 2525.
- 135 (a) Chowdhury, R. and Ghosh, S.K. (2009) *Org. Lett.*, **11**, 3270; (b) Chowdhury, R. and Ghosh, S.K. (2011) *Synthesis*, 1936.
- 136 Mase, N., Fukasawa, M., Kitagawa, N., Shibagaki, F., Noshiro, N., and Takabe, K. (2010) *Synlett*, 2340.
- 137 Zhu, M.-K., Cun, L.-F., Mi, A.-Q., Jiang, Y.-Z., and Gong, L.-Z. (2006) *Tetrahedron: Asymmetry*, **17**, 491.
- 138 Chen, H., Wang, Y., Wei, S., and Sun, J. (2007) *Tetrahedron: Asymmetry*, **18**, 1308.
- 139 Pansare, S.V. and Pandya, K. (2006) *J. Am. Chem. Soc.*, **128**, 9624.
- 140 Mase, N., Watanabe, K., Yoda, H., Takabe, K., Tanaka, F., and Barbas, C.F. III (2006) *J. Am. Chem. Soc.*, **128**, 4966.
- 141 Vishnumaya M.R. and Singh, V.K. (2007) *Org. Lett.*, **9**, 1117.
- 142 Lo, C.-M. and Chow, H.-F. (2009) *J. Org. Chem.*, **74**, 5181.
- 143 (a) Alexakis, A. and Andrey, O. (2002) *Org. Lett.*, **4**, 3611; (b) Andrey, O., Alexakis, A., and Bernardinelli, G. (2003) *Org. Lett.*, **5**, 2559; (c) Andrey, O., Vidonne, A., and Alexakis, A. (2003) *Tetrahedron Lett.*, **44**, 7901; (d) Andrey, O., Alexakis, A., Tomassini, A., and Bernardinelli, G. (2004) *Adv. Synth. Catal.*, **346**, 1147; (e) Mossé, S. and Alexakis, A. (2005) *Org. Lett.*, **7**, 4361; (f) Mossé, S. and Alexakis, A. (2006) *Org. Lett.*, **8**, 3577; (g) Belot, S., Sulzer-Mossé, S., Kehrl, S., and Alexakis, A. (2008) *Chem. Commun.*, 4694; (h) Sulzer-Mossé, S., Alexakis, A., Mareda, J., Bollot, G., Bernardinelli, G., and Filinchuk, Y. (2009) *Chem. Eur. J.*, **15**, 3204.
- 144 Jin, S., Li, C., Ma, Y., Kan, Y., Zhang, Y.J., and Zhang, W. (2010) *Org. Biomol. Chem.*, **8**, 4011.
- 145 Xu, D.-Z., Liu, Y., Shi, S., and Wang, Y. (2010) *Tetrahedron: Asymmetry*, **21**, 2530.
- 146 Pansare, S.V., Lingampally, R., and Kirby, R.L. (2010) *Org. Lett.*, **12**, 556.
- 147 Anwar, S., Lee, P.-H., Chou, T.-Y., Chang, C., and Chen, K. (2011) *Tetrahedron*, **67**, 1171.

- 148 Quintard, A., Bournaud, C., and Alexakis, A. (2008) *Chem. Eur. J.*, **14**, 7504.
- 149 (a) Karlsson, S. and Högberg, H.-E. (2002) *Tetrahedron: Asymmetry*, **13**, 923; (b) Karlsson, S. and Högberg, H.-E. (2003) *Eur. J. Org. Chem.*, 2782.
- 150 Wabnitz, T.C., Saaby, S., and Jørgensen, K.A. (2004) *Org. Biomol. Chem.*, **2**, 828.
- 151 Breistein, P., Karlsson, S., and Hedenström, E. (2006) *Tetrahedron: Asymmetry*, **17**, 107.
- 152 (a) Thayumanavan, R., Dhevalapally, B., Sakthivel, K., Tanaka, F., and Barbas, C.F. III (2002) *Tetrahedron Lett.*, **43**, 3817; (b) Ramachary, D.B., Chowdari, N.S., and Barbas, C.F. III (2002) *Tetrahedron Lett.*, **43**, 6743.
- 153 McNally, A., Evans, B., and Gaunt, M.J. (2006) *Angew. Chem., Int. Ed.*, **45**, 2116.
- 154 (a) Ramachary, D.B., Ramakumar, K., and Kishor, M. (2005) *Tetrahedron Lett.*, **46**, 7037; (b) Ramachary, D.B., Ramakumar, K., Bharanishashank, A., and Narayana, V.V. (2010) *J. Comb. Chem.*, **12**, 855.
- 155 Sundén, H., Ibrahim, I., and Córdova, A. (2006) *Tetrahedron Lett.*, **47**, 99.
- 156 Ramachary, D.B. and Mondal, R. (2006) *Tetrahedron Lett.*, **47**, 7689.
- 157 Ramachary, D.B. and Sakthidevi, R. (2008) *Org. Biomol. Chem.*, **6**, 2488.
- 158 (a) Aureggi, V., Franckevicius, V., Kitching, M.O., Ley, S.V., Longbottom, D.A., Oelke, A.J., and Sedelmeier, G. (2008) *Org. Synth.*, **85**, 72; (b) Franckevicius, V., Knudsen, K.R., Ladlow, M., Longbottom, D.A., and Ley, S.V. (2006) *Synlett*, 889.
- 159 Reviews: (a) Limbach, M. (2006) *Chem. Biodivers.*, **3**, 119; (b) Longbottom, D.A., Franckevicius, V., and Ley, S.V. (2007) *Chimia*, **61**, 247; (c) Longbottom, D.A., Franckevicius, V., Kumarn, S., Oelke, A.J., Wascholowski, V., and Ley, S.V. (2008) *Aldrichim. Acta*, **41**, 3.
- 160 For example, see: (a) Kano, T., Takai, J., Tokuda, O., and Maruoka, K. (2005) *Angew. Chem., Int. Ed.*, **44**, 3055; (b) Seebach, D., Beck, A.K., Badine, D.M., Limbach, M., Eschenmoser, A., Treasurywala, A.M., Hobi, R., Prikoszovich, W., and Linder, B. (2007) *Helv. Chim. Acta*, **90**, 425; (c) Isart, C., Burés, J., and Vilarrasa, J. (2008) *Tetrahedron Lett.*, **49**, 5414; (d) Companyó, X., Valero, G., Crovetto, L., Moyano, A., and Rios, R. (2009) *Chem. Eur. J.*, **15**, 6564.
- 161 Torii, H., Nakadai, M., Ishihara, K., Saito, S., and Yamamoto, H. (2004) *Angew. Chem., Int. Ed.*, **43**, 1983.
- 162 (a) Hartikka, A. and Arvidsson, P.I. (2004) *Tetrahedron: Asymmetry*, **15**, 1831; (b) Hartikka, A. and Arvidsson, P.I. (2005) *Eur. J. Org. Chem.*, 4287.
- 163 Jheengut, V. and Ward, D.E. (2007) *J. Org. Chem.*, **72**, 7805.
- 164 (a) Cobb, A.J.A., Shaw, D.M., and Ley, S.V. (2004) *Synlett*, 558; (b) Cobb, A.J.A., Shaw, D.M., Longbottom, D.A., Gold, J.B., and Ley, S.V. (2005) *Org. Biomol. Chem.*, **3**, 84.
- 165 Arnó, M., Zaragoza, R.J., and Domingo, L.R. (2005) *Tetrahedron: Asymmetry*, **16**, 2764.
- 166 Chowdari, N.S., Ahmad, M., Albertshofer, K., Tanaka, F., and Barbas, C.F. III (2006) *Org. Lett.*, **8**, 2839.
- 167 Hayashi, Y., Urushima, T., Aratake, S., Okano, T., and Obi, K. (2008) *Org. Lett.*, **10**, 21.
- 168 Tong, S.-T. (Amy), Harris, P.W.R., Barker, D., and Brimble, M.A. (2008) *Eur. J. Org. Chem.*, 164.
- 169 Xue, F., Zhang, S., Liu, L., Duan, W., and Wang, W. (2009) *Chem. Asian J.*, **4**, 1664.
- 170 Odedra, A. and Seeberger, P.H. (2009) *Angew. Chem., Int. Ed.*, **48**, 2699.
- 171 Ward, D.E., Jheengut, V., Beyé, G.E., Gillis, H.M., Karagiannis, A., and Becerril-Jimenez, F. (2011) *Synlett*, 508.
- 172 Funabiki, K., Itoh, Y., Kubota, Y., and Matsui, M. (2011) *J. Org. Chem.*, **76**, 3545.
- 173 Zhou, S., Zhou, Y., Xing, Y., Wang, N., and Cao, L. (2011) *Chirality*, **23**, 504.
- 174 Komatsu, Y., Watanabe, R., Ikishima, H., Nakano, K., Ichikawa, Y., and Kotsuki, H. (2012) *Org. Biomol. Chem.*, **10**, 2993.
- 175 (a) Cobb, A.J.A., Longbottom, D.A., Shaw, D.M., and Ley, S.V. (2004) *Chem. Commun.*, 1808; (b) Mitchell, C.E.T., Cobb, A.J.A., and Ley, S.V. (2005) *Synlett*, 611; (c) Mitchell, C.E.T., Brenner, S.E., and Ley, S.V. (2005)

- Chem. Commun.*, 5346; (d) Knudsen, K.R., Mitchell, C.E.T., and Ley, S.V. (2006) *Chem. Commun.*, 66; (e) Mitchell, C.E.T., Brenner, S.E., García-Fortanet, J., and Ley, S.V. (2006) *Org. Biomol. Chem.*, 4, 2039; (f) Wascholowski, V., Knudsen, K.R., Mitchell, C.E.T., and Ley, S.V. (2008) *Chem. Eur. J.*, 14, 6155.
- 176 Arnó, M., Zaragoza, R.J., and Domingo, L.R. (2007) *Tetrahedron: Asymmetry*, 18, 157.
- 177 Meciárová, M., Toma, S., and Sebesta, R. (2009) *Tetrahedron: Asymmetry*, 20, 2403.
- 178 (a) Hansen, H.M., Longbottom, D.A., and Ley, S.V. (2006) *Chem. Commun.*, 4838; (b) Wascholowski, V., Hansen, H.M., Longbottom, D.A., and Ley, S.V. (2008) *Synthesis*, 1269.
- 179 Hartikka, A. and Arvidsson, P.I. (2007) *J. Org. Chem.*, 72, 5874.
- 180 Wu, Y.-Y., Chai, Z., Liu, X.-Y., Zhao, G., and Wang, S.-W. (2009) *Eur. J. Org. Chem.*, 904.
- 181 Su, W., Li, J., and Li, J. (2008) *Aust. J. Chem.*, 61, 860.
- 182 Reviews: (a) Merino, P. and Tejero, T. (2004) *Angew. Chem., Int. Ed.*, 43, 2995; (b) Yamamoto, H. and Momiyama, N. (2005) *Chem. Commun.*, 3514; (c) Janey, J.M. (2005) *Angew. Chem., Int. Ed.*, 44, 4292; (d) Marigo, M. and Jørgensen, K.A. (2006) *Chem. Commun.*, 2001; (e) Guillena, G. and Ramón, D. (2006) *Tetrahedron: Asymmetry*, 17, 1465.
- 183 (a) Momiyama, N., Torii, H., Saito, S., and Yamamoto, H. (2004) *Proc. Natl. Acad. Sci. U.S.A.*, 101, 5374; (b) Jiao, P., Kawasaki, M., and Yamamoto, H. (2009) *Angew. Chem., Int. Ed.*, 48, 3333; (c) Jiao, P. and Yamamoto, H. (2009) *Synlett*, 2685.
- 184 (a) Yamamoto, Y., Momiyama, N., and Yamamoto, H. (2004) *J. Am. Chem. Soc.*, 126, 5962; (b) Momiyama, N., Yamamoto, Y., and Yamamoto, H. (2007) *J. Am. Chem. Soc.*, 129, 1190.
- 185 Ramachary, D.B. and Barbas, C.F. III (2005) *Org. Lett.*, 7, 1577.
- 186 Kim, S.-G. and Park, T.-H. (2006) *Tetrahedron Lett.*, 47, 9067.
- 187 (a) Kumarn, S., Shaw, D.M., Longbottom, D.A., and Ley, S.V. (2005) *Org. Lett.*, 7, 4189; (b) Kumarn, S., Shaw, D.M., and Ley, S.V. (2006) *Chem. Commun.*, 3211; (c) Kumarn, S., Oelke, A.J., Shaw, D.M., Longbottom, D.A., and Ley, S.V. (2007) *Org. Biomol. Chem.*, 5, 2678.
- 188 Lu, M., Zhu, D., Lu, Y., Hou, Y., Tan, B., and Zhong, G. (2008) *Angew. Chem., Int. Ed.*, 47, 10187.
- 189 Capitta, F., Frongia, A., Ollivier, J., Piras, P.P., and Secci, F. (2011) *Synlett*, 89.
- 190 Tong, S.-T., Brimble, M.A., and Barker, D. (2009) *Tetrahedron*, 65, 4801.
- 191 Chowdari, N.S. and Barbas, C.F. III (2005) *Org. Lett.*, 7, 867.
- 192 (a) Oelke, A.J., Kumarn, S., Longbottom, D.A., and Ley, S.V. (2006) *Synlett*, 2548; (b) Oelke, A.J., France, D.J., Hofmann, T., Wuitschik, G., and Ley, S.V. (2010) *Angew. Chem., Int. Ed.*, 49, 6139; (c) Oelke, A.J., Antonietti, F., Bertone, L., Cranwell, P.B., France, D.J., Goss, R.J.M., Hofmann, T., Knauer, S., Moss, S.J., Skelton, P.C., Turner, R.M., Wuitschik, G., and Ley, S.V. (2011) *Chem. Eur. J.*, 17, 4183.
- 193 Baumann, T., Bächle, M., Hartmann, C., and Bräse, S. (2008) *Eur. J. Org. Chem.*, 2207.
- 194 Shigenaga, A., Yamamoto, J., Nishioka, N., and Otaka, A. (2010) *Tetrahedron*, 66, 7367.
- 195 Review: Ni, B. and Headley, A.D. (2010) *Chem. Eur. J.*, 16, 4426.
- 196 (a) Luo, S., Mi, X., Zhang, L., Liu, S., Xu, H., and Cheng, J.-P. (2006) *Angew. Chem., Int. Ed.*, 45, 3093; (b) Luo, S., Mi, X., Liu, S., Xu, H., and Cheng, J.-P. (2006) *Chem. Commun.*, 3687; (c) Luo, S., Mi, X., Zhang, L., Liu, S., Xu, H., and Cheng, J.-P. (2007) *Tetrahedron*, 63, 1923; (d) Zhang, L., Luo, S., Mi, X., Liu, S., Qiao, Y., Xu, H., and Cheng, J. (2008) *Org. Biomol. Chem.*, 6, 567.
- 197 Xu, D., Luo, S., Yue, H., Wang, L., Liu, Y., and Xu, Z. (2006) *Synlett*, 2569.
- 198 Yacob, Z., Shah, J., Leistner, J., and Liebscher, J. (2008) *Synlett*, 2342.
- 199 Li, P., Wang, L., Zhang, Y., and Wang, G. (2008) *Tetrahedron*, 64, 7633.
- 200 Luo, S., Xu, H., Mi, X., Li, J., Zheng, X., and Cheng, J.-P. (2006) *J. Org. Chem.*, 71, 9244.

- 201 (a) Yan, Z.-Y., Niu, Y.-N., Wei, H.-L., Wu, L.-Y., Zhao, Y.-B., and Liang, Y.-M. (2006) *Tetrahedron: Asymmetry*, **17**, 3288; (b) Wu, L.-Y., Yan, Z.-Y., Xie, Y.-X., Niu, Y.-N., and Liang, Y.-M. (2007) *Tetrahedron: Asymmetry*, **18**, 2086.
- 202 Alza, E., Cambeiro, X.C., Jimeno, C., and Pericas, M.A. (2007) *Org. Lett.*, **9**, 3717.
- 203 (a) Chandrasekhar, S., Tiwari, B., Parida, B.B., and Reddy, Ch.R. (2008) *Tetrahedron: Asymmetry*, **19**, 495; (b) Chandrasekhar, S., Mallikarjun, K., Pavankumarreddy, G., Rao, K.V., and Jagadeesh, B. (2009) *Chem. Commun.*, 4985; (c) Chandrasekhar, S., Kumar, T.P., Haribabu, K., and Reddy, C.R. (2010) *Tetrahedron: Asymmetry*, **21**, 2372.
- 204 (a) Miao, T. and Wang, L. (2008) *Tetrahedron Lett.*, **49**, 2173; (b) Miao, T., Wang, L., Li, P., and Yan, J. (2008) *Synthesis*, 3828; (c) Wang, L., Liu, J., Miao, T., Zhou, W., Li, P., Ren, K., and Zhang, X. (2010) *Adv. Synth. Catal.*, **352**, 2571.
- 205 Zhao, Y.-B., Zhang, L.-W., Wu, L.-Y., Zhong, X., Li, R., and Ma, J.-T. (2008) *Tetrahedron: Asymmetry*, **19**, 1352.
- 206 Lv, G., Jin, R., Mai, W., and Gao, L. (2008) *Tetrahedron: Asymmetry*, **19**, 2568.
- 207 Karthikeyan, T. and Sankararaman, S. (2008) *Tetrahedron: Asymmetry*, **19**, 2741.
- 208 Zhang, L., Cui, L., Li, X., Li, J., Luo, S., and Cheng, J.-P. (2010) *Eur. J. Org. Chem.*, 4876.
- 209 Wang, W., Wang, J., Li, H., and Liao, L. (2004) *Tetrahedron Lett.*, **45**, 7235.
- 210 Wang, W., Wang, J., and Li, H. (2004) *Tetrahedron Lett.*, **45**, 7243.
- 211 Wang, W., Li, H., Wang, J., and Liao, L. (2004) *Tetrahedron Lett.*, **45**, 8229.
- 212 (a) Wang, W., Wang, J., and Li, H. (2005) *Angew. Chem., Int. Ed.*, **44**, 1369; (b) Wang, J., Li, H., Zu, L., and Wang, W. (2006) *Adv. Synth. Catal.*, **348**, 425; (c) Zu, L., Wang, J., Li, H., and Wang, W. (2006) *Org. Lett.*, **8**, 3077; (d) Wang, J., Li, H., Lou, B., Zu, L., Guo, H., and Wang, W. (2006) *Chem. Eur. J.*, **12**, 4321; (e) Wang, J., Zhang, M.-M., Zhang, S., Xu, Z.-A., Li, H., Yu, X.-H., and Wang, W. (2011) *Synlett*, 473.
- 213 (a) Wang, W., Li, H., and Wang, J. (2005) *Tetrahedron Lett.*, **46**, 5077; (b) Zu, L., Xie, H., Li, H., Wang, J., and Wang, W. (2008) *Org. Lett.*, **10**, 1211; (c) Mei, K., Zhang, S., He, S., Li, P., Jin, M., Xue, F., Luo, G., Zhang, H., Song, L., Duan, W., and Wang, W. (2008) *Tetrahedron Lett.*, **49**, 2681.
- 214 Enders, D. and Chow, S. (2006) *Eur. J. Org. Chem.*, 4578.
- 215 Diez, D., Gil, M.J., Moro, R.F., Marcos, I.S., García, P., Basabe, P., Garrido, N.M., Broughton, H.B., and Urones, J.G. (2007) *Tetrahedron*, **63**, 740.
- 216 (a) Ni, B., Zhang, Q., and Headley, A.D. (2007) *Green Chem.*, **9**, 737; (b) Ni, B., Zhang, Q., and Headley, A.D. (2007) *Tetrahedron: Asymmetry*, **18**, 1443; (c) Ni, B., Zhang, Q., Dhungana, K., and Headley, A.D. (2009) *Org. Lett.*, **11**, 1037.
- 217 Cao, C.-L., Sun, X.-L., Kang, Y.-B., and Tang, Y. (2007) *Org. Lett.*, **9**, 4151.
- 218 Lu, A., Wu, R., Wang, Y., Zhou, Z., Wu, G., Fang, J., and Tang, C. (2010) *Eur. J. Org. Chem.*, 2057.
- 219 (a) Ban, S., Du, D.-M., Liu, H., and Yang, W. (2010) *Eur. J. Org. Chem.*, 5160; (b) Luo, C. and Du, D.-M. (2011) *Synthesis*, 1968.
- 220 Dahlin, N., Bøgevig, A., and Adolfsson, H. (2004) *Adv. Synth. Catal.*, **346**, 1101.
- 221 (a) Liu, P.-M., Chang, C., Reddy, R.J., Ting, Y.-F., Kuan, H.-H., and Chen, K. (2010) *Eur. J. Org. Chem.*, 42; (b) Liu, P.-M., Magar, D.R., and Chen, K. (2010) *Eur. J. Org. Chem.*, 5705.
- 222 Review: Lu, L.-Q., An, X.-L., Chen, J.-R., and Xiao, W.-J. (2012) *Synlett*, 490.
- 223 (a) Cao, C.-L., Ye, M.-C., Sun, X.-L., and Tang, Y. (2006) *Org. Lett.*, **8**, 2901; (b) Cao, C.-L., Sun, X.-L., Zhou, J.-L., and Tang, Y. (2007) *J. Org. Chem.*, **72**, 4073; (c) Cao, C.-L., Zhou, Y.-Y., Zhou, J., Sun, X.-L., Tang, Y., Li, Y.-X., Li, G.-Y., and Sun, J. (2009) *Chem. Eur. J.*, **15**, 11384; (d) Cao, X.-Y., Zheng, J.-C., Li, Y.-X., Shu, Z.-C., Sun, X.-L., Wang, B.-Q., and Tang, Y. (2010) *Tetrahedron*, **66**, 9703.
- 224 (a) Cao, Y.-J., Lu, H.-H., Lai, Y.-Y., Lu, L.-Q., and Xiao, W.-J. (2006) *Synthesis*, 3795; (b) Cao, Y.-J., Lai, Y.-Y., Wang, X., Li, Y.-J., and Xiao, W.-J. (2007) *Tetrahedron Lett.*, **48**, 21; (c) Chen, J.-R., An, X.-L., Zhu, X.-Y., Wang, X.-F., and Xiao, W.-J. (2008) *J. Org. Chem.*, **73**,

- 6006; (d) Chen, J.-R., Cao, Y.-J., Zou, Y.-Q., Tan, F., Fu, L., Zhu, X.-Y., and Xiao, W.-J. (2010) *Org. Biomol. Chem.*, **8**, 1275.
- 225 Shen, Z., Zhang, Y., Jiao, C., Li, B., Ding, J., and Zhang, Y. (2007) *Chirality*, **19**, 307.
- 226 Carley, A.P., Dixon, S., and Kilburn, J.D. (2009) *Synthesis*, 2509.
- 227 Lu, A., Gao, P., Wu, Y., Wang, Y., Zhou, Z., and Tang, C. (2009) *Org. Biomol. Chem.*, **7**, 3141.
- 228 Chen, J.-R., Fu, L., Zou, Y.-Q., Chang, N.-J., Rong, J., and Xiao, W.-J. (2011) *Org. Biomol. Chem.*, **9**, 5280.
- 229 Lin, J., Tian, H., Jiang, Y.-J., Huang, W.-B., Zheng, L.-Y., and Zhang, S.-Q. (2011) *Tetrahedron: Asymmetry*, **22**, 1434.
- 230 Yu, C., Qiu, J., Zheng, F., and Zhong, W. (2011) *Tetrahedron Lett.*, **52**, 3298.
- 231 Kokotos, C.G., Limnios, D., Triggidou, D., Trifonidou, M., and Kokotos, G. (2011) *Org. Biomol. Chem.*, **9**, 3386.
- 232 Pansare, S.V. and Lingampally, R. (2009) *Org. Biomol. Chem.*, **7**, 319.
- 233 Riguert, E. (2009) *Tetrahedron Lett.*, **50**, 4283.
- 234 Morokuma, K., Taira, Y., Uehara, Y., Shibahara, S., Takahashi, K., Ishihara, J., and Hatakeyama, S. (2008) *Tetrahedron Lett.*, **49**, 6043.
- 235 (a) Li, J., Yang, G., Qin, Y., Yang, X., and Cui, Y. (2011) *Tetrahedron: Asymmetry*, **22**, 613; (b) Li, J., Yang, G., and Cui, Y. (2011) *J. Appl. Polym. Sci.*, **121**, 1506.
- 236 Wang, L., Cai, C., Curran, D.P., and Zhang, W. (2010) *Synlett*, 433.
- 237 Chuan, Y.-M., Chen, G.-H., Gao, J.-Z., Zhang, H., and Peng, Y.-G. (2011) *Chem. Commun.*, **47**, 3260.
- 238 (a) Burroughs, L., Vale, M.E., Gilks, J.A.R., Forintos, H., Hayes, C.J., and Clarke, P.A. (2010) *Chem. Commun.*, **46**, 4776; (b) Burroughs, L., Clarke, P.A., Forintos, H., Gilks, J.A.R., Hayes, C.J., Vale, M.E., Wade, W., and Zbytniewski, M. (2012) *Org. Biomol. Chem.*, **10**, 1565.
- 239 Xiao, J., Lu, Y., Liu, Y.-L., Wong, P.-S., and Loh, T. (2011) *Org. Lett.*, **13**, 876.
- 240 Melchiorre, P. and Jørgensen, K.A. (2003) *J. Org. Chem.*, **68**, 4151.
- 241 Ishii, T., Fujioka, S., Sekiguchi, Y., and Kotsuki, H. (2004) *J. Am. Chem. Soc.*, **126**, 9558.
- 242 (a) Xu, D.-Q., Yue, H.-D., Luo, S.-P., Xia, A.-B., Zhang, S., and Xu, Z.-Y. (2008) *Org. Biomol. Chem.*, **6**, 2054; (b) Xu, D.-Q., Wang, L.-P., Luo, S.-P., Wang, Y.-F., Zhang, S., and Xu, Z.-Y. (2008) *Eur. J. Org. Chem.*, 1049; (c) Xu, D.-Q., Xia, A.-B., Luo, S., Tang, J., Zhang, S., Jiang, J.-R., and Xu, Z.-Y. (2009) *Angew. Chem., Int. Ed.*, **48**, 3821.
- 243 Mandal, T. and Zhao, C.-G. (2007) *Tetrahedron Lett.*, **48**, 5803.
- 244 (a) Díez, D., Antón, A.B., García, P., Garrido, N.M., Marcos, I.S., Basabe, P., and Urones, J.G. (2008) *Tetrahedron: Asymmetry*, **19**, 2088; (b) Díez, D., Antón, A.B., Pena, J., García, P., Garrido, N.M., Sanz, M.F., Basabe, P., and Urones, J.G. (2010) *Tetrahedron: Asymmetry*, **21**, 786.
- 245 (a) Liu, F., Wang, S., Wang, N., and Peng, Y. (2007) *Synlett*, 2415; (b) Wang, C., Yu, C., Liu, C., and Peng, Y. (2009) *Tetrahedron Lett.*, **50**, 2363; (c) Chuan, Y.-M., Yin, L.-Y., Zhang, Y.-M., and Peng, Y.-G. (2011) *Eur. J. Org. Chem.*, 578.
- 246 Zeng, X. and Zhong, G. (2009) *Synthesis*, 1545.
- 247 Syu, S., Kao, T.-T., and Lin, W. (2010) *Tetrahedron*, **66**, 891.
- 248 Bauer, J.O., Stiller, J., Marqués-López, E., Strohfeldt, K., Christmann, M., and Strohmman, C. (2010) *Chem. Eur. J.*, **16**, 12553.
- 249 Liu, X.-W., Le, T.N., Lu, Y., Xiao, Y., Ma, J., and Li, X. (2008) *Org. Biomol. Chem.*, **6**, 3997.
- 250 Zeng, Z., Luo, P., Jiang, Y., Liu, Y., Tang, G., Xu, P., Zhao, Y., and Blackburn, G.M. (2011) *Org. Biomol. Chem.*, **9**, 6973.
- 251 Lacoste, E., Landais, Y., Schenk, K., Verlhac, J.-B., and Vincent, J.-M. (2004) *Tetrahedron Lett.*, **45**, 8035.
- 252 Frisch, K., Landa, A., Saaby, S., and Jørgensen, K.A. (2005) *Angew. Chem., Int. Ed.*, **44**, 6058.
- 253 Kano, T., Hato, Y., and Maruoka, K. (2006) *Tetrahedron Lett.*, **47**, 8467.
- 254 Díez, D., Núñez, M.G., Benítez, A., Moro, R.F., Marcos, I.S., Basabe, P.,

- Broughton, H.B., and Urones, J.G. (2009) *Synlett*, 390.
- 255 Juhl, K. and Jørgensen, K.A. (2003) *Angew. Chem., Int. Ed.*, **42**, 1498.
- 256 Samanta, S., Krause, J., Mandal, T., and Zhao, C.-G. (2007) *Org. Lett.*, **9**, 2745.
- 257 (a) Halland, N., Braunton, A., Bachmann, S., Marigo, M., and Jørgensen, K.A. (2004) *J. Am. Chem. Soc.*, **126**, 4790; (b) Halland, N., Lie, M.A., Kjærsgaard, A., Marigo, M., Schiøtt, B., and Jørgensen, K.A. (2005) *Chem. Eur. J.*, **11**, 7083.
- 258 (a) Marigo, M., Fielenbach, D., Braunton, A., Kjærsgaard, A., and Jørgensen, K.A. (2005) *Angew. Chem., Int. Ed.*, **44**, 3703; (b) Bertelsen, S., Halland, N., Bachmann, S., Marigo, M., Braunton, A., and Jørgensen, K.A. (2005) *Chem. Commun.*, 4821.
- 259 Fadeyi, O.O., Schulte, M.L., and Lindsley, C.W. (2010) *Org. Lett.*, **12**, 3276.
- 260 Umezawa, T., Shibata, M., Kaneko, K., Okino, T., and Matsuda, F. (2011) *Org. Lett.*, **13**, 904.
- 261 (a) Aggarwal, V.K., Lopin, C., and Sandrinelli, F. (2003) *J. Am. Chem. Soc.*, **125**, 7596; (b) Aggarwal, V.K. and Fang, G.Y. (2005) *Chem. Commun.*, 3448.
- 262 Kano, T., Mii, H., and Maruoka, K. (2009) *J. Am. Chem. Soc.*, **131**, 3450.
- 263 O'Hagan, D., Royer, F., and Tavasli, M. (2000) *Tetrahedron: Asymmetry*, **11**, 2033.
- 264 Ho, C.-Y., Chen, Y.-C., Wong, M.-K., and Yang, D. (2005) *J. Org. Chem.*, **70**, 898.
- 265 (a) Sparr, C., Schweizer, W.B., Senn, H.M., and Gilmour, R. (2009) *Angew. Chem., Int. Ed.*, **48**, 3065; (b) Sparr, C., Tanzer, E.-M., Bachmann, J., and Gilmour, R. (2010) *Synthesis*, 1394.
- 266 Shi, Z., Tan, B., Yi Leong, W.W., Zeng, X., Lu, M., and Zhong, G. (2010) *Org. Lett.*, **12**, 5402.

