

## Index

### a

- absolute adsorption 600
- acetylacetonate 107
- 4,4',4'',4'''-(adamantane-1,3,5,7-tetrayl)tetrabenzoic acid (H<sub>4</sub>ATB) 293
- adsorption
  - enthalpy
    - calorimeter 591
    - estimations 591
    - ligand functionalization 595
    - probing open metal sites 592
    - structure transitions 596
  - high pressure adsorption
    - absolute adsorption 600
    - Gibbs excess adsorption isotherms 600
    - gravimetric technique 598
    - manometric method 598
    - surface excess 598
    - total adsorption 600
  - molecular simulation
    - adsorbent/adsorbate interactions 774
    - extended charge equilibration PQeq strategy 774
    - force field 774
    - GCMC, *see* grand canonical Monte Carlo (GCMC) simulations
    - LJ parameters 774
    - Monte Carlo simulations 773, 774
    - rationalization 780
    - and NPD
      - redox processes 713
      - adsorption chamber 709
      - Co<sub>2</sub>(*m*-dobdc) 711, 712
      - Co<sub>2</sub>(dobdc) 711
      - Cu<sub>3</sub>(btc)<sub>2</sub> 710
      - Fourier map analysis 710
      - hydrogen and methane adsorption 709
      - neutron scattering properties 709
      - NOTT-101 material 712
      - research reactors 709
      - spallation sources 709
      - X-rays scattering 709
  - porosity, *see* porosity determination
  - and powder XRD
    - acetylene 700
    - Co(BDP) 704
    - CO<sub>2</sub> and CH<sub>4</sub> mixtures 702
    - Co<sub>2</sub>(dobdc) 708
    - CPL-1 697, 699
    - DUT-8(Ni) 705
    - ELM-11 700
    - large pore (lp) 702
    - methanol adsorption isotherm 699
    - micropores 699
    - MIL-53(Cr) 699, 701
    - MIL-53(Fe)-X 703
    - narrow pore (np) 702
    - NOTT-300 707
    - rigid/flexible MOFs, pores of 709
    - single crystals 708
    - structural transformation 708
    - synchrotron radiation 699
    - 3<sup>rd</sup> generation PCPs 698
  - sample preparation 576
- Ag(TCNQ) 448
- Al<sub>2</sub>(OH)<sub>2</sub>(H<sub>2</sub>TCPP)] (Al-PMOF) 117
- alkaline-earth metal based MOFs (AE-MOFs)
  - advantages 73, 93
  - linkers
    - carboxylate-based linkers 75
    - participating linkers 75
    - phosphonate and sulfonated linkers 75
    - role 75
  - properties and applications 86
  - catalysis 93
  - gas-adsorption and separation 86, 89

- alkaline-earth metal based MOFs (AE-MOFs) (*contd.*)
- luminescence-based chemical sensing 94
  - proton and ionic conductivity 96, 97
  - selected physical 87
  - structures
    - barium 84
    - beryllium 76
    - calcium 80
    - factors 85
    - magnesium 77
    - organic linker(s), shape and size of 85
    - strontium 82
    - synthetic parameters 85
    - synthesis
      - hydrothermal 74
      - ionic liquids 75
      - ionothermal 74
      - solvothermal 74
  - aluminium
    - annual production 105
    - low toxicity 105
    - properties 106
    - structures 119
    - synthesis 119
  - aluminium fumarate 126, 825
  - amino-MOFs 128, 129, 139, 347, 351, 367, 372, 388–396, 412, 415, 437, 503, 534, 543, 623, 662, 743
  - 3-aminopropyltrialkoxysilane (APS) 361
  - angular-dispersive diffraction (ADXRD) 732, 733
  - atomic force microscopy (AFM) 536, 754
  - atomic layer deposition (ALD) 166
  - atropisomery 402
  - automated assembly of structure building unit (AASBU) approach 771
  - Avrami-Erofe'ev model 735, 742
  - azolates 309, 310
- b**
- Basolite A520 825
  - 1,4-benzenedicarboxylate (BDC) 139
  - benzenedicarboxylic acid 115
  - 5,5',5'',5''',5''''*sit*,5''''''-(benzene-1,2,3,4,5,6-hexylhexakis(methylene)hexakis(oxy))-hexaisophthalic acid (H<sub>12</sub>PHHI) 303
  - 1,2,4,5-benzenetetracarboxylic acid (H4btcc) 94
  - 1,3,5-benzenetribenzoic acid (btb) 76
  - 4,4',4''-benzene-1,3,5-triyl-tribenzoic acid (H3BTB) 291
  - 5,5',5''-((benzene-1,3,5-tricarbonyl)tris(azanediyl))triisophthalic acid (H<sub>6</sub>TPBTM) 299
  - 4,4',4''-(Benzene-1,3,5-triyl-tris(benzene-4,1-diyl))tribenzoic acid (H<sub>3</sub>BBC) 291
  - 5,5',5''-(((benzene-1,3,5-triyltris(benzene-4,1-diyl))tris(ethyne-2,1-diyl))-tris(benzene-4,1-diyl))tris(ethyne-2,1-diyl))triisophthalic acid (H<sub>6</sub>BHEHPI) 300
  - 5,5',5''-(benzene-1,3,5-triyltris(buta-1,3-diyne-4,1-diyl))triisophthalic acid (H<sub>6</sub>BHEI) 300
  - 4,4',4''-(benzene-1,3,5-triyltris(ethyne-2,1-diyl))tribenzoic acid (H<sub>3</sub>BTE) 291
  - 5,5',5''-(((benzene-1,3,5-triyl-tris(ethyne-2,1-diyl))tris(benzene-4,1-diyl))tris(ethyne-2,1-diyl))triisophthalic acid (H<sub>6</sub>TTEI) 300
  - 6,6',6''-(benzene-1,3,5-triyl)tris(2-naphthoic acid) (H<sub>3</sub>BTN) 292
  - 3,4,3',4',3'',4''-benzo[18]annulenehexabenzoc acid (H<sub>6</sub>PhTHBA) 301
  - benzoic acid 163
  - beryllium 86
  - 9,9'-([1,1'-biphenyl]-4,4'-diyl)bis(9H-carbazole-3,6-dicarboxylic acid) (H<sub>4</sub>BBCDC) 297
  - 9,9',9'',9'''-([1,1'-biphenyl]-3,3',5,5'-tetrayl)tetrakis(9H-carbazole-3,6-dicarboxylic acid) (H<sub>8</sub>BTTCD) 302
  - 1,12-bis(3',5'-bis(hydroxycarbonyl)phen-1'-yl)-1,12-dicarba-closo-dodecaborane (H<sub>4</sub>CBI) 295
  - 5',5''-bis(4-carboxyphenyl)-2',2''-diethoxy-[1,1'3',1''3'',1'''-quaterphenyl]-4,4'''-dicarboxylic acid (H<sub>4</sub>CQDA-OEt) 296
  - 5,5''-((2,2-bis((3,5-dicarboxyphenoxy)methyl)propane-1,3-diyl)bis(oxy))diisophthalic acid (H<sub>8</sub>TDM) 302
  - 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl (BINAP) 405
  - bismuth tri- and tetraarylcarboxylates 744
  - “bottom up” approach 771
  - Brønsted acidic ionic liquid (BAIL) 650
  - Bragg reflectors 540
  - Bragg stacks (BSs) 540
  - Brunauer–Emmett–Teller (BET) surface area 49, 159
- c**
- carbon nanoparticles 494
  - carboxylate-based linkers 75
  - carboxylic functional groups (CFGs) 167

- 5'-(4-carboxyphenyl)-[1,1'3',1''-terphenyl]-3,3'',5,5''-tetracarboxylic acid (H<sub>5</sub>PTPCA) 298
- chain-based pillared-column frameworks 324
- charge-coupled detector (CCD) 745
- charge-density wave (CDW) 206, 436–438
- china lantern' complex 438
- and chromatography
- – analytical separations 410
- chiral linker systems
- amino acids
  - – azides with alkynes 391
  - – chir-H2bpdc 389
  - – chiral pool 388
  - – cyclization reaction 390
  - – electrophilic properties 391
  - – enantiopure derivatives 391
  - – naphthalene tetracarboxylic acid 390
  - – natural  $\alpha$ -amino acids 388
  - – peptides 389, 392
  - – Schiff base derivatives 391
  - – single  $\alpha$ -amino acids 388
  - axial chirality
  - – arylboronic ester 402
  - – atoms 401
  - – atropisomery 402
  - – BINAP 406
  - – BINOL 402, 405, 408
  - – biphenyl- and binaphthyl-derivates 401
  - – bromine groups 405
  - – calixarenes 408
  - – chiral pool 401
  - – CMOF-4b 404
  - – enantiopurification 401
  - – isophthalic acid derivative 404
  - – multidentate linkers 405
  - – rigid backbone 407
  - – rotamers 402
  - – tetra-bromide precursor 404
  - – tetracarboxylic acid 407
  - central chirality 401
  - enantioselective separation and chromatography
  - – analytical separations 409, 410
  - – GC 410
  - – HPLC 410
  - – preparative separations 409, 413
  - – simulation 414
  - natural sources 396
  - rigid linkers 397
  - salen linkers 399
- chiral separation 524
- chirality 387
- clays 105
- cluster-based networks 318
- CMOFs 359
- 3-c net 9
- Co-por-2 ligand 357
- Co<sub>4</sub>O(pz)<sub>6</sub> clusters 322
- commercially available MOF 825
- confocal fluorescence microscopy (CFM) 811
- continuous wave (CW) 629
- coordination network compounds (CNCs) 795
- coordination sequence 24, 25
- Copper(II)-benzene-1,3,5-tricarboxylate (Cu-BTC) 554
- CPM-99 159
- cross polarization (CP) 609
- crystalline nanoparticles 495
- crystallisation
- atomic force microscopy 754
  - crystalline intermediate phases 759
  - defects, MOF 796
  - electrospray ionisation mass spectrometry 758
  - HKUST-1 757
  - in situ diffraction
  - – ADXRD 733
  - – CAU-1-(NH<sub>2</sub>) 743
  - – CAU-1-(OH)<sub>2</sub> 743
  - – diffraction data measurement 731
  - – EDXRD, *see* energy-dispersive diffraction (EDXRD)
  - – kinetics 734
  - – MOF-74 739
  - – monochromatic X-rays 745
  - – SAXS 734
  - – SAXS/WAXS 746
  - large image-plate detectors 759
  - light scattering 749
  - MIL-101 760
  - MOF, solvothermal synthesis of 731
  - multi-technique analyses 759
  - NMR spectroscopy 751
  - PDF 760
  - polycrystalline powders 730
  - Raman scattering 760
  - SAMs method 757
  - small-angle neutron scattering 760
  - small-angle X-ray scattering 731
  - solvothermal conditions 729
  - UiO-66 759
  - vibrational spectroscopy 750

- crystallisation (*contd.*)
- XAFS 753
  - zeolite 730
- Cu-BTC (HKUST-1) pellet 567
- cuboctahedral (cuo) 61, 64
- cyclohexane 355
- d**
- deconstruction of MOFs 8
- defect-linker (DL) 796, 807
- density functional theory (DFT) 159, 587, 668
- DFT/GCMC method 588
- 4,4',4'',4'''-((1E,1'E,1''E,1'''E)-((R)-2,2'-diethoxy-[1,1'-binaphthalene]-4,4',6,6'-tetrayl)tetrakis(ethene-2,1-diyl))tetrabenzoic acid (H<sub>4</sub>BNETBA-OEt) 297
- diethylenetriamine (DETA) 361
- diethylformamide (DEF) 359, 471
- diffuse reflectance infrared fourier transform spectroscopy (DRIFTS) 166, 664
- N,N-diisopropylformamide (DIF) 78
- dimethylformamide (DMF) 146, 471
- 4,4'-((1E,1'E)-(2,5-dimethoxy-1,4-phenylene)bis(ethene-2,1-diyl))dibenzoic acid (H<sub>2</sub>PVDC) 289
- 4,4'-((2,5-dimethoxy-1,4-phenylene)bis(ethyne-2,1-diyl))dibenzoic acid (H<sub>2</sub>DMBDA) 289
- 1,4-dioxido-2, 5-benzenedicarboxylate (DOBDC) 144
- dithieno[3,2-b + 2',3'-d]thiophene-2,6-dicarboxylate (DTTDC) linker 163
- ditopic linkers 288
- H<sub>2</sub>CONQDA 288
  - H<sub>2</sub>DMBDA 289
  - H<sub>2</sub>HHTFDA 288
  - H<sub>2</sub>POMDA 289
  - H<sub>2</sub>PVDC 289
  - H<sub>4</sub>DH<sub>1</sub>PhDC 288
- DMOF, basic-hydrophobic balance 365
- dodecatopic linkers
- H<sub>12</sub>PHHI 303
  - H<sub>12</sub>TDDPB 303
- DUT-6 278, 291, 470
- DUT-11 32
- DUT-32 614
- DUT-49 65, 66, 279, 297
- DUT-67(Zr) (Dresden University of Technology-67) 832
- dye sensitized solar cells (DSSCs) 179
- dynamic nuclear polarization (DNP) 377, 610
- e**
- edge-transitive nets 27
- electron nuclear double resonance (ENDOR) spectroscopy 629
- electron paramagnetic resonance (EPR) spectroscopy
- crystalline powders 635
  - CW EPR 629
  - ENDOR 629
  - ESEEM 629
  - ESEEM 636
  - free radicals, linker based 651
  - inorganic radicals (NO, O<sub>2</sub>-) 652
  - magnetic resonance condition 629
  - MOF, *see* metal organic framework (MOF)
  - nitroxide radicals 651
  - paramagnetic species 629
  - principles of 630
  - pulsed ENDOR spectroscopy 637
  - S = 1/2 and I = 1/2 model system 636
  - Schrödinger equation 631
  - spin Hamiltonian formalism
    - – electron Zeeman interaction 631
    - – hyperfine interaction 633
    - – nuclear quadrupole interaction 635
    - – nuclear Zeeman interaction 632
    - – phenomenological second-rank tensors 631
    - – zero-field splitting 634
  - TMI
    - – Co<sup>2+</sup> 650
    - – Cr<sup>2+</sup> 648
    - – Cu<sup>2+</sup> 644
    - – paramagnetic 3d 650
    - – Ti<sup>3+</sup> and V<sup>4+</sup> 646
  - X-band experiments 629
- electron spin echo envelope modulation (ESEEM) 629, 636
- electron spin resonance (ESR) spectroscopy 629
- electron Zeeman interaction 630, 631
- energy dispersive X-ray spectroscopy (EDS) 812
- energy-dispersive diffraction (EDXRD)
- advantages 732
  - bismuth tri- and tetraarylcarboxylates 744
  - Bragg equation 732
  - CAU-n family 742
  - chalcogenide materials 733
  - Cu(II) phosphonatoethanesulfonates 742
  - Daresbury SRS 733
  - DESY facility 733
  - disadvantage 732
  - HKUST-1 736

- lanthanide phosphonatobutanesulfonates 742
  - MIL-53(Fe) 737
  - MOF-14 739
  - neb phase 742
  - UiO-66 744
  - ultrasound, implementation of 744
  - zeolite 733
  - ZIF-8 740
  - zirconium fumarate MOF 741
  - ESEEM spectroscopy, *see* electron spin echo envelope modulation (ESEEM)
  - 5,5',5'',5'''-(ethene-1,1,2,2-tetrayltetrakis (benzene-4,1-diyl))tetrakis(ethyne-2,1-diyl)tetrakisophthalic acid (H<sub>8</sub>TDPEPE) 302
  - 4',4''',4''''',4''''''|-(ethene-1,1,2,2-tetrayl)tetrakis((1,1'-biphenyl]-4-carboxylic acid)) (H<sub>4</sub>ETTC) 296
  - ethylene diamine (ED) 361
  - European Synchrotron Radiation Facility (ESRF) 746
  - ex situ* methods 691
  - extended organic linkers
    - binding sites 285
    - construction 283
    - core unit 283
    - designing linker synthesis 280
    - ditopic 285
    - extending units 283
    - hexatopic 298
    - octatopic 301, 302
    - pentatopic 298
    - tetratopic 293
    - tritopic 289
  - extended X-ray absorption fine structure (EXAFS) 714
    - clusters 715
    - CPO-27(Ni) 715
    - crystallisation 753
    - Cu atoms 714
    - defects, MOF 812
    - DUT-8(Ni) 717
    - fit parameters 720
    - HKUST-1 716
    - K<sub>2</sub>PtCl<sub>4</sub> and UiO-66 - NH<sub>2</sub> 721
    - O<sub>2</sub>-loaded, pre-edge data 720
    - parameter values 723
    - post-activation data 723
  - extrusion
    - advantages 562
    - Caleva Mini Screw Extruder 563
    - Cu-BTC 564
    - extrudates 562
    - MOF monoliths 563
    - process 563
    - STYs 563, 565
    - synthetic approach 563
    - twin screw extruder 565
- f**
- face symbols 18
  - ferrocenylamide 129
  - finite graphs 8
  - flash-photolysis time-resolved microwave conductivity (FP-TRMC) technique 424, 426
  - 2-fold symmetric M(pz)<sub>2</sub> chains 328
  - 4-fold symmetric M(pz)<sub>2</sub> chains 328
  - 6-fold symmetric M(pz)<sub>3</sub> chains 329
  - four-probe method 422, 423, 437
  - framework breathing 332
  - freeze drying (FD) 577
  - functional linkers
    - MOFs 345
    - design, characterizations 376
    - post-synthetic modification
      - metalation 366
      - organocatalyst grafting 361
    - self-assembled frameworks
      - metal functionalized ligands 352
      - organocatalyst ligands 347
  - functionalized
    - dihydroxy-undeciphenyldicarboxylic acid (H<sub>4</sub>DH<sub>11</sub>PhDC) 288
- g**
- Galinstan™ 105
  - gallium (Ga)
    - annual production 105
    - framework structures 121
    - in semiconductor technology 105
    - low toxicity 105
    - properties of 106
  - gallium arsenide 105
  - gas chromatography (GC) 410, 413
  - Gd<sup>3+</sup> containing nanoMOFs 262
  - genus 22
  - Gibbs excess adsorption isotherms 600
  - grand canonical Monte Carlo (GCMC) simulations 581, 589
    - adsorption isotherm 778, 780
    - global theoretical isotherm 780
    - *in situ* IR experiments 780
    - MIL-47(V) [115] 780
    - MIL-53(Cr) 777
    - prediction 775

- granulation
  - CPO-27(Ni) spheres 558
  - Cu-BTC (HKUST-1) spheres 556
  - formulation parameters and process equipment 558
  - hydrocolloids 557
  - mechanical strength 553
  - MIL-100(Fe) spheres 556, 557
  - parameters and relative interactions 554
  - physicochemical properties 554, 559
  - process 553, 554
  - products, with different diameters 558
  - properties 554
  - self-supporting and self-completing process 557
  - transformation 554
  - VSSA 557
- graphics processing units (GPUs) 775
- graphs 7, 8
- group 13 elements (G13-MOFs)
  - aluminium
    - – frame work structures 119, 120
    - – MIL-110 121
    - – properties 106
    - – synthesis 119
  - application 125
  - breathing process 127
  - characterisation 110
  - distribution 106
  - gallium
    - – framework structures 121
    - – properties 106
  - vs. IBUs 107
  - indium
    - – properties 106
    - – structures 122
  - inorganic building units 110
  - post-synthetic modification 128
  - structures 110, 116
  - synthesis 109
    - – Al-CAU-13 126
    - – Al-Fum 126
    - – Al-MIL-53 126
- group 3 MOFs, *see* Lanthanide containing MOFs (Ln-MOFs)
- group 4 metals-based MOFs 166
  - chemical stability 138
  - hafnium 137
  - titanium, *see* titanium-based MOFs
  - valence of 138
  - zirconium, *see* zirconium-based MOFs
- Gualtieri equation 736
- h**
  - Heck reaction 285
  - 4,4',4''-(1,3,3a1,4,6,7,9-heptaazaphenalene-2,5,8-triyl)tribenzoic acid (H<sub>3</sub>HTB) 291
  - 4,4'-(9,9,9',9'',9'',9''-hexahexyl-9H,9'H,9''H-[2,2''7',2''-terfluorene]-7,7''-diyl)-dibenzoic acid (H<sub>2</sub>HHTFDA),288
  - hexatopic linkers 298
    - H<sub>6</sub>TDCPB 300
    - H<sub>6</sub>BHEHPI 300
    - H<sub>6</sub>BHEI 300
    - H<sub>6</sub>PhTHBA 301
    - H<sub>6</sub>TTA 299
    - H<sub>6</sub>TPBTM 299
    - H<sub>6</sub>TTEI 300
  - high resolution neutron powder diffraction (NPD) experiments 672
  - high-performance liquid adsorption (HPLC) 410
  - HKUST-1 (Hong Kong University of Science and Technology-1) 345, 829
  - <sup>1</sup>H NMR spectra 350, 364, 376
  - Horvath-Kawazoe (HK) method 587
  - hybrid osmotic Monte Carlo (HOMC) scheme 777
  - hydrocolloids 557
  - hydrogen-bond-donating (HBD) 349
  - hydrolytic kinetic resolution (HKR) 360
  - hyperfine sublevel correlation (HYSCORE) 379, 636
  - HYSCORE-EPR technique 380
- i**
  - ideal adsorbed solution theory (IAST) 669
  - imidazolate coordination modes 311
    - polyimidazolate 315
    - zeolitic and zeolite-like frameworks 311
  - 1H-imidazo[4,5-f][1,10]phenanthroline (Hip) 337
  - in situ diffraction
    - adsorption, *see* adsorption
    - ADXRD 733
    - CAU-1-(NH<sub>2</sub>) 743
    - CAU-1-(OH)<sub>2</sub> 743
    - diffraction data measurement 731
    - EDXRD
      - – advantages 732
      - – bismuth tri- and tetraarylcarboxylates 744
      - – Bragg equation 732
      - – CAU-n family 742
      - – chalcogenide materials 733

- – Cu(II) phosphonatoethanesulfonates 742
- – Daresbury SRS 733
- – DESY facility 733
- – disadvantage 732
- – HKUST-1 736
- – lanthanide phosphonatobutanesulfonates 742
- – MIL-53(Fe) 737
- – MOF-14 739
- – neb phase 742
- – UiO-66 744
- – ultrasound, implementation of 744
- – zeolite 733
- – ZIF-8 740
- – zirconium fumarate MOF 741
- instrumentation
  - – adsorption chamber 692
  - – adsorption measurements 697
  - – beryllium chamber 696
  - – capillary based cells 693
  - – close cycle cryocoolers 698
  - – diffractometer and experimental set-up 693, 694
  - – gas flow cryostream systems 693
  - – glass/quartz capillary 693
  - – phase transformation 696
  - – powder XRD and gas adsorption 697
  - – sample holder 694, 696
  - – SNBL 693
  - – temperature control 696
  - – temperature profiling 695
  - – kinetics 734
  - MOF-74 739
  - monochromatic X-rays 745
  - SAXS 734
  - SAXS/WAXS 746
- in situ* method 691
- indium
  - annual production 105
  - low toxicity 105
  - properties of 106
  - structural properties 122
- indium tin oxide (ITO) 105
- inductively coupled plasma-mass spectrometry (ICP-MS) 372
- infinite graphs 8
- infrared reflection-absorption microscopy (IRRAS) 536
- infrared spectroscopy 377
- inherent defects 796
- inorganic building units (IBUs) 105
- interfacial crystal growth method 330
- internal defects 806
- ion and liquid assisted grinding (ILAG) 745
- Ir-based catalyst, solid nitrogen-containing chelating ligand 372, 373
- IRMOF-3 367, 368
- k**
  - Kelvin equation methods 587
  - kinetically tuned dimensional augmentation 140
- l**
  - large-scale defects 810
  - linkers
    - carboxylates
      - – applications 424
      - – Cu<sub>3</sub> (BTC)<sub>2</sub> film 430
      - – FP-TRMC 427
      - – M<sub>2</sub>(DOBDC) 426, 427
      - – Mn<sub>2</sub>(DSBDC) 426
      - – MOF-5 424
      - – photocurrent transients 427
      - – pores 428
      - – structure design/doping technique 426
      - – TCNQ 430
      - – “through-bond” approach 426
      - – “through-space” approach 426
    - catechol
      - – Co-CAT-1 442
      - –  $\pi$ -conjugated nickel bis(dithiolene) complex nanosheet 445
      - – 2D conducting materials 443, 444
      - – derivatives 442
      - – diradical Ni-bisdiimine linkages 444
      - – high conductivity 442
      - – molecular formula 442
      - – nickel bis(dithiolene) complex nanosheet 444
      - – Pt-HTT 446
    - conductivity data 457
    - Cu[Cu(pdt)<sub>2</sub>] 435
    - cyanide
      - – PB 453
      - – PBAs 454, 456
      - – Rb<sub>0.5</sub>Co<sup>III</sup>Co<sup>II</sup><sub>0.25</sub>[Fe<sup>II</sup>(CN)<sub>6</sub>].5.9H<sub>2</sub>O 456
      - – Ru<sup>III</sup>Ru<sup>II</sup> 455
    - electric current, time dependence 432
    - halogen
      - – MMX 438
      - – MX complexes 436
    - H<sub>2</sub>dtoaCu 431
    - (HOC2H4)<sub>2</sub>-dtoa-Cu 432

- linkers (*contd.*)
    - Na[Cu<sup>III</sup>(pdt)<sub>2</sub>]·2H<sub>2</sub>O 433
    - Na[Ni(pdt)<sub>2</sub>]·2H<sub>2</sub>O 434
    - nitrogen-containing heterocyclic compounds 440
    - TCNQ
      - – Ag(TCNQ) 448
      - – chemical structure 447
      - – Cu(TCNQ) 447, 448
      - – electrical on/off switching characteristics 447
      - – electrically conductive properties 446
      - – electron acceptor molecule 446
      - – 1–1 M(TCNQ) (M=Cu(I), Ag(I)) 447
      - – Ni(TCNQ)<sub>2</sub> 450
      - – organic linker 446
      - – phase I and phase II 447
      - – Ru<sub>2</sub><sup>II,II</sup>/TCNQ(R) 451
      - – single crystal X-ray diffraction data 451
      - – Ti(TCNQ) 449
  - Langmuir surface 146
  - lanthanide containing MOFs (Ln-MOFs)
    - carboxylate ligands 236
    - chemical properties 232, 237
    - design and synthesis 235
    - electronic properties 246
    - 4f-states 232, 233, 247–249, 262
    - ligands and metal ions 237
    - luminescence of 249
    - noncarboxylate ligands 241
    - sensing applications 257
  - layer-by-layer (LBL) deposition method 523
  - ligand field stabilization energy(LFSE) 146
  - linkers, optical functionality
    - chemical sensing 483
    - design principles
      - – computational modeling 474
      - – DEF 471
      - – DFT calculations 474
      - – DMF 472
      - – enzymes and proteins structure 465
      - – inorganic and organic components 465
      - – optoelectronic applications 465
      - – primary structure 465, 473
      - – quantum chemistry methods 474
      - – quaternary structure 471, 472, 474
      - – SBU 465
      - – secondary structure 470, 473
      - – tertiary structure 471, 473
      - – electronic structure 463
    - light-harvesting
      - – guest-facilitated energy transfer 478
      - – organometallic complexes 477
      - – porphyrins 476
      - – MRI 480
      - – optical imaging 481
      - – radiation detection 485
      - – X-ray computed tomography imaging 482
  - linking triangular M<sub>3</sub>(pz)<sub>3</sub> clusters 319
  - liquid assisted grinding (LAG) 745
  - liquid-phase epitaxy (LPE)
    - laterally structured substrates 527
    - magMOF particles 531
    - nan and microparticles 529
    - non-structured substrates 526
  - local defects 810
  - luminescent MOFs (LMOFs) 94
- m**
- M<sub>3</sub>(pz)<sub>3</sub> cluster 319
  - M<sub>4</sub>(pz)<sub>4</sub> clusters 319
  - MagGel 546
  - magic angle spinning (MAS) 608
  - magnetic resonance imaging (MRI) 480
  - Maxwell–Stefan (M–S) diffusivity 784
  - metal azolate frameworks (MAFs) 309, 310
    - imidazolate coordination modes 311
    - – polyimidazolate 315
    - – zeolitic and zeolite-like frameworks 311
    - pyrazolate coordination modes 317
    - – chain-based pillared-column frameworks 324
    - – cluster-based networks 318
  - tetrazolate modes 336
  - triazolate coordination modes
    - – 1,2,3-modes 332
    - – 1,2,4-modes 329
  - metal-catecholates 442
  - metal coordination number (mCN) 807, 809
  - metal functionalized ligands 352
  - metal-ion vacancies 809, 812
  - metal organic framework (MOF)
    - metal to ligand charge transfer transitions (MLCT) 212
  - metal–metalloporphyrin frameworks (MMPFs) 125
  - metal–organic frameworks (MOFs) 73, 273
    - additives 171
    - applications 73
    - azolate ligands 178
    - catalysis 345
    - cation possess 172
    - characterization methods 3, 766
    - charge density 171
    - components 2
    - CPO-27(Fe)/MOF-74(Fe) crystal structure 179



- crystallisation, *see* crystallisation
- $\text{Cu}^{2+}$ , in  $\text{Cu}_{2.97}\text{Zn}_{0.03}(\text{btc})_2$
- - adsorbate complex, geometry of 642, 644
- - axial ligand field 640
- - CW EPR 638
- - dehydration 640
- - g and A tensor parameters 638, 640
- - HF interactions 641
- - OS ENDOR experiments 641, 642
- - OS HYSORE experiments 642
- - oxygen coordination 640
- - parameters 640
- - powder patterns 639
- - PW units 638, 640
- - spin Hamiltonian parameters 640
- - temperature range 638
- defects
- - Schottky- or Frenkel-type point 809
- - characterization 811
- - defective linker 796
- - defects engineering 796
- - distribution and interactions 810
- - inherent defects 796
- - metal-ion vacancies 809
- - substitutional point defects 810
- - concentration 796
- - defects engineering 805
- - definition 796
- - DL 807
- - HKUST-1 813
- - internal defects 806
- - interstitial 809
- - linker-vacancies 807
- - mCUs 809
- - metal-sites, modification of 808
- - MOF-5 813
- - PXRD 805
- - single solid 795
- - spectroscopic techniques 805
- - surface defects 806
- - types of 807
- - typical solids 796
- - UiO-66 816
- - uncoordinated (dangling) linker functions, ligator-site 809
- - XRD patterns 805
- design, characterizations 376
- device integration 1
- dimeric paddle wheel motif 180
- economy 2
- electrical properties measurement 422
- electronic conductivity 421
- first-row transition metal centers 73
- flexibility 1
- functions 2
- heterogeneity 795
- highly porous 279
- industrial applications 74
- innovation cycles 1
- ligands 421
- linkers, *see* linkers
- materials genome 1
- metal carboxylates
- - crystal structures 176
- - dicarboxylate ligands 174
- -  $\text{M}^{3+}$  poly-carboxylate structures 174–176
- - SBUs 174, 175
- - 'unusual' iron/vanadium polycarboxylate 176, 177
- metal(II) dioxohydroquinone structures 179
- metallic clusters 137
- metal phosphonates 172
- molecular simulations, *see* molecular simulation
- molybdenum(VI) 172
- nanoparticles 3
- nanoscale synthesis 181
- node and spacer principle 43
- organic linkers, *see* extended organic linkers
- paramagnetic species 629
- POM-MOFs 172
- poor hydrothermal stability 1
- post-synthetic modification 361
- - organocatalyst grafting 361
- - post-synthetic metalation 366
- Pourbaix diagrams 171
- prebuilt units 181
- properties
- - acidic behavior 184
- - bioapplications 186
- - catalytic 189
- - flexible 182
- - fluid separation 192
- - gas storage 192
- - inclusion 190
- - mechanical 183
- - redox 188
- - stability issues 185
- redox-active metals 422
- roadmaps 2
- self-assembled frameworks 347
- sensor integration 1
- SBU, *see* secondary building unit (SBU)
- $\sigma$ -bond 421
- suppliers 823

- metal–organic frameworks (MOFs) (*contd.*)
  - ultrahigh porosity, design 273
  - predictability 730
- metal–organic polyhedra (MOPs) 45
  - cuo-MOP 64
  - rht 62
  - cages 62
  - fcu net 62
  - functionalized mBDC derivatives 61
  - MOP-1, construction of 61
  - quadruple cross-linking 63
- metal-to-ligand charge transfer (3MLCT) 477
- 4,4',4'',4'''-((methanetetrayltetrakis(benzene-4,1-diyl))tetrakis(ethyne-2,1-diyl))tetra-benzoic acid (H<sub>4</sub>MTBA) 293
- 4',4''',4''''',4''''''-methanetetrayltetrakis-(1,1'-biphenyl)-4-carboxylic acid (H<sub>4</sub>MTBP) 293
- 4',4''',4''''',4''''''-methanetetrayltetrakis([1,1'-biphenyl]-3,5-dicarboxylic acid) (H<sub>8</sub>MTBDA) 301
- 5',5''''-methylenebis([1,1'3',1''-terphenyl]-4,4''-dicarboxylic acid) (H<sub>4</sub>PMTB) 297
- microporosity 84, 87, 124
- microwave synthesis 500
- MIL-53 175, 177, 183, 191, 346, 578, 701, 704
- MIL-68 174, 175, 378, 380
- MIL-88 175, 187, 781
- MIL-100(Al) (Material of Institute Lavoisier-100) 827
- MIL-101(Cr) (chromium(III) terephthalate, Material of Institute Lavoisier-101) 371, 828
- MIL-125
  - alcohol oxidation 141
  - photocatalytic properties 142
  - structure 141
- minimal transitivity principle 34
- mixed metal–organic frameworks (M'MOFs)
  - asymmetric catalysts 207
  - BINAP-MOF system 207
  - 2,2'-bipyridyl-derived linkers 209
  - bpy-UiO-Ir system 209
  - catenation isomer 209
  - chiral active catalysts 209
  - emission and high Stern–Volmer values 212
  - functional sites 207
  - hybrid solid 207
  - layer-by-layer method 212
  - metalloligand approach 207
  - MLCT 212
  - palladium(II) and platinum (II) coordination bonds 212
  - photoactive Ru(II) trisbipyridine based systems 212
  - photocatalytic properties 212
  - MOF structure 212
  - ultra nanoporous nature 213
- Mn-por-2 357
- modulation approach 808
- MOFomics 580
- MOFs, *see* metal–organic frameworks (MOFs)
- molecular dynamics (MD) simulation 777
  - benzene 785
  - corkscrew-type diffusion mechanism 786
  - corrected diffusivity 784
  - 2D-free energy map 786
  - intra-crystalline diffusivities 784
  - molecular diffusion 783
  - pore 786
  - rotational motions 784
  - super-mobility 784
  - self-diffusivity 783, 785
  - thermodynamic properties 788
  - translational motion 784, 785
  - transport/Fickian diffusivity 784
  - unusual diffusion 784
- molecular modelling 765
- molecular simulation
  - A520 770
  - adsorption
    - – adsorbent/adsorbate interactions 774
    - – extended charge equilibration PQeq strategy 774
    - – force field 774
    - – GCMC simulations 775
    - – LJ parameters 774
    - – Monte Carlo simulations 773, 774
    - – rationalization 780
  - cell parameters 769
  - computational tool 788
  - corrected diffusivity 784
  - isoreticular MOFs 767
  - ligand replacement strategy 767
  - MD
    - – corkscrew-type diffusion mechanism 786
    - – unusual diffusion 784
    - – 2D-free energy map 786
    - – benzene 785
    - – intra-crystalline diffusivities 784
    - – molecular diffusion 783
    - – pore 786

- – super-mobility 784
  - – self-diffusivity 783, 785
  - – thermodynamic properties 788
  - – translational and rotational motions 784
  - – transport/Fickian diffusivity 784
  - molecular modelling 765
  - NMR spectroscopy 768
  - pre-defined Building Blocks 766
  - PXRD 766
  - SBUs 766
  - structure prediction
    - – AASBU approach 771
    - – “bottom up” approach 771
    - – NU-100 771
  - – trial-and-error experimental methods 770
  - – UiO-66(Zr) 771
  - – UiO-66(Zr)-(CO<sub>2</sub>H)<sub>2</sub> 773
  - tools 766
  - translational motion 785
  - monoglyceride synthesis
    - Sn-EOF catalyst 682
    - ZIF-8 catalyst 680
  - multiple Linear Regression (MLR) methods 781
- n**
- nanoparticles 3
    - carbon 494
    - crystalline 495
    - nanoscale-effects 496
    - nanostructured materials 495
    - nanotoxicology 514
    - natural 494
    - NMOFs, *see* nanoparticles made of metal–organic frameworks (NMOFs)
    - size 495
    - synthesis 497
  - nanoparticles made of metal–organic frameworks (NMOFs)
    - applications
      - – catalytic properties 513
      - – for diagnostics 510
      - – gas sorption properties 512
      - – nanocarrier 507
      - – thin films 511
    - block-copolymers 505
    - bottom-up approaches 504
    - core-shell nanoparticles 505
    - external surface functionalization 503
    - factors 498
    - liquid–solid interfacial reactions 505
    - liquid–liquid interfacial reaction 505
    - microwave synthesis 500
    - morphology modulation 501
    - one-dimensional (1D) 504
    - reverse microemulsion method 501
    - solvo-thermal method 500
    - spontaneous precipitation method 499
    - three dimensional (3D) 504
    - top-down approaches 502, 504
    - two-dimensional (2D) 504
    - ultrasonic sound 501
    - zero-dimensional (0D) 504
    - nanoscale-effects 496
    - nanoscale Ln-MOFs 262
    - nanotoxicology 514
    - narrow pore (NP) phase 648
    - natural nanoparticles 494
    - natural tilings 26
    - net/topology, MOFs 6
      - derived nets 28
      - MIL-47/53 simplifications 33
      - minimal transitivity principle 34
      - RCSR database 27
      - symbols
        - – face symbols 18
        - – point symbols 17
        - – RCSR symbols 19
        - – Schläfli symbols 13
        - – vertex symbols 13
    - neutron powder diffraction (NPD)
      - adsorption chamber 709
      - Co<sub>2</sub>(m-dobdc) 711, 712
      - Co<sub>2</sub>(dobdc) 711
      - Cu<sub>3</sub>(btc)<sub>2</sub> 710
      - Fourier map analysis 710
      - hydrogen and methane adsorption 709
      - neutron scattering properties 709
      - NOTT-101 material 712
      - redox processes 713
      - X-rays scattering 709
    - [Ni<sub>3</sub>(μ<sub>4</sub>-OH)4(μ<sub>4</sub>-H<sub>2</sub>O)<sub>2</sub>(pz)<sub>12</sub>] clusters 324
    - 4,4',4'',4''',4''''-,4'''''-((nitrilotris(benzene-4,1-diyl)tris(azanetriyl))hexabenzic acid (H<sub>6</sub>TTA) 299
    - 4',4''',4'''''-nitrilotris([1,1'-biphenyl]-4-carboxylic acid) (H<sub>3</sub>TCBPA) 291
    - NU-1000 system 164
    - nuclear magnetic resonance (NMR) 110, 629, 635, 751, 768
      - <sup>1</sup>H 614
      - <sup>2</sup>H NMR measurements 623
      - host–guest interactions
        - – <sup>13</sup>C 619
        - – <sup>129</sup>Xe 616
      - hydrocarbon diffusion 623

- nuclear magnetic resonance (NMR) (*contd.*)
    - ortho/para ratio 621
    - solid-state
      - $^{13}\text{C}$  612, 613
      - $^2\text{H}$  614
      - CP 609
      - DNP 610
      - inorganic MOF components 615
      - line broadening interactions 607
      - probing spatial proximities 612
      - SEOP 611
  - nuclear Zeeman interaction 632
  - 4,4',4'',4'''-(6,7,9,10,17,18,20,21-octahydrodibenzo[b,k][1,4,7,10,13,16]hexaoxacyclo-octadecine-2,3,13,14-tetrayl)tetraazobenzene acid [H<sub>4</sub>TBADB-18Cr6] 296
  
  - O**
  - octatopic linkers
    - H<sub>8</sub>MTBDA 301
    - H<sub>8</sub>TBCPPP-H<sub>2</sub> 302
    - H<sub>8</sub>BTTCDD 302
    - H<sub>8</sub>TDM 302
    - H<sub>8</sub>TDPEPE 302
  - open metal sites (OMS)
    - fof topology 50
    - nbo topology 51
    - tbo net 49
    - 4,4-c nbo type net 50, 51
    - BET 49
    - HKUST-1 49
    - MOF-11 48
  - organic light-emitting diodes (OLEDs) 474
  - organic linkers 278
  - organocatalyst grafting 361
  - organocatalyst ligands 347
  
  - P**
  - paddle-wheel complex 438, 638, 640
  - pair distribution function (PDF) analysis 760
  - PCN-223 159, 162
  - PCN-224 162
  - pentatopic linker 298
  - 3-periodic nets 15
  - 4,4'-((1,4-phenylenebis(oxy))bis(methylene))dibenzoic acid (H<sub>2</sub>POMDA) 289
  - platinum group metal–organic frameworks
    - Hofmann type
      - azpy series 217
      - bpb ligand 221
      - bromide and iodide derivatives 217
      - chemical analyses 219
      - chemical migration 217
      - chloride derivative 217
      - clathrate compounds 219
      - crystal packing effects 215
      - diffraction efficiency values 218
      - 2D networks 214
      - electronic and steric factors 218
      - HS and LS states 216
      - hydration–dehydration process 220
      - large binding energy and adsorption position 217
      - pore size and functionality 217, 218
      - porosity 221
      - porous properties and SCO phenomenon 215
      - pyrazine ligands bridges 215
      - redox active Pt(II) centers 217
      - ST behavior 220
      - thermal hysteresis loops 215
      - X-ray single crystal structural analysis 218, 219
    - M' MOFs
      - asymmetric catalysts 207
      - BINAP-MOF system 207
      - 2,2'-bipyridyl-derived linkers 209
      - bpy-UiO-Ir system 209
      - catenation isomer 209
      - chiral active catalysts 209
      - emission and high Stern–Volmer values 212
      - functional sites 207
      - hybrid solid 207
      - layer-by-layer method 212
      - metalloligand approach 207
      - MLCT 212
      - palladium(II) and platinum (II) coordination bonds 212
      - photoactive Ru(II) trisbipyridine based systems 212
      - photocatalytic properties 212
      - MOF structure 212
      - ultra nanoporous nature 213
  - one-dimensional coordination polymers 222
  - single node 204
  - two-dimensional coordination polymers 224, 225
- point/isolated defects 810
- point symbols 17
- polyimidazolate 315
- polyoxometallates (POMs) 172
- porosity determination
  - adsorptive choice 583
  - BET equation 586
  - geometrical surface area
  - computational method 580

- – crystallographic data 580
  - – experimental isotherms 582
  - – factors 582
  - – GCMC simulations 581
  - – MOFomics 580
  - – molecular simulations 580
  - – molecule/crystal 578
  - – nitrogen molecule 580
  - – PLATON 579
  - – pore volume and surface area 581
  - – Poreblazer 580
  - – probe sphere 578
  - – solvent accessible 578
  - – solvent excluded 578
  - – van der Waals surface 578
  - pore volume 587
  - PSD 587
  - porous coordination polymers (PCPs) 698
  - porous interpenetrated zirconium organic frameworks (PIZOFs) 163
  - 4',4''',4''''',4''''''-(porphyrin-5,10,15,20-tetrayl)tetrakis((1,1'-biphenyl)-4-carboxylic acid)) (H<sub>4</sub>TCBPP-H<sub>2</sub>) 295
  - post-synthetic metalation 366
  - post-synthetic modification (PSM) 110, 361, 524
    - internal surface modification 534
    - metalation 366
    - organocatalyst grafting 361
    - SURGELS 534
  - powder X-ray diffraction (PXRD) patterns 76, 766, 772, 805
  - pressing method
    - Cu-BTC (HKUST-1) pellet 567
    - deformation properties 566
    - HKUST-1 567
    - industrial applications 566
    - MIL-100(Fe) pellets 569
    - pellet 566
    - SIM-1 567
    - specific surface area 568
    - two-step process 567
    - UiO-66(Zr) 569
    - ZIF-8 567
  - pressure-swing-adsorption (PSA) processes 600
  - primitive cubic (pcu) topology 52, 62
  - 1-propylene-3-methylimidazolium chloride (AmimCl) 75
  - Prussian Blue (PB) 452
  - Prussian blue analogues (PBAs) 454, 456
  - pulsed ENDOR spectroscopy 637
  - PXRD, *see* powder X-ray diffraction (PXRD) patterns
  - pyrazolate coordination modes 317
    - chain-based pillared-column frameworks 324
    - cluster-based networks 318
  - 4,4',4'',4'''-(pyrene-1,3,6,8-tetrayl)tetrabenzoic acid (H<sub>4</sub>TBAPy) 296
  - 3,5-pyridinedicarboxylate (pdc) 75, 81
- q**
- QENS 785
  - quantitative structure property relationship (QSPR) 780, 782
  - quantum chemistry methods 474
  - quantum dots 496
  - quartz crystal microbalance (QCM) 525, 536
- r**
- radiation detection 485
  - Raman scattering 760
  - Raman spectroscopy 107, 658, 660, 662
  - rare earth ions (REI) 629
  - RCSR symbols 19
  - reasonably priced MOFs 825
  - redox processes
    - NPd 713
    - XAS 714
  - reverse microemulsion method 501
  - Rh-based catalysts 376
  - 24-c rhombicuboctahedral (rco) 61
  - rhombicuboctahedron triangle (rht) 62
  - RHO-[Zn(bim)<sub>2</sub>] 312
  - RIXS techniques 379
  - rotational echo adiabatic passage double resonance (REAPDOR) experiment 612
  - rotational echo double resonance (REDOR) technique 612
  - RuDCBPY 477
- s**
- Saito-Foley (SF) method 587
  - salen-based MOF oxidation catalysts 358
  - scanning electron microscopy (SEM) 536
  - Schläfli symbols 13
  - SciFinder® 105
  - secondary building unit (SBU) 137, 174, 465, 766, 772
    - cluster geometries 45
    - concept 43
    - eight extension points 55
    - four extension points 46
    - infinite secondary building units 65
    - MOP 45
    - – cuo-MOP 64
    - – cages 62

- secondary building unit (SBU) (*contd.*)
  - design principles 43
  - fcu net 62
  - functionalized mBDC derivatives 61
  - MOP-1, construction of 61
  - quadruple cross-linking 63
  - nine extension points 56
  - OMS
  - nbo topology 51
  - tbo net 49
  - 4,4-c nbo type net 50, 51
  - BET 49
  - HKUST-1 49
  - MOF-11 48
  - six extension points
  - carboxylate moieties 55
  - MOF-3 51, 52
  - MOF-5 52, 53
  - MOF-177 54
  - octahedral and trigonal prismatic geometries 51
  - octahedral clusters 52
  - pcu topology 52
  - qom 54
  - rutile rtl and pyrite pyr nets 54
  - toz topology 54
  - trigonal prismatic clusters 54
  - UCMC-150 56
  - $Zn_4O(-COO)_6$  54
  - three extension points 45
  - twelve extension points 57
- self-assembled frameworks
  - metal functionalized ligands 353
  - organocatalyst ligands 347
- self-assembled monolayer (SAM) 524
- shaping methods
  - extrusion
    - advantages 562
    - Caleva Mini Screw Extruder 563
    - Cu-BTC 564
    - extrudates 562
    - MOF monoliths 563
    - process 563
    - STYs 563, 565
    - synthetic approach 563
    - twin screw extruder 565
  - granulation, *see* Granulation
  - industrial/large-scale applications 551
  - plasticizers 552
  - porous materials 552
  - powder forms 551
  - pressing
    - Cu-BTC (HKUST-1) pellet 567
    - deformation properties 566
    - HKUST-1 567
    - industrial applications 566
    - MIL-100(Fe) pellets 569
    - pellet 566
    - SIM-1 567
    - specific surface area 568
    - two-step process 567
    - UiO-66(Zr) 569
    - ZIF-8 567
  - pressurized methods 552
  - spray drying 560
  - techniques 551
- SIM catalyst 364
- simulated annealing and parallel tempering techniques 766
- single-crystal XRD 163
- small phase incremental alteration (SPINAL) 609
- small-angle X-ray scattering (SAXS) 731, 734
- Soc-MOF 588
- SOD-[Zn(bim)<sub>2</sub>] 312
- solvent-assisted ligand incorporation (SALI) 166, 168
- solvent-assisted linker exchange (SALE) 355
- solvothermal reaction 334, 500
- Sonogashira coupling 284
- space-time yields (STYs) 563, 565
- Spallation Neutron Source (SNS) 709
- spectroscopic monitoring
  - carbon dioxide adsorption
    - adsorption–desorption profiles 672
    - CO<sub>2</sub>-mmen-M<sub>2</sub>(dobpdc) 673–675
    - 2D IR Map 673
    - doubly degenerate bending vibration  $\nu_2$  669
    - FTIR spectra 669, 670
    - hysteresis loop 672
    - “in situ” IR spectra evolution 672
    - M<sub>2</sub>(dobdc) analogues 672
    - molecular symmetry 669
    - NPD experiments 672
    - $\nu_3$  stretching mode 670
    - symmetric stretching  $\nu_1$  669
    - vibrational frequencies 671
  - carbon monoxide adsorption
    - DFT 668
    - high and low frequency 667
    - IAST 669
    - IR absorption bands 665
    - Madelung effects 667
    - Mg<sup>2+</sup>-CO interaction 666
    - molecular probe 665
    - outgassed sample 667
    - perturbing effect 668

- transition metal ions 665
  - catalytic reactions
  - Cr-MIL-101 encapsulated Keggin phosphotungstic acid 678
  - Fe<sub>2</sub>(dobdc) catalyst 675
  - Sn-EOF catalyst 682
  - ZIF-8 catalyst 680
  - spin density wave (SDW) 437
  - spin transition (ST) behavior 220
  - spin-exchange optical pumping (SEOP) 611
  - 4,4',4'',4'''-(9,9'-Spirobi[fluorene]-2,2',7,7'-tetrayl)tetrabenzoic acid (H<sub>4</sub>SFTT) 298
  - spontaneous precipitation method 499
  - spray drying method 560
  - Stern–Volmer plot 330
  - stilbenedicarboxylic acid 114
  - substitutional point defects 810
  - 3,3',4,4'-sulfonediphenyltetracarboxylic acid 117
  - supercritical drying (SCD) 577
  - surface plasmon resonance (SPR) 525
  - surface-mounted MOFs (SURMOFs)
    - applications
    - BSs 540
    - cell culture substrate 544
    - conductive Polymer 546
    - electronic properties 540
    - MagGel 546
    - reversible and remote-controlled modification 544
    - chiral separation 524, 539
    - heterostructures
    - heteroepitaxial growth 531
    - heterogeneous synthesis approach 531
    - lattice parameters 532
    - solvothermal synthesis 531
    - HKUST-1 525
    - LBL deposition method 523, 525
    - LPE-based synthesis
    - laterally structured substrates 527
    - magMOF particles 531
    - nan and microparticles 529
    - non-structured substrates 526
    - pinhole-free films 523
    - PSM 524
    - internal surface modification 534
    - SURGELs 534
    - QCM/SPR 525
    - surface barrier phenomenon 536
    - surface preparation methods 524
    - surface-sensitive techniques 535
    - with SAM 524
  - SURMOFs, *see* Surface-mounted MOFs (SURMOFs)
  - Suzuki coupling 284
  - Swiss-Norwegian beamline (SNBL) 693
  - Systre 10
- t**
- tailoring activity 365
  - tetraberyllium oxohexaacetate 76
  - 4,4'-(5,6,12,13-tetrachloro-1,3,8,10-tetraoxo-1,3,8,10-tetrahydroanthra[2,1,9-def 6,5,10-d'e'f']diisoquinoline-2,9-diy) dibenzoic acid (H<sub>2</sub>CONQDA) 288
  - 7,7,8,8-tetracyanoquinodimethane (TCNQ)
    - 1–1 M(TCNQ) (M = Cu(I), Ag(I)) 447
    - Ag(TCNQ) 448
    - chemical structure 447
    - Cu(TCNQ) 447, 448
    - electrical on/off switching characteristics 447
    - electrically conductive properties 446
    - electron acceptor molecule 446
    - Ni(TCNQ)<sub>2</sub> 450
    - organic linker 446
    - phase I and phase II 447
    - Ru<sub>2</sub><sup>II,II</sup>/TCNQ(R) 451
    - single crystal X-ray diffraction data 451
    - Tl(TCNQ) 449
  - tetradedral M<sub>4</sub>(pz)<sub>4</sub> cluster 319
  - tetragonal-prismatic clusters 323
  - tetrahydrofuran 79
  - 1,3,6,8-tetrakis(*p*-benzoicacid)pyrene (H<sub>4</sub>TBAPy) ligand 166
  - 5,10,15,20-tetrakis(4-carboxybiphenyl) porphyrin 125
  - tetrakis(4-carboxyphenyl)porphyrin (TCPP) ligand 156
  - 5,10,15,20-tetrakis(4-carboxyphenyl)porphyrin 125
  - tetratopic linkers
    - H<sub>4</sub>ADBTD 296
    - H<sub>4</sub>BBCDC 297
    - H<sub>4</sub>BNETBA-OEt 297
    - H<sub>4</sub>CBI 295
    - H<sub>4</sub>CQDA-OEt 296
    - H<sub>4</sub>PMTB 297
    - H<sub>4</sub>SFTT 298
    - H<sub>4</sub>TBADB-18Cr6 296
    - H<sub>4</sub>ETTC 296
    - H<sub>4</sub>ATB 293
    - H<sub>4</sub>MTBA 293
    - H<sub>4</sub>MTBP 293
    - H<sub>4</sub>QPTCA 295
    - H<sub>4</sub>TBAPy 296

- tetratopic linkers (*contd.*)  
 – H<sub>4</sub>TCBPP-H<sub>2</sub> 295  
 tetravalent metals 137  
 tetrazolate modes 335  
 thermal gravimetric analysis(TGA) 151  
 thiazoledicarboxylate (thz) 85  
 “through-bond” approach 426  
 “through-space” approach 426  
 Ti<sub>8</sub>O<sub>8</sub>(OH)<sub>4</sub>(CO<sub>2</sub>)<sub>12</sub> 141  
 Ti<sub>8</sub>O<sub>8</sub>(OOCR)<sub>16</sub> 140  
 tilings 26  
 titanium-based MOFs 167  
 – high stability and photocatalytic activity 139  
 – MIL-125  
 – – alcohol oxidation 141  
 – – photocatalytic properties 142  
 – – structure 141  
 – oxo clusters 139, 140  
 – vs. Zr-MOFs 139  
 TMI, *see* transition metal ions (TMI)  
 “top down” approach 771  
 topological density 24  
 topology 9  
 – zirconium-porphyrin MOF 157  
 ToposPro 10, 25  
 total adsorption 600  
 transfer of populations in double resonance (TRAPDOR) 612  
 transition metal ions (TMI) 629, 630  
 – Co<sup>2+</sup> 650  
 – Cr<sup>3+</sup> 648  
 – Cu<sup>2+</sup> 644  
 – paramagnetic 3d 650  
 – Ti<sup>3+</sup> and V<sup>4+</sup> 646  
 transitivity 26  
 4, 4', 4''-s-triazine-1,3,5-triyltri-p-aminobenzoic acid (H<sub>3</sub>TATAB) 293  
 4,4',4''-(triazine-2,4,6-triyl-tris(benzene-4,1-diyl))tribenzoic acid (H<sub>3</sub>TAPB) 292  
 4,4',4'';-(1,3,5-triazine-2,4,6-triyl)tribenzoic acid (H<sub>3</sub>TATB) 292  
 triazolate coordination modes  
 – 1,2,3-modes 332  
 – 1,2,4-modes 329  
 triethanolamine (TEA) 143  
 triethanolamine (TEOA) 142  
 triphenylene-2,6,10-tricarboxylic acid (H<sub>3</sub>TTCA) 292  
 1,3,5-tris(3,5-di(3,5-dicarboxyphenyl-1-yl)phenyl-1-yl)benzene (H<sub>12</sub>TDDPB) 303  
 tritopic linkers 289  
 – H<sub>3</sub>TATAB 293  
 – H<sub>3</sub>BBC 291  
 – H<sub>3</sub>BTB 291  
 – H<sub>3</sub>BTE 291  
 – H<sub>3</sub>BTN 292  
 – H<sub>3</sub>HTB 291  
 – H<sub>3</sub>TAPB 292  
 – H<sub>3</sub>TATB 292  
 – H<sub>3</sub>TCBPA 291  
 – H<sub>3</sub>TTCA 292  
 tritopic trimetric acid 118  
 twisted boracite (tbo) net 49  
 two pulse phase modulation (TPPM) 609
- u**  
 UiO-66 (Universitetet i Oslo 66, zirconium terephthalate) 145–156, 163, 372, 582, 595, 596, 831  
 UiO-67 framework 375, 376, 380  
 ultra high-performance liquid chromatography (UHPLC) 410  
 ultrasonic sound 501  
 UMCM-1-AMpz 371  
 UMCM-1-AMsal 371  
 urea functional linkers 349  
 urea-functionalized NU-601 MOF 349
- v**  
 Vantablack-material 496  
 vertex symbols 13  
 vertex-transitive nets 27  
 vibrational spectroscopy 750  
 volume-specific surface area (VSSA) 557  
 volumetric method  
 – advantage 598
- x**  
 X-ray absorption fine-structure spectra (XAFS) 753  
 X-ray absorption near edge structure (XANES) 379, 714  
 – CPO-27(Ni) 715  
 – crystallisation 753  
 – defects, MOF 812  
 – Fe K-edge 720, 721  
 – HKUST-1 714, 715  
 – pre-edge 722, 723  
 X-ray absorption spectroscopy (XAS)  
 – capillary and close cycle helium cryostat 714  
 – energy value 713  
 – EXAFS 714  
 – – clusters 715  
 – – CPO-27(Ni) 715  
 – – Cu atoms 714



- DUT-8(Ni) 717
- fit parameters 720
- HKUST-1 716
- $K_2PtCl_4$  and UiO-66 -  $NH_2$  721
- $O_2$ -loaded, pre-edge data 720
- parameter values 723
- post-activation data 723
- limitation 714
- $PtCl_4^{2-}$  infiltration 719
- redox processes 714
- scattering techniques 723
- UiO-66 716
- XANES 714
- CPO-27(Ni) 715
- Fe K-edge 720, 721
- HKUST-1 714, 715
- pre-edge 722, 723
- X-ray diffraction (XRD) 110, 138, 159
- X-ray photoelectron spectroscopy (XPS) 812
- X-ray powder diffraction 107
- XANES, *see* X-ray absorption near edge structure (XANES)
  
- Z**
- zeolites 7, 105
- zeolitic and zeolite-like frameworks 311
- zeolitic imidazolate frameworks (ZIFs) 440, 740, 747, 830
- zeotype topology (ZMOF) 388
- zirconium-based MOFs 138, 167, 372, 373
- zirconium-based MOFs
  - advantages 144
  - NU-100 system 164
  - PIZOF ligands 163
  - porphyrinic MOFs
    - elongated ligands 160, 161
    - structure of 162
    - topology 157
    - $Zr_6O_8$  cluster and TCPP ligand, connections of 156, 158
  - stability of 144
  - structure 144
  - UiO-66
    - and derivatives 145
    - applications 151
    - modulated synthesis 148
    - post-synthetically exchange 147
    - stability 146
  - Zr(IV) based metal-organic frameworks 163
- sql*-[Zn(bim)<sub>2</sub>] 312
- [Zn<sub>2</sub>(Httb)<sub>2</sub>] 336
- Zn<sub>4</sub>O(pz)<sub>3</sub>(COO)<sub>3</sub> clusters 321
- Zn<sub>4</sub>O(pz)<sub>4</sub>(COO)<sub>2</sub> clusters 320
- ZnPOR-1 catalyzed acyl transfer reaction 354

