

## Index

### a

- acid recovery using diffusion dialysis 386–389
- acylation
  - of alcohols 111–112
  - of amides 118–119
  - of amines 112–114
  - of carboxylic acids 116
  - of crown ether 149
  - of enols 114–116
  - of ketones 116–118
  - of sulfonamides 119–121
- alkaline anion-exchange membranes (AAEMs) 107, 123, 148, 184
- alkaline membranes 10, 180, 182, 191, 201, 203, 205, 215, 382, 385
- amidoxime of dibenzodioxin-based PIMs 233–234
- anion exchange membranes (AEMs) 1, 22, 191
  - imidazole-type side-chain 65
  - poly-crown ether-based 147
  - polyimide-based 138
  - SPEK-based 144
- anionic active polymerization technology 61, 63
- area resistance 27–29, 256, 308, 312, 313, 339
- aromatic carboxylic monomers 60
- aryl ether-free
  - 2,7-diphenylfluorene-based copolymers 186
- azaspirocyclic cations 191

### b

- benzyltrimethylammonium (BTMA) 180, 245, 385

- binary superacid 171
- biphenyl-2,5-diol 49
- bipolar membrane electro dialysis (BMED)
  - acid production 362–364
  - alkali production 360–362
  - basic theory of bipolar membranes 358–360
  - production of CO<sub>2</sub> conversion 364–365
  - schematic diagram 356–357
- 9,9-bis(3,5-dimethoxy-4-hydroxyphenyl) fluorene 66
- 1,3-bis(4-fluorophenoyl) benzene 56
- bisphenol A-type PES 50, 53, 62, 63, 69
- 2,2-bis-(p-hydroxyphenyl)-pentafluoropropane sulfonate 57
- 4-(3-bromoalkyl oxygen) benzaldehyde 69
- 4-bromo-N,N,N-trimethylbutane-1-ammonium bromide 71
- 7-bromo-1,1,1-trifluoroheptan-2-one 184
- Brønsted acid system 108–109, 170, 172, 179
- Brønsted/Lewis superacids 171, 173
- 4-(3-butenyl) styrene 70

### c

- 6-(9*H*-carbazol-9-yl)-*N,N,N*-trimethylhexan-1-aminium bromide 231
- carbocations 174, 179
- carbonyl-containing electrophilic monomers 182
- catalyst 4f 113
- cation-dipole interactions
  - alkaline stable ion conduction channel establishment 91–92

- cation-dipole interactions (*contd.*)  
 ion conduction channel establishment 88–91
- cation exchange membranes (CEMs) 1  
 polyimide-based 135  
 SPEK-based 140  
 sulfonic acid-type side-chain 63
- cationic COF membrane 296–300
- chlorine-free potassium fertilizers 370, 371
- 4-(chloromethyl) styrene (VBC) 83, 84, 253, 254, 256
- Coion diffusion 352
- comb-shaped poly(arylene indole piperidinium) copolymer (PITP-CnQx) 190
- commercial aromatic polymers 47
- commercial polybenzimidazole 50
- concentration polarization 23, 26, 28–30, 34, 308, 342
- conductivity, of membrane 38, 85, 91, 135, 141, 154, 190–191
- copper-catalyzed azide-alkyne click reaction 68
- covalent bonding-triggered (CBT)  
 pore-filling strategy 252, 253
- covalent organic framework (COF)  
 membrane 5, 285–292, 314  
 introduction 285–286  
 pore structure design 288  
 pore surface engineering 288–292  
 preparation  
 cationic COF membrane 296–300  
 self-standing COF (TpTAPA) membrane 300–305  
 sub-2-nm channels 292–296
- cross-linked quaternized TB polymers 5
- cross-linked sulfonated polyimide (SPI) membrane 137
- cross-linking side chain construction  
 hydrogen bonds 82  
 unsaturated end groups 81–82  
 unsaturated groups of main chains 82
- crown ethers 124, 137, 146, 149, 154
- current-voltage (I-V) curves 29–32, 34, 35, 358
- d**
- DD alkali recovery 333
- diamine monomers 54, 57–60, 72, 140, 289
- diazacyclooctane polymers 229
- dibenzo-15-crown-5-based lithium  
 conductive membranes 153–156
- diffuse layer 35, 36
- diffusion coefficient 25–26, 30, 239, 293, 295, 296, 346, 347
- diffusion dialysis (DD)  
 advantages of 327  
 basic theory of 327–328  
 components 330–331  
 recovery of steel waste acid 331–333
- difluorophenonitrile 56
- 4,4'-difluorophenophenone 56
- 2,8-m-dinaphthol 6-sulfonate 56
- diphenol monomers 49, 52, 54, 56, 57, 61
- 2,6-diphenyl phenol 62
- Donnan dialysis 343–347
- Donnan equilibrium 21–22, 24, 39, 345, 346
- Donnan exclusion potential 1
- dynamic transfer region construction  
 in AEMs 86–88  
 in CEMs 85–86
- e**
- efficient superacid-catalyzed  
 Friedel-Crafts polycondensations 207
- electrodialysis (ED)  
 advantages of 348, 350  
 approaches to improve the  
 concentration on 352–353  
 Coion diffusion 352  
 concentration 351  
 desalination 349–351  
 disadvantages of 348–351  
 limit current density 352  
 resource conversion 353–355  
 water migration 352
- electrodialysis metathesis (EDM)  
 high-salinity wastewater conversion 369–370  
 ionic liquids 367–369  
 potassium fertilizers 370–372
- electronanofiltration membranes (ENMs)  
 introduction 305–306  
 performance of 307–313  
 preparation of 306–307
- electrospinning 141, 245, 246, 265, 329
- ether-free polymer backbones 194, 213

**f**

- film-forming PAPipQs 185
- fluorinated sulfuric acid 169
- fluorinated triazine framework  
membranes 225
- Friedel–Crafts polycondensation reaction  
72, 73
- Friedel–Crafts reaction 65, 175
- fuel cells, application of 378–383

**g**

- guanidinium anion exchange groups 78

**h**

- N*-heteroaryl-4-(1-(2-oxo-chromen-3-yl)  
ethylideneamino) benzene-  
sulfonamide compounds 121
- heterocyclic anion exchange groups  
78–80
- heterocyclic nitrogen 178
- heteronafenyl phenolone 56
- high ion-permselective CC3 membrane  
281–285
- highly conductive poly [(terphenyl  
piperidinium)-*co*-(oxindole  
terphenylene)] (PTP-*x*) AEMs  
188, 190
- high-performance diffusion dialysis  
membranes 328–330
- high-salinity wastewater conversion  
369–370
- hydrolysis of dibenzodioxin-based  
microporous polymers 233
- hydroxyalkylation reaction 173, 175
- hyperbranched networks construction  
83–84
- hypercrosslinked ionic polymers (HIPs)  
138

**i**

- imidazole-type side-chain AEMs 65, 66
- inductively coupled plasma–optical  
emission spectrometry (ICP–OES)  
26, 293
- inorganic secondary building units  
(SBUs) 261
- ion-distillation technology 372–378
- ion exchange capacity 8, 9, 39–40, 77,  
114, 154, 184, 191, 229, 231, 236,  
247, 248, 252, 329, 346

- ion exchange membrane (IEM)  
applications 10
  - bipolar membrane electro dialysis  
356–367
  - diffusion dialysis (DD)  
basic theory of 327–330  
components 330–331  
recovery of steel waste acid 331–333
  - Donnan dialysis 343–347
  - electrodialysis 347–349
  - electrodialysis metathesis (EDM)  
367–372
  - fuel cells 378–383
  - industrial  
acid recovery using diffusion dialysis  
386–389  
resource recovery using  
electrodialysis 389–394
  - ion-distillation technology 372–378
  - reverse electro dialysis 334–343
  - water electrolysis 384–386
  - ion exchange membranes (IEMs)  
definition and classifications 1  
main categories of 1, 3  
potentials 11–13  
preparation of 7–10  
conventional methods 245–246  
in situ polymerization 255–256  
pore filling 250–253  
semi-interpenetrating polymer  
networks 246–250  
solvent-free strategy 253–255  
profile of 1–6
  - ion flux 26–27, 29, 269, 271, 282, 283,  
285, 306, 312, 328
  - ionic liquids 201, 367–369
  - ionic membranes, preparation of 179,  
180
  - ion permeability 25–26, 237, 277
  - ion transport behaviour,  
nanospace-confined membranes  
219–220
  - Iso-Reticular Metal–Organic Framework  
(IRMOF) 261, 264
- l**
- Lewis acid 60, 65, 108, 112, 116,  
170–172, 179, 229, 230
  - limiting current density 29–32, 35, 376
  - linear polycyclic aromatic hydrocarbons  
181

**m**

- MBTF proton membrane 212, 214
- mechanical strength of ion exchange membranes 40
- membrane electrode assembly (MEA) 204, 206, 207, 379–382, 386
- membrane potential 1, 22–25, 337
- membrane scaling and fouling 34–35
- membrane technology 105, 286, 330, 356
- Menshutkin reaction 71, 72, 74, 139, 181, 183, 185, 211
- metal cationic anion exchange groups 81
- metal-organic framework (MOF) membranes
  - high stability 264–265
  - introduction 261
  - preparation
    - polyMOF membrane 271–272
    - UiO-66(Zr/Ti)-NH<sub>2</sub>/polyamide mixed matrix membrane 268–271
    - UiO-66-NH<sub>2</sub> membrane 266–267
    - UiO-66-SO<sub>3</sub>H membrane 268
  - structural diversity 261–262
  - structural tunability 262–264
- 5-methoxy-2-iodophenylboronic acid (MIBA) 113
- methyl trifluoromethyl ketone 184
- Michaelis–Arbuzov reaction 210, 211
- microphase-separated IEMs 4, 5
- microphase separation of grafted polymers 47
- microporous polymers
  - negatively charged 232–233
  - positively charged 225–232
  - principle of 220–225

**n**

- Nafion® 4, 133, 135, 141, 142, 143, 237, 380
  - chemical structure of 46
  - functionalized monomers 71–73
  - inspiration of 45–47
  - morphologies of 60–61
  - side-chain AEMs with 73–74
  - synthesis from side-chain sulfonated monomers 54–60
  - synthesis via post-functionalization 47–54

- Nafion-H 172, 180
- naphthalene-based polybenzimidazole (NPBI) 204
- negatively charged microporous polymers 232–239
- nitronium superelectrophiles 172
- NPBI-QAx membrane 204
- nucleophilic substitution reactions 52, 62, 65, 66, 80, 143

**p**

- permselectivity 26–27, 236, 266, 268, 272, 274, 285, 310, 312–313, 339, 346, 352
- perylene-based polyimides
  - polyacylation route 127–131
  - polyimide-based AEMs 138–140
  - polyimide-based CEMs 135–138
  - polyimide-based IEMs 131–132
  - traditional route 125–128
- phenyl monomers 182, 184
- polyacylation
  - acylation of alcohols 111–112
  - acylation of amides 118–119
  - acylation of amines 112–114
  - acylation of carboxylic acids 116
  - acylation of enols 114–116
  - acylation of ketones 116–118
  - acylation of sulfonamides 119–121
  - advantages 123–124
  - challenges/opportunities 156–157
  - crown ethers 146
  - limitations 124
  - need for novel and sustainable membrane materials 105–107
  - perylene-based polyimides 125–140
  - of polymers 121–124
  - principle of 108–109
  - SPEK-based AEMs 144–146
  - SPEK-based CEMs 140–144
  - synthesis of aromatic polyelectrolytes 107–108
- poly(arylene ether sulfone) 180, 181, 254
- poly(arylene perfluorophenylphosphonic acid)s 211
- poly(arylene piperidinium)s (PAPipQs) 184
- poly(arylene sulfide)s 144
- poly(aryl piperidine) homopolymer ionomers 186
- poly(aryl piperidinium) membranes 204

- poly(biphenyl piperidine) anion exchange membrane 191
- poly(carbazole)-based anion-exchange polymer 209
- poly-crown ether-based AEMs 147
- poly-crown ether-based noncharged selective membrane (PCENS-M) 153–155
- polycyclic aromatic hydrocarbons 181, 182
- poly(diphenyl-terphenyl piperidinium) copolymer 186, 187
- polyelectrolyte membranes (PEMs) 105, 107, 108, 129, 137, 142–144, 211, 247, 251, 380, 385, 386
- poly[(fluorene alkylene)-co-(biphenyl alkylene)] (PFBA) 188
- poly(fluorenyl aryl piperidinium) ionomers 187
- polyimide-based AEMs 135, 138–140
- polyimide-based CEMs 135, 137–138
- polyimide-based ion exchange membranes (IEMs) 131, 134–135
- polymers of intrinsic microporosity (PIMs) 5, 221, 222
- poly(meta terphenyl piperidinium) copolymers 191
- polyMOF membrane 271–272
- poly(p-terphenyl perfluoroalkylene)s 210
- poly(spirobiindane-aryl ether sulfone) copolymers 231
- poly(terphenyl alkylene)s 185
- pore filling 250–253
- pore surface engineering 288–292
- porous organic cage membranes  
introduction 272–274  
preparation  
  high ion-permselective CC3 membrane 281–285  
  versatile channels 276–281  
structural properties of 274–276
- porous polyimide frameworks (PIFs) 138
- positively charged microporous polymers 225–232
- positively charged PIMs 5, 231, 232
- postsulfonation of PIMs or bottom-up approach 234–239
- potassium fertilizers 370–372
- pressure-retarded osmosis (PRO)  
operating mechanism diagram 334, 335
- q**
- QA38PPO membrane 88, 89
- QA38PPO-P membrane 88, 89, 91
- quasidiphenol monomers 54
- quaternary ammonium (QA) anion exchange groups 76–78
- quaternary phosphine anion exchange groups 80
- quaternized poly-carbazole (QPC-TMA) 207, 208
- quaternized Tröger's base (TB) polymers 5
- r**
- radiation-induced grafting 245
- resource recovery using electro dialysis 389–394
- reverse electro dialysis  
basic theory of 336–338  
electrode system 342–343  
ion exchange membrane 339–341  
spacers 341–342
- s**
- salinity gradient power (SGP) 334
- self-standing COF (TpTAPA) membrane 300
- semi-interpenetrating polymer network  
anion exchange membranes (semi-IPN AEMs) 248
- semi-interpenetrating polymer networks (semi-IPNs) 246
- side-chain AEMs design  
with grafted structures 75–76  
with similar Nafion structures 64–71
- side-chain CEMs design  
with grafted structures 61–64  
with similar Nafion structures 47–54
- side chain engineering  
microphase separation of grafted polymers 47
- Nafion 45–47  
theoretical basis of 45
- side-chain perfluoroalkyl sulfonate aromatic polymers 53
- side-chain sulfonated aromatic diamine monomers 54, 57, 58

- side-chain sulfonated aromatic dihalide monomers 54–56
- side-chain sulfonated aromatic diphenol monomers 54, 56
- side-chain sulfonated PES 51
- side-chain type microporous polymers 224, 225
- small angle X-ray diffraction (SAXS) 46, 88, 139, 193, 200–203
- solid superacid 171, 172, 179
- soluble conjugated microporous polymers 223
- solution casting 7, 9, 141, 142, 148, 225, 231, 245, 271
- solvent-free strategy 253–255
- SPEK-based AEMs 144–146
- SPEK-based CEMs 140, 145
- spiral wound diffusion dialysis (SWDD) component 330, 333
- spirobifluorene-based ladder polymer 237
- steric effect 50, 79, 81, 219, 220, 234
- Stern layer 35, 36
- sulfonated poly(arylene ether sulfone) desalination membranes 254
- sulfonated polybenzene 207
- sulfonated poly(ether ether ketone ketone) (SPEEKK) 133, 143
- sulfonated polyimides (SPIs) 107, 128, 129, 135, 136
- sulfonated poly(phenyl-alkane)s 236, 237
- sulfonated polyxanthene 219, 237
- sulfonated side-chain aromatic diamine monomers 59
- sulfonic acid-type side-chain CEMs 63
- superacid  
 acidity range of 171  
 definition and types of 169–172  
 solutions 169
- superacid-catalyst polymerization for IEM preparation 179–213
- superacid-catalyst, principle of 172
- superacid-catalyzed Friedel–Crafts polycondensation 186, 207
- superacid-catalyzed polyhydroxy-alkylation 177, 178, 181, 182
- superacid-catalyzed reactions 173  
 for polymer synthesis 175–178
- superelectrophiles 172, 173, 179
- superelectrophilic activation 172, 173, 175, 179, 182
- superelectrophilic reagent 172
- t**
- tertiary sulfur anion exchange groups 80–81
- thiophene-substituted conjugated microporous polymers 224
- Tröger's base (TB) 5, 122, 221
- transference numbers 23–25, 29–31, 35
- transition-metal-catalyzed cross-coupling reactions 183
- trifluoromethanesulfonic acid 72, 111, 143, 169, 225, 234, 239
- trimethyl benzyl ammonium-grafted ladder polymers 5
- triphenyl carbocations 174
- 4,4'-tris-methylene bis(1-methyl-piperidine) 70
- tunable multication-crosslinked poly(biphenyl piperidinium) (PBP-xQ4) membranes 190
- u**
- UiO-66-NH<sub>2</sub> membrane 266–267
- UiO-66-SO<sub>3</sub>H membrane 268, 269
- UiO-66(Zr/Ti)-NH<sub>2</sub>/polyamide mixed matrix membrane 268–271
- w**
- water electrolysis 139, 151, 155, 203, 206, 207, 215, 358, 384–386
- water migration 352, 353, 371
- water transport 32–34, 353
- water treatment membrane 106
- water uptake and swelling ratio 40, 199
- wide angle X-ray diffraction (WAXD) 46
- Wittig reaction 69, 82
- z**
- zeta potential 35–37, 280, 299, 312, 313
- Ziegler–Natta catalyst 73