

Index

a

- abstract shape analysis
 - applications 583–584
 - comparative
 - from aligned sequences 587–588
 - from unaligned sequences 588–592
 - definition 580–582
 - functions 580
 - general caveats 582–583
 - probabilistic shape analysis 585–587
 - RNA folding space 579–580
 - RNAlshapes parameter 593
 - RNAsHapes parameter 592–593
 - shape representative structures computing 581
- acetal levulinyl ester (ALE) 130
- ADE2 gene 1071, 1080
- ADH1 promoter 1073
- 3'-adapter
 - ligation 862–863, 889–890
 - preadenylated, for Illumina sequencing 864–865
- 5'-adapter
 - ligation 869, 890–891
 - oligoribonucleotide, compatible with Illumina sequencing 865
- Advanced Photon Source (APS) 409, 411–412
- affinity matrices, for RNA binding proteins
 - isolation 921–924, 948–952
 - applications
 - nuclear tRNase Z purification from wheat germ 927–930
 - tRNA-splicing ligase purification from wheat germ 930
 - materials 924–925
 - methods 925–927
- affinity purification. *See* chromatography
- affinity-selected material analysis 943–944
- Affymetrix Integrated Genome Browser 724
- AFM. *See* atomic force microscopy imaging
- Agilent eArray system 796
- Ago2 immunoprecipitation, Western blot analysis 1090
- ALIFOLDZ algorithm 727
- aliphatic isocyanates synthesis 160–161
- altritol 1251
- AMBER (Assisted Model Building with Energy Refinement) program, MD simulation 687, 695
- 2'-amino approach
 - with aromatic isothiocyanates 158–159
 - postsynthetic modification 156–158
 - with aliphatic isocyanates 159
 - with succinimidyl esters 158
- 2'-amino-modified RNA postsynthetic labeling 161–164
- aminoglycosides 493–494
 - as functional and structural probes 401–403
- amino-propyltriethoxy silane (APTES) 534
- analytical crosslinking 246–247
- anion-exchange. *See* chromatography
- antibodies testing, for immunoprecipitation 882
- antifading agents 510–520
- antipeptide antibodies generation 958–962
- antisense (AS)
 - accessible sites in target RNA/RNP, by SELEX 1165–1166, 1169–1170, 1180–1181
 - binders, in SELEX 1100
 - *cis*-AS RNAs 794
 - kinetics of AS RNA:target RNA pairing
 - low-abundance *cis*-antisense transcripts to protein-coding genes 728

- antisense (AS) (*contd.*)
 - ncRNA sequence alignment 789
 - Northern blot probes 729, 747
 - oligonucleotide, phosphorothioate-modified 1246
 - primer extension on RNA 774–776
 - primer PCR 38–39, 755, 757–758, 760–761, 763, 768, 772, 1173–1174, 1177–1178
 - regulation by sRNAs 739, 743
 - – RybB-*ompC* interaction 767
 - RNA elements favorable for AS pairing 740
 - rRNA capture by AS oligonucleotides 722
 - sRNA-mRNA interaction 719–720
 - strand, e-shRNA, 1230
 - strand, lhRNA 1226–1227, 1230
 - strand, shRNA 1225–1227
 - strand, siRNA 1243–1245, 1249–1251, 1253–1255, 1263–1264
 - strand of genome, microarrays 720
 - strand transcription 1169
 - to miRNAs, oligonucleotides 837
 - to protein-coding genes, long ncRNAs 814
 - to RNA/RNP, biotinylated oligonucleotides 178–180, 936–948, 957
 - – displacement DNA oligonucleotide 940, 945–948
 - true and false AS transcripts 795
- APART (Automated Pipeline for Analysis of RNA Transcripts) 1194–1196
- aptamers 735–736
- aptamer selection, against biological macromolecules 1097–1098
 - general strategy 1098–1100
 - – carbohydrate targets 1101–1102
 - – library design and preparation 1103–1104
 - – protein targets 1100–1101
 - – selection arrays 1103
 - – suitable target choosing 1100–1102
 - – target immobilization 1102–1103
 - *in vitro* selection cycle 1104–1106
 - protocols 1108–1131
 - selection outcome analysis 1106
- 3D architectures 667
 - S2S and Assemble2 668–673
 - – installation 673–684
 - – semi-automatic architectures 672–673
- aromatic isothiocyanates 158–159
- arylazide photocrosslinking agent attachment, to 5'-terminal phosphorothioate 236–238
 - 3'-addition of arylazide photocrosslinking agent 238–240
- ASH1 gene 1080, 1082
- Assemble2 671–672
 - S2S 668–671
 - – installation 673–684
 - – semi-automatic architectures 672–673
- atomic force microscopy (AFM) imaging 527
 - example protocol 537–538
 - force spectroscopy 540–543
 - of RNA structures
 - – experiment example for salt-dependent RNA folding 535–537
 - – general preconditions 528–531
 - – imaging in liquid 535
 - – surface preparation conditions 531, 533–535
- b**
- bacterial aging 743
- bait 1067–1068, 1073, 1078, 1188
- barcoded cDNA libraries for miRNA profiling, by next-generation sequencing 861–862
 - method overview 862–863
 - – materials 863–866
 - – procedure 866–872
- base-labile protecting groups deprotection 137
- BED sequence alignment format 823, 826
- bimolecular transitions 452
 - absorbance parameters temperature dependence 455
 - basic equations about melting curves 454
 - entropic considerations 452–453
 - fitting or shape method 456–457
 - half-width method 456
 - higher order transitions 455
 - results 457–458
 - slope method 455–456
- binning 505–506
- 3'-biotinylated deoxyoligonucleotide 178–179
- biochemical studies and RNP affinity selection 941–945
- biotin attachment 120–121
- biotin-based affinity purification of RNA-protein complexes 935–937
 - affinity matrices 937–938
 - biotinylated probes 937
 - buffers and solutions 938–939
 - cell extracts 938
 - methods
 - – nuclear extract depletion 951–952
 - – RNA-protein complexes affinity purification 939–948

- specific RNA binding proteins affinity purification 948–951
- BLASTN program 598, 600–602, 623–624, 736, 808
- BLAST program 596, 598, 620, 625, 627, 726, 788–789
- Boltzmann weighting 433
- BOWTIE sequence read alignment program 823, 826, 1196–1197
- breathing 449
- burst mode (smFRET) 500
- BWA sequence read alignment program 823, 826
- C**
- calf intestinal alkaline phosphatase (CIAP) 56–57, 172
- CAN1 gene 1074
- cantilever 528–529, 535, 538, 540–542
- capillary electrophoresis (CE) 206–207, 220–221, 265
- carbohydrate targets 1101–1102
- CASP (Critical Assessment of techniques to protein Structure Prediction) 702
- catalytic RNA 29, 33, 261, 304, 308
- cDNA library of ncRNAs. *See also* barcoded cDNA libraries for miRNA profiling, by nex-generation sequencing
- computational analysis of ncRNA sequences 811, 816
- construction and screening 804–808
- exclusion of cDNAs for abundant known RNA species 808
- Northern blot analysis 808
- sequence analysis 807–808
- specialized library 808–811
- RNA sample enriched by immunoprecipitation 809
- RT-PCR procedure with specialized primers 809–811
- PAR-CLIP 882–883
- preparation from recovered RNA fragments 889–893
- RNP-derived construction 814
- RNA preparation and construction of library 815–816
- RNP extracts and glycerol gradient centrifugation preparation 814–815
- cell lysate/extract
- affinity purification of RNA binding protein with immobilized RNA 735
- affinity purification of snRNPs and CA-repeat RNA binding factor with biotinylated antisense oligonucleotides 940–952
- β -galactosidase activity assay 1078
- crosslinking of RNA 241
- endogenous RNase activity 902
- EMSA 981
- immunoprecipitation/immunoaffinity chromatography of RNA binding proteins and RNPs 805, 882, 884, 986, 988–990, 1020–1021, 1090, 1093
- formaldehyde-crosslinked 1017–1018, 1021–1022
- isolation of proteins associated with *in vivo* expressed tagged RNAs 736, 963, 966, 969
- magnetic beads for RNP affinity purification 938
- preparation 938
- ribosome-free (S100) 176
- cell-penetrating peptides (CPPs) 1260
- CHARMM (Chemistry at HARvard Molecular Mechanics), MD simulation 688, 695
- chemical RNA synthesis, purification, and analysis 129–132
- deprotection 136–138
- purification 138–139
- anion-exchange HPLC purification 139–140
- desalting by HPLC 142–143
- trityl-on RNA detritylation 142
- trityl-on RNA reversed-phase HPLC purification 140–142
- purified RNA analysis 143–144
- RNA solid-phase synthesis 132–136
- automated RNA synthesis 136
- manual RNA synthesis 134–136
- chemical sequencing 187–189
- chromatography
- affinity 921, 1191
- biotinylated antisense 2'-O-methyl RNA oligonucleotides to select RNPs *via* immobilization to streptavidin agarose 941–943, 976
- 3'-biotinylated DNA oligonucleotides to affinity-purify RNAs *via* immobilization to streptavidin agarose 178–180
- Cibacron Blue, HighTrap Blue 927–931
- heparin 927–931
- immunoaffinity using anti-peptide antibodies specific to snRNPs 957–962, 976

- chromatography (*contd.*)
- – immunoaffinity using 5'-cap-specific antibodies 986
 - – Ni-NTA agarose, proteins 1110–1111
 - – purification of RNA binding proteins (depletion of cell extracts) *via* biotinylated RNA and streptavidin agarose 948–952
 - – RNA aptamer-based affinity purification 963–971, 976, 986–993, 1006, 1008
 - – SELEX, covalent target immobilization 1140–1146
 - – SELEX, non-covalent target immobilization, biotinylated target captured *via* (strept)avidin matrix 1147
 - – tRNA-sepharose 927–931
 - fast-performance liquid chromatography (FPLC) 21–23, 107, 112, 482, 976
 - flash column (CH₂Cl₂), synthesis of aromatic isothiocyanates and aliphatic isocyanates 160
 - gel (exclusion, permeation) chromatography, gel filtration 114–115, 121, 213, 320, 509, 927–929, 1110
 - – NAP columns 19, 80, 142, 146, 512
 - – Sephadex G-25 19, 142, 744, 1046, 1091
 - – Sephadex G-50 122, 323, 350, 403, 537, 1046
 - – Sephadex G-75 122
 - – spin column 122, 323, 350, 402–403, 744, 1046, 1091
 - – Superdex 200 930–931
 - ion-exchange 235, 921, 930, 932, 1110
 - – weak anion-exchange, diethylaminoethyl (DEAE) sepharose 18, 105, 107, 111–115
 - – DEAE, snRNP enrichment 938, 941, 945
 - – strong anion-exchange 138–140, 928
 - – strong cation exchange 22–23, 930–931
 - – MonoQ 213, 976
 - isocratic competitive affinity chromatography 1161–1162
 - high-performance liquid chromatography (HPLC) 18, 132, 138–147, 162–166, 195, 290–291, 361, 471, 948, 1098, 1108, 1110, 1143
 - – Dowex 50 cation-exchange, conversion of sodium to ammonium salt of RNA 146–147
 - – oligoribonucleotide desalting with Sephadex G-25 142–143
 - – separation of peptides 929–930
 - HPLC, anion-exchange 132, 138–140, 142–144, 146
 - – DEAE 235–237
 - – Dionex 471, 482–483
 - HPLC, reversed-phase 132, 138, 140–145, 147, 162, 164–165, 512
 - – analysis of ribonucleosides 893–896
 - – of trityl-on RNA 138, 140–142
 - – phosphorothioate stereoisomers 290–291
 - ion-pair reversed-phase liquid chromatography (IP-RPLC) 142–143, 213
 - thin-layer (TLC)
 - cellulose acetate strips/DEAE cellulose 175, 190–194
 - cellulose 195–197, 200–201
 - polyethyleneimine (PEI) 196, 201, 236
 - ultra performance liquid chromatography (UPLC) 143–144
 - circular potato spindle tuber viroid (cPSTVd) 434–435
 - cleavage reactions by Tb³⁺ 256–258
 - click reaction 509
 - between alkyne-containing RNA oligonucleotide and azide-functionalized fluorophore 511–512
 - CLIP (crosslinking-immunoprecipitation) 899, 1088, 1199
 - cloverleaf structure 76
 - CLUSTALW program 598, 620
 - COLORES program (SAXS) 418–419
 - comparative genomics 836–837
 - concatenation method 737
 - concentration calculation, of RNA 60, 1279–1281
 - Cordycepin 182–183, 337–339. *See* radioactive labeling methods
 - Coulombic field 319, 322
 - coupling chemistries 119, 121–122
 - covalent coupling of antibodies, to protein A sepharose 1047–1048
 - covalent immobilization
 - epoxy-activated matrices 1143–1145
 - NHS-activated matrices 1145
 - pyridyl disulfide-activated matrices 1146
 - covariance model (CM). *See* INFERNAL
 - cross-correlation function (CCF) 419
 - crosslinking 8, 50, 1199. *See also* PAR-CLIP
 - RNA-RNA crosslinking
 - – disulfide crosslinking 156–160
 - – photocrosslinking for probing of RNA structure 231–233, 236, 238–244, 246–249
 - RNA-matrix crosslinking 1048

- EDC [1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide] crosslinking 89, 91–92, 95–97, 100, 1091, 1094
- RNA-protein crosslinking 736, 977
- formaldehyde crosslinking 1017–1019, 1021, 1026
- iCLIP 899–902, 904, 908
- UV 91, 363, 746, 883, 899–901, 905, 999, 1003–1005, 1008–1009, 1026, 1029, 1030–1038, 1040–1042, 1191
- crude transcription reaction 105, 107, 114
- cryo-electron microscopy (cryo-EM) structures 690
- crystallization of RNA 481–482
 - assays evaluation 488–489
 - complexes with organic ligands 493–494
 - optimization process 489, 491
 - purification 482
 - gel electrophoresis 483–484
 - HPLC 482–483
 - recovery 484
 - renaturation 485
 - RNA constructs designing, with improved crystallization capabilities 491–493
 - search for conditions 485–488
- CURVES+ program (conformational analysis of nucleic acids) 688, 703
- d**
- DAMAVAR/DAMFILT/DAMMIF/DAMMIN programs (SAXS analysis) 416–417
- DAP Goldstar® Polymerase (Eurogentec) 1120
- Dashboard 633, 636–639, 641, 650, 652, 657, 659, 661–663
- Debye–Hückel approximation 462
- decoy 636, 659–661, 1082, 1235
- deep sequencing. *See* next-generation sequencing (NGS)
- dephosphorylation protocols 180
 - DNA 5′-ends 110
 - ribonucleoside preparation for HPLC analysis 895
 - RNA 5′-ends 64, 181, 337–338, 350, 885, 980
 - RNA 3′-ends 39–40, 56–57, 121, 909
- deprotection 136–138
- desalting, by HPLC 142–143
- detritylation 132, 138, 141–142, 145
- Diels-Alderase ribozyme 501
- diethylaminoethyl (DEAE) 107, 112–114, 176, 190–192, 235, 938, 941–943, 945, 1127
- diethyl pyrocarbonate (DEPC). *See also* probing RNA structure
 - RNase-free water preparation 20, 59, 305, 309–310, 313, 744, 803, 1107, 1173
- differential melting curve (DMC) 448, 450, 451–452, 454, 456–457, 471–474
- differential scanning calorimetry (DSC) 453–454, 457–458, 462, 472
- diffusion elution 19
- 5′-digoxigenin (DIG)-labeled probes 89, 91, 946,
 - detection 95, 101–102
 - DIG Easy Hyb Granules 97, 101,
 - probe generation by T7 transcription using DIG-11-UTP 89, 92, 100–101
- dimethylsulfate (DMS) modification. *See* probing RNA structure
- divalent metal ion binding sites, in nucleic acid-metabolizing enzymes 397–398
 - probing 398
 - aminoglycosides as functional and structural probes 401–403
 - Fe(II)-mediated hydroxyl radical cleavage 398–399
 - mapping 399–401
- DNAMAN sequence analysis program 811
- DNase treatment 18, 53, 59–60, 67, 80, 83, 213, 278, 745, 752, 769, 908, 980, 1009, 1111–1112, 1153, 1167–1168, 1172, 1176, 1179, 1190, 1216–1217
- 3DNA program (nucleic acid structure analysis, rebuilding, and visualization) 688, 703
- DNA splint 49–52, 57–59, 66–70, 75, 78, 80, 82–83, 286, 291–293, 338, 362
- DNazyme 35, 380, 1166
- donor substrate. *See* ligation
- dot plot folding comparisons 565
- double-stranded RNA (dsRNA) 527, 530, 534, 537, 1221, 1254, 1256–1257
- dRNA-seq approach 722–724, 732
- dynamic programming 222, 552, 554–555, 572, 586, 604
- e**
- EDC crosslinking. *See* crosslinking
- Edman degradation 929
- electroelution 19, 41, 223, 886, 888, 897, 1178
- electron paramagnetic resonance (EPR) 319
 - active probes 156
 - spin labeling reagent 4-isocyanato TEMPO 157
 - spin probe/label TEMPO 159

- electrophoretic mobility shift assay (EMSA)
 - 924, 976–978, 981–985, 996, 998, 1005, 1010, 1067, 1103, 1106, 1124–1128, 1199
 - method 977–978, 981–986
 - protein identification in RNP 984–986
 - supershift method 983–984
- Elutrap[®] 19
- emacs (text editor system) 621, 625
- EMSA. *See* electrophoretic mobility shift assay
- end heterogeneity of RNA 6, 29, 45, 54, 57, 84, 121, 223, 338
- end-labeling of RNA 257, 346, 1130. *See also* radioactive labeling methods
 - at 3'-end 33, 73, 181–182, 213, 378–379, 382–383, 385, 389, 980
 - – with fluorescent dye 121, 509
 - – with biotin 119–121
 - at 5'-end 5–7, 10–11, 32, 92, 180–181, 213, 380–381, 383, 389, 774, 883, 886, 980, 995
 - – with fluorescent 2-aminopurine 73
 - – with 3'-amino ATP (for biotinylation) 73
 - – with 5'-biotin-GMP 7
 - – with GMPs 7, 233–234
 - – with 6-thio-GMP 7, 235, 242
- 3'-ends 6, 33, 53–54, 182–183
 - addition of arylazide photocrosslinking agent 238–240
 - generation by autocatalytic ribozyme cassettes
 - cassettes construction 36–37
 - *cis*-cleaving 30–31
 - dephosphorylation protocols 39–40
 - PCR protocols 37–39
 - removal of 2',3' cyclic phosphate 32–33
 - *trans*-cleaving 33
 - generation by RNase P 33–35
 - protocol for RNase P cleavage 40–41
 - RNase H approach to generate homogeneous transcript 3'-ends 54–56
- 5' ends 5–6, 35, 183–184, 195, 199
 - generation by *cis*-cleaving autocatalytic ribozyme cassettes 30–31
 - cassettes construction 36–37
 - modification by transcription priming 6–7, 11–12, 17–18, 51, 61, 233–234
 - – 5'-monophosphorylation of nucleosides 234–236
 - PCR protocols 37–39
 - radioactive labeling, of RNA termini 64–65, 180–183, 337–339, 909–910
- endogenous RNAs associated with RNA binding proteins, immunopurification 1017
 - methods 1017–1022
 - microarray analysis of immunopurified RNA 1022–1024
 - – next-generation sequencing of immunopurified RNA 1025
 - – RT-PCR analysis of immunopurified RNA 1024
 - – critical points and common problems 1025–1026
 - – microarray data analysis 1026–1027
 - – uncrosslinked or crosslinked RNA immunoprecipitation 1026
- endothermic process 451
- energy dot plot 558–561, 563, 567
- enhanced green fluorescent protein (eGFP) 1234
- ENSEMBL database and genome browser 620, 623–624, 629, 816, 851
 - bacterial genome browser 602
- entropic considerations 452–453
- enzymatic hydrolysis 184–187, 191, 224, 226, 895
- enzymatic probing 205–206, 214–215, 224, 243, 435, 462, 602, 634–635, 998
- enzymatics 1150–1151
 - *in vitro* transcription 1153
 - polymerase chain reaction 1152–1153
 - reverse transcription 1151–1152
- epoxy-activated matrices 1143–1145
- EPR. *See* electron paramagnetic resonance
- equilibrium dialysis 320, 323–325, 1159–1160
- equilibrium filtration analysis 1160–1161
- ERPIN program (RNA motif identification) 597–599, 609–616
- ethanol (EtOH) precipitation
 - protocols for DNA 1176, 1215, 1218
 - protocols for RNA 19, 61–62, 98, 120, 142, 162, 177, 213–214, 219–221, 223, 226, 248, 258, 278, 306, 309–310, 336, 350, 355, 381–382, 385, 752, 769, 775–776, 806, 810, 867–868, 871–872, 888, 890, 897, 944, 962, 971, 998, 1021, 1044, 1060–1061, 1064–1065, 1114, 1217
- EUROSCARF (*Saccharomyces cerevisiae* database, strain and plasmid collection) 1030
- EvoFold (RNA structure identification in multiple-sequence alignments) 572, 788–789
- exonic splicing enhancers (ESEs) 1207–1208, 1210, 1212
- exonic splicing silencers (ESSs) 1207–1208

exothermic process 451
 extended shRNA (e-shRNA) 1224,
 1226–1227, 1229–1230, 1233

f

FASTA (format) 556–557, 566, 598,
 600–602, 608, 611–612, 616, 623, 625, 650,
 674–675, 678–679, 681
 fast-performance liquid chromatography
 (FPLC). *See* chromatography
 Fe(II)-mediated hydroxyl radical cleavage
 301–312, 397–401. *See also* Fenton
 reaction/cleavage
 FeBABE reagent 302
 Fenton chemistry 308, 314
 Fenton cleavage
 – mapping divalent metal ion binding sites in
 proteins 397–401
 – – methods description 402–404
 – – aminoglycosides as functional and
 structural probes 401–402
 – of RNA 301, 303, 310, 312–313
 – – methods description 305–312
 – – drug-directed (tetracycline) 310
 Fenton footprinting 304
 Fenton reaction 301, 306, 398
 flow cytometry 738, 756–757, 764
 – protocol 764–766
 fluorescence resonance energy transfer
 (FRET) 50, 66, 68, 263, 499–503
 – dye pairs 504
 – – choice of 507–508
 – initial spectrometric characterization by
 bulk measurements 513–514
 – smFRET. *See* single molecule fluorescence
 resonance energy transfer
 fluorescent labeling 121, 123, 796–797
 5-fluoroorotic acid (5-FOA) 1074, 1081
 footprinting 207, 214, 219, 225, 255–257,
 259–261, 302, 304, 636, 660, 662, 739,
 977–978, 981, 991, 1067, 1106, 1199.
See also terbium(III) footprinting
 – semi-automated footprinting analysis
 (SAFA) software 221
 forced dialysis 320–324, 326
 force spectroscopy AFM, of RNA 527,
 540–543
 Förster radius 502–504, 507
 Freiburg RNA Tools 620
 FRET. *See* fluorescence resonance energy
 transfer
 Friedel law 472
 fusion protein (FP)
 – MS2-MBP 957, 966, 969, 989–990, 992

– in the yeast three-hybrid system 1067
 – – FP1 with DNA binding domain
 1068–1070
 – – FP2 with transcriptional activation
 domain 1068–1070
 – – FP1, LexA fused to MS2 coat protein
 1069, 1073
 – – FP2, activation domain of yeast Gal4 fused
 to RNA binding tester protein
 1069–1070, 1073–1074

g

GAL4 activation 1069–1070, 1073–1075,
 1077–1078
 β -galactosidase 738, 1076, 1078–1079, 1081
 – filter assay 1079
 gel electrophoresis, of DNA 220, 889, 902,
 916, 1108, 1120, 1153
 gel electrophoresis, of RNA 722, 1192
 – denaturing polyacrylamide (denaturing
 PAGE) 6, 15, 18–20, 29, 31, 39, 41, 53,
 59, 61–64, 67, 69–70, 80–84, 92, 105,
 113–114, 123–124, 142–143, 162–164,
 183, 207, 212–213, 216, 223, 226,
 244–245, 248, 257, 261, 292–294, 306,
 324, 372, 377, 379–384, 389, 392, 422,
 471, 483–484, 495, 520–521, 537, 740,
 768, 770, 774, 862, 899–901, 942, 944,
 961, 967–968, 980, 992, 1005, 1036, 1047,
 1059–1060, 1063–1064, 1112, 1123–1124,
 1130, 1153, 1187
 – native polyacrylamide
 (native/non-denaturing PAGE) 18, 31,
 99, 224, 344, 352, 481, 948, 982–986,
 1123–1126, 1181
 – temperature-gradient (TGGE) 427–433,
 462
 – two-dimensional 176, 192–193
 gel exclusion (permeation) chromatography,
 gel filtration 114–115, 121, 213, 320, 509,
 927–929, 1110
 – NAP columns 19, 80, 142, 146, 512
 – Sephadex G-25 19, 142, 744, 1046, 1091
 – Sephadex G-50 122, 323, 350, 403, 537,
 1046
 – Sephadex G-75 122
 – spin column 122, 323, 350, 402–403, 744,
 1046, 1091
 – Superdex 200 930–931
 gel fractionation 221, 996
 gel purification, of RNA 15, 21, 52, 56, 65,
 122–123, 242–243, 247, 257, 331, 361, 379,
 482, 740, 770, 914, 969, 1176
 – preparative 84

- gel purification, of RNA (*contd.*)
 - kit 751, 880
- gel running buffer
 - 1 x TBE 19
 - 5 x TBE 59
- GenBank 433, 556, 560, 563, 566, 569, 571, 595, 619, 622, 624, 808, 821
- gene expression profiling computational methods. *See* next-generation sequencing (NGS)
- general buffers and procedures (GBP) 59
- gene silencing methods 1221
 - background information 1221–1223
 - extended shRNA (e-shRNA) and long hairpin RNAs (lhRNAs), construction 1229–1230
 - lentiviral vectors, production 1230–1234
 - miRNA vectors, construction 1228–1229
 - shRNA vectors, construction 1223–1228
- genome(s) 625–628, 667, 719–720, 731, 736, 787–791, 794, 811, 822, 829, 1165–1166, 1168, 1185–1186, 1193, 1202
 - annotation 628, 787, 794, 1197, 1223
 - APART pipeline 1196–1198
 - *Arabidopsis thaliana* 929
 - assembly 623–624, 628, 829
 - avian 626, 628
 - bacterial 595, 600, 602, 616, 726–727, 787, 790–791, 1187
 - BLAST search 623
 - browser 602, 620, 623, 625, 628, 722–724, 823, 826, 829
 - – Affimetrix integrated 724
 - – UCSC 620, 625, 823, 826, 829
 - comparative analysis 721, 787–788
 - coverage 1188
 - coverage by tiling microarray probes 795–796
 - database 596, 599, 608, 614, 624, 726
 - encoded aptamers and regulatory sequences 1185, 1199
 - eukaryotic projects 620
 - gene copies 622
 - HIV-1 206
 - human, mammalian 623–625, 627–628, 801, 849, 864, 877, 1168, 1170, 1187
 - Illumina Genome Analyzer 902, 904, 916
 - invertebrate 623
 - lizard 625
 - low complexity region 598
 - mitochondrial 583
 - mRNA profiling on microarrays, genome-wide 729, 733
 - ncRNA prediction, genome-wide 787–791
 - NGS read alignment 826–827
 - of lentiviral vector 1232–1233
 - organellar 620
 - phage T4 46
 - phage T7 4
 - probe complementarity 93
 - PSTVd 434
 - (nc)RNA cDNA read mapping 816, 916, 1195–1197
 - SELEX (genomic) 1185
 - sequence(s) 175, 595, 604, 624, 726, 1185
 - sequencing 1081
 - shuffled database 609
 - specific primer 1187
 - transcriptome analysis 732–733, 825–826
 - vertebrate 620, 623
 - wallaby 627
 - whole genome oligo microarrays, mouse 1023
 - yeast 1069, 1187
- genome mapping, of cDNA reads 723, 1195–1197
- genome screening, for RNA genes/motifs 595–596
 - search for bacterial sRNA homologs 600
 - – alignment and structure prediction 602–604
 - – BLASTN 600–602
 - – ERPIN 609–614
 - – HMMER 604–605
 - – INFERNAL 614–615
 - – RNAMOTIF 606–609
 - RNA search procedure overview 597–598
 - search program choice 596–597
 - search specificity assessment 598–600
- genomic aptamer(s) 1185–1186, 1188, 1192, 1194, 1196
 - biochemical analysis 1199–1202
 - – RNA-protein interaction validation 1199
 - expression analysis 1199–1200
 - size of native transcript 1200
 - function of RNA-protein interaction 1200
- genomic SELEX 720, 728, 1165–1166, 1171, 1185–1186
 - bait choice 1188
 - computational analysis of sequences 1194–1198
 - library construction 1186–1188
 - procedure 1188–1194
 - specialized SELEX method to identify antisense and protein target sites in RNA or hnRNPs 1165–1166

- generation of RNA 20-mer library using *Mme* I 1166–1169
 - natural RNA substrates for proteins, identification 1171
 - optimal antisense sites identification 1169–1170
 - procedure and protocols 1171–1182
 - glyoxy-oxygen-scavenging enzyme stock 515–516
 - glycerol gradient (centrifugation) 802, 813–816, 940, 948, 960–962, 964, 966–968, 970, 976, 986, 1056–1061
 - graph theory 552
 - green fluorescent protein (GFP) 732, 1228–1229
 - affinity tag 1017–1019
 - anti-GFP antibody 768–769, 771, 1018, 1021
 - enhanced GFP (eGFP) 1234
 - flow cytometry 764–767
 - GFP-based reporter system for sRNA target validation 755–757
 - lentiviral vector encoding Emerald Green Fluorescent Protein (EmGFP) 1228–1229
 - read out approaches 763–764
 - translational GFP fusion, cloning 759
 - pXG-10 759–761
 - pXG-20 761–762
 - pXG-30 762–763
 - GROMACS package, MD simulation 688, 695
 - GROMOS (GROningen MOlecular Simulation computer program package) 688, 695
 - GSNAP sequence read alignment program 823, 826
 - guanosine 5'-monophosphorothioate (GMPS) 7–8, 233–238
 - guide strand. *See* small interfering RNA (siRNA)
 - Gwyddion software (AFM) 530
- h**
- hammerhead ribozyme 6, 30–33, 36–39, 75, 79–80, 121, 123–124, 152, 158, 264, 286–287, 337, 374, 500
 - HaMMY software package, smFRET analysis 507
 - hepatitis delta virus (HDV) ribozyme 6, 30–39, 261–264, 337
 - heterogeneous nuclear ribonucleoprotein (hnRNP) 900, 950–952, 975–976, 984–986, 998–1000, 1003–1007, 1055, 1072, 1165–1182
 - Hfq 719–720, 725, 728–729, 731, 734–738, 740–743, 770–771, 1197
 - Hfq coimmunoprecipitation. *See* immunoprecipitation
 - Hidden Markov Model (HMM) 507, 604–605
 - high-performance liquid chromatography (HPLC). *See* chromatography
 - high-throughput RNomics 794–799
 - high-throughput sequencing (HTS) 89, 206, 720–723, 742, 802, 812, 877, 899, 901–902, 904, 916, 1106, 1186, 1194. *See also* RNA-seq
 - HTSeq, high-throughput sequencing data analysis software 823, 829
 - Hill equation 297
 - HIS3* gene 1069, 1075, 1078, 1080
 - histidine-tag (His-tag) 56, 372–373, 865, 1030, 1050, 1128–1129
 - HMMER program (search for RNA homologs) 596–599, 604–605, 614–616
 - holoenzyme 289, 292, 294, 326, 375, 387, 492
 - reaction 294
 - homogeneous 5'- and 3'-ends. *See* T7 RNA polymerase
 - homology-based programs 596–597
 - homology search 596–597, 602, 604, 811
 - for small structured ncRNAs 619–621
 - mascRNA example 621–629
 - initial BLAST search 622–624
 - initial secondary structure model 624–625
 - structure-based searches 625–629
 - web-service-independent software 621
 - web services 620
 - RsaE RNA test case 600
 - alignment and secondary structure prediction using LOCARNA 602–604
 - BLASTN first training set 600–602
 - descriptor- vs. homology-based programs 596–597
 - RNA motif search specificity and protocol 598–600
 - RNA motif search with sequence and secondary structure constraints 595
 - searching with ERPIN 609–614
 - searching with INFERNAL, covariance models 614–615
 - searching with HMMER 604–605
 - searching with RNAMOTIF 606–609
 - program versions and download sites 616
 - Hooke's law 528

- hot phenol method 90, 94
 - RNA extraction from cells 98
- HQS. *See* 8-hydroxyquinoline-5-sulfonic acid
- HTSeq, high-throughput sequencing data analysis software 823, 829
- hybridization 117, 352, 527, 555, 648, 650, 836, 871, 936, 1000, 1018–1019, 1248
 - cross-hybridization 827, 861
 - experimental RNomics 803, 807–808, 816
 - identification of small non-coding RNAs 725, 729, 737, 744, 746–749, 768
 - immunopurification 1018–1019, 1022–1023
 - microarray hybridization 796–799
 - Northern blot 89–92, 94, 96–102, 794, 1091–1092, 1094
 - oligonucleotides 179, 427, 434–435, 437–438, 952
 - primer extension 220, 223, 226, 280, 310, 995, 1046
 - to DNA splint 58, 67–68
- hybrid RNA 1068–1075, 1077–1081
- hydroxyl radicals 206, 209, 225, 301–302, 304, 313–314, 398, 421, 740, 1007. *See also* Fenton chemistry/cleavage/footprinting/reaction
- 8-hydroxyquinoline-5-sulfonic acid (HQS) 320, 324–325, 327
- hyperchromism 446, 449
- hypochromism 446, 449, 464
- i**
- iCLIP (individual-nucleotide resolution crosslinking and immunoprecipitation. *See* crosslinking)
- ImageJ software 530, 536, 744, 747, 774
- ImageSXM software (AFM) 530
- immunoprecipitation 725, 805, 809, 840, 963, 1008–1009, 1020–1027, 1030, 1037–1038, 1040–1044, 1046–1048, 1171
 - coimmunoprecipitation 720, 1073, 1075, 1119
 - – Hfq coimmunoprecipitation 722–725, 728, 734
 - – RNA coimmunoprecipitation 1017, 1020–1027
 - iCLIP 899–902, 905–909
 - identification of miRNA targets 1087–1095
 - PAR-CLIP 877, 882, 884–887
 - RISC components 838–839
- INFERNAL program (structural RNA sequence alignment based on covariance models) 597–599, 614–616, 621, 626–628
- in-line probing 206, 210–211, 213, 216, 218, 225–226, 417, 1106
- inner-sphere coordination, cation : RNA 319, 370, 373
- interference suppression 330–332, 361, 363, 375, 378
- intergenic region (IGR) 719–720, 722, 724, 726–728, 748, 788, 790–791, 793–796, 1195
 - Intergenic Sequence Inspector (ISI) 726
- intermittent contact mode (tapping), AFM 529
- intron ribozyme activity (group II) 339–344
- in vitro* evolution 442, 1099–1100
- in vitro* probing. *See* probing
- in vitro* selection 16, 442, 1097, 1099–1101, 1104–1107, 1116. *See also* SELEX
 - against proteins and carbohydrates 1097–1131
 - against small targets: 1139–1162
 - selection cycle: 1099, 1104–1106, 1141
- in vivo* SELEX. *See* SELEX
- iodine cleavage (iodine-induced hydrolysis) 330–331, 341–342, 344, 346–347, 354–355, 385–386, 391–393
- ion-exchange chromatography. *See* chromatography
- isocratic competitive affinity chromatography. *See* chromatography
- isopycnic ultracentrifugation 1056, 1061. *See also* ultracentrifugation
- isopropanol precipitation
 - of plasmid DNA 110–111
 - of RNA 99, 752, 1157–1158, 1180, 1217
- isothermal titration calorimetry (ITC) 453
- k**
- Klenow polymerase (Klenow Pol/Klenow fragment/large fragment of *E. coli* DNA polymerase I)
 - 3′-³²P-end-labeling of RNA 337–338, 341
 - Fe(II)-mediated hydroxyl radical cleavage 397, 399–403
 - fill-in reaction, *in vitro* selection/SELEX 1149–1150, 1168, 1171, 1174–1176
 - genomic SELEX 1186–1188, 1193
- K-Means Clustering 661–662
- knockdown (KD) 837, 839, 851, 902, 908, 911, 922, 1200
- gene silencing 1221, 1223–1224, 1228
- chemical modifications to enhance siRNA performance 1243, 1247, 1252–1253, 1260, 1262
- “K-turn” (structure/motif) 501, 671, 975, 982

I

- 3'-labeling. *See* end-labeling of RNA
 - 5'-labeling. *See* end-labeling of RNA
 - lacZ* (gene) 735, 738–739, 756, 762, 1068–1070, 1075
 - lanthanide (ion) 255
 - lentivirus/lentiviral
 - vector production 1230–1235
 - LEU2* gene 1070, 1073, 1080–1081
 - Lewis
 - acid 270, 286, 370
 - base 286, 370
 - lhRNA. *See* long hairpin RNA
 - ligand coupling (*in vitro* selection) 1144–1146
 - ligation of RNA 6, 29, 33, 45–84, 152, 521, 801
 - acceptor substrate 46, 49, 70–73, 75–76, 78–80
 - donor substrate 46, 49, 70–73, 75, 78–80
 - ligation efficiency 46, 54, 70, 72, 75–76, 78, 82, 1169, 1172, 1179, 1214, 1216
 - ligation site accessibility 78
 - reaction mechanism 46, 48–49
 - simultaneous splint ligation of RNA fragments for FRET experiments 66–70
 - T4 DNA ligase (T4 Dnl) 47, 49–66. *See also* T4 DNA ligase
 - dephosphorylation and phosphorylation 56–57
 - homogeneous acceptor 3'-ends, generation 53–54
 - large-scale transcription and purification 53
 - RNA ligation 57–58
 - RNA ligation substrates generated with RNase H 54–57, 63–64
 - T4 polynucleotide ligase 46–47
 - T4 RNA ligase(s) 47, 70–84. *See also* T4 RNA ligase(s)
 - LOCARNA program (alignment and folding of RNA sequences) 573, 598–599, 602–604, 616, 620, 624–625
 - Locomotif tool (abstract shape analysis) 583
 - long hairpin RNA (lhRNA) 1224, 1226–1227, 1229–1230
- m**
- magnesium (Mg^{2+}) ions bound to RNA, quantification 319–320
 - Mg^{2+} complexes with 8-hydroxyquinoline-5-sulfonic acid (HQS) 320
 - separation of free Mg^{2+} from RNA-bound Mg^{2+} 320–324
 - equilibrium dialysis 323–324
 - forced dialysis 321–323
 - size exclusion chromatography 323–324
 - Mg^{2+} concentration determination using HQS 324–326
 - affinity and stoichiometry of Mg^{2+} binding 322–323
 - RNase P holoenzyme 326
 - magnetic bead(s)
 - aptamer selection 1099, 1116, 1120–1121
 - capture of rRNA using antisense oligonucleotides 722
 - immunoprecipitation
 - PAR-CLIP 882–886
 - iCLIP 900
 - miRNA targets 1088–1089, 1094
 - streptavidin-coated 937–938
 - MALAT-1 RNA/transcript 622, 625, 628
 - maltose-binding protein (MBP) 293, 736, 935, 957, 963, 966–970, 976, 987–993, 1006, 1008–1009
 - masCRNAs 621–629
 - BLAST search 623–624
 - secondary structure model 624–625
 - structure-based searches 625–629
 - mate-pair reads, next-generation sequencing 825
 - MC-Tools, RNA 2D and 3D structure prediction 633–634
 - equipment and input 634
 - MC-Cons 636, 651–654
 - MC-fold 635, 637–651
 - MC-Sym 636, 654, 657–661, 663
 - menRNA 627–629
 - metal ion coordination interactions with RNA
 - identification and characterization 285–286
 - thiophilic metal ion rescue of RNA phosphorothioate modifications 286–290
 - purification of phosphorothioate stereoisomers by RP-HPLC 290–291
 - phosphorothioates incorporation into RNA, techniques 291–293
 - kinetic analysis of thiophilic metal ion rescue 293–297
 - NAIM/NAIS analyses 329–332, 370–374
 - metal ion-induced cleavage/hydrolysis 256, 270–271, 275, 1104. *See also* probing
 - metalloenzyme 287, 397–398, 401
 - MICROExpress kit 722

- Micro Bio-Spin[®] chromatography column 122
- microRNA (miRNA) targets 833–836.
 - See also* barcoded cDNA libraries for miRNA profiling
 - comparative genomics 836–837
 - experimental identification 1087–1088
 - Ago2 immunoprecipitation, Western blot analysis 1090
 - cell culture and transfection 1089
 - gene expression array analysis of Ago2-associated mRNAs 1092–1093
 - Northern blot analysis for miRNA detection 1091–1092
 - qRT-PCR of Ago2-associated mRNAs 1092–1093
 - RNA precipitation 1090–1091
 - sample preparation and immunoprecipitation 1089–1090
 - functional binding site properties
 - data set type, quality and reproducibility 839–840
 - evolutionary conservation 841
 - miRNA-mRNA duplex stability 841
 - seed binding criterion 840–841
 - sequence composition 842
 - spatial effects 842–843
 - structural accessibility 841–842
 - targeting prediction 843–847
 - perturbation and omics 837–838
 - RISC components, immunoprecipitation 838–839
 - translation repression inferred from polysome profiles 839
- microRNA (miRNA) expression vector for gene silencing 1221–1223
- miRNA polycistron 1224
- vector construction 1226–1229
- minimum free energy (MFE), definition 549–555
- Mme* I restriction enzyme/site 1167–1170, 1173–1174
 - cleavage protocol 1178
 - digestion 1170, 1172
- mobility shift analysis
 - terminal RNA sequence identification 184, 190–194
 - electrophoretic mobility shift assay (EMSA) 166, 540, 899, 924, 976, 1067, 1103, 1124–1126, 1199
- modified nucleotides
 - incorporation by phosphoramidite chemistry 153–154, 359, 361
 - T7 transcription 6–8, 16–18, 61, 231–234
 - in NAIM/NAIS 333–336, 371–372
 - modified nucleotides determination, by postlabeling methods 194–201
 - position and identity determination 198–201
 - total nucleotide content analysis 195–198
 - modified nucleotide incorporation techniques 232–233
 - modified RNAs 151–152
 - 2'-amino approach 156–158
 - 2'-amino groups reaction
 - with aliphatic isocyanates 159
 - with aromatic isothiocyanates 158–159
 - with succinimidyl esters 158
 - experimental protocols 159–166
 - 2'-amino-modified RNA, postsynthetic labeling 161–164
 - 4-thiouridine-modified RNA, postsynthetic labeling 164
 - aromatic isothiocyanates and aliphatic isocyanates, synthesis 160–161
 - label incorporation verification 164–165
 - phosphoramidite method 152–154
 - postsynthetic labeling strategy 154–156
 - molecular barcodes 825
 - molecular dynamics (MD) simulations of RNA systems 304, 687–704
 - comparison with experimental data 702
 - conformational parameters 703
 - consistency checks 702
 - convergence issues 703
 - data analysis 704
 - validation based on structural databases 703
 - visualization 702
 - methods 689
 - setups
 - box size 693
 - clustering artifacts and ion parameters 696
 - conformational checks 690–691
 - divalent ions 694
 - duration of simulations 699–701
 - electrostatic interactions treatment 697
 - environment and ion types 693
 - equilibration 699
 - force fields 695
 - hydrogen atoms addition 693
 - initial solute and solvent configurations 694
 - minimal salt conditions 694
 - monovalent ions 693–694
 - multiple molecular dynamics (MMD) simulations 701

- parameterization of modified nucleotides, ligands, and ions 696
- programs 688, 695
- protonation issues 692
- rare non-covalent interactions 691–692
- shake, time steps, and non-bonded pair list update 698
- solvent 692–693
- starting structure selection 689–690
- temperature and pressure 698
- thermodynamic ensemble 697–698
- water models 696–697
- water molecules 694
- mRNA analysis
 - enzymatic and chemical probing 213–227
 - gene-expression-array analysis 1092–1093
 - qRT-PCR analysis 1092
- mRNAs (short) and dual-function RNAs 794
- MS2 aptamer/tag, for the phage MS2 coat protein 735–736, 1071–1073, 1075
- MS2-LexA fusion in three-hybrid analysis 1069–1070, 1080
- MS2-MBP affinity chromatography 736, 935, 957, 963–971, 976, 987–993, 1006, 1008–1009
- multi-FASTA format 598, 600
- multiple molecular dynamics (MMD) simulations 701
- n**
- NAIM. *See* nucleotide analog interference mapping
- NAIS. *See* nucleotide analog interference suppression
- NAMD program, MD simulation 418, 688, 695
- NcDAlign program (genome-wide alignments of non-protein-coding sequences) 788–789
- ncRNAs. *See* non-coding RNAs
- neutravidin 937, 946–947, 951
- next-generation sequencing (NGS). *See also* RNA-seq
 - cDNA library preparation 889–893
 - computational methods for gene expression profiling 821–822
 - association of reads with transcripts 827–828
 - data generation: experimental background information 823–825
 - expression determination and uncertainty 828
 - library generation 823–825
 - normalization 828
 - output and viewing 828–829
 - read alignment 826
 - sequencing 825
 - of immunopurified RNA 1025
 - associated with RBPs 1018
 - sequencing platforms
 - 454 Sequencing™ (Roche Diagnostics) 721–724, 812, 825, 1025
 - Solexa™ (Illumina) 721, 812, 822, 825, 861–862, 864–866, 872, 887, 889, 893, 902, 904, 916, 1025
 - SOLiD™ (ABI) 721, 807–808, 812, 1025
 - NF- κ B transcription factor 1073, 1082
 - N-hydroxysuccinimide (NHS)
 - acetal-PEG-NHS coupling to tip, AFM 542
 - activated sepharose matrices
 - tRNA coupling 922, 924–927, 929–930
 - tobramycin coupling 987–988
 - small ligand coupling for SELEX 1144–1145
 - biotin-NHS
 - for AFM 542
 - for SELEX experiments 1102, 1147
 - maleimide-PEG-NHS, AFM 543
 - esters, for RNA modification 67, 156, 508
 - nitrocellulose (NC)
 - filter assay 1079
 - filter binding 1101–1102, 1116, 1125–1127
 - membrane transfer 1090
 - iCLIP 900, 910–911
 - EMSA coupled to SDS-PAGE and Western blot 984–986
 - NMR (nuclear magnetic resonance spectroscopy)
 - isotopically labeled nucleotides, incorporation by *in vitro* transcription 34, 106
 - spectroscopic analysis 119, 164, 166, 286, 301, 407
 - ^{19}F NMR of nucleic acids 165
 - of mutant RNAs 439
 - RNA analysis and purity 29, 132, 143
 - RNA-metal ion interactions 304
 - structural dynamics of RNA 499
 - structure(s)
 - RNA 211, 263, 503
 - RNA/protein 409
 - in MD simulations of RNA, 688–690, 692
 - non-canonical base pairs 596
 - in NAIM analyses 340
 - in RNA structure prediction 560, 633–635, 645–648

- non-canonical base pairs (*contd.*)
 - in sRNA-mRNA interaction 736
 - in tRNAs 34
 - in UV spectroscopy 446
- non-coding RNAs (ncRNAs) 595, 667, 671, 1197. *See also* “small non-coding RNA (sRNA)” and “homology search”
 - characterization and prediction of miRNA targets 833–834
 - dual-function RNAs 794
 - experimental identification of miRNA targets 1087–1088
 - experimental RNomics 794–799, 801–802
 - cDNA library construction and analysis 804–811
 - computational analysis 811, 816
 - expression verification of cloned ncRNAs 808
 - immunoprecipitation of ncRNA-protein complexes 809
 - laboratory protocols 744–776, 797–799, 804–811, 813–816
 - identification and characterization in bacteria 719–720, 787–788
 - expression-based discovery 720–726
 - computational search/prediction 726–728, 787–794
 - computational searches for ncRNA targets 736–737
 - Hfq coimmunoprecipitation 724–726
 - high-throughput sequencing (RNA-Seq) 721–724
 - microarray 720–721, 794–797
 - mRNA target validation 737–742
 - laboratory protocols 744–776, 797–799
 - strategies to decipher the biological role 728–737
 - miRNA profiling, barcoded cDNA libraries for NGS 861–872
 - Pb²⁺-induced cleavage 275
 - purely computational approaches. *See also* homology search
 - computational methods for expression profiling based on RNA-seq data 821–823, 827
 - homology search for small structured ncRNAs 595–616, 619–621
 - mascRNA 621–629
 - RsaE RNA 600
 - shape analysis, ncRNA consensus structure prediction 583
 - screening for RNA genes and motifs 595–596
 - identification *via* APART 1197
- non-coding RNP (ncRNP) 801–802
 - ncRNP transcriptome 814
- non-radioactive labeling of RNA 89, 92, 100–101, 1103–1104
 - digoxigenin, internally 92, 100–101
 - biotin 8, 66, 117–125, 937
 - fluorescent dye 117–125, 152, 158, 238, 796–797
- phosphorothioates and iodine cleavage 329–364, 369–393
- Northern blot analysis 89–102, 744–747, 808, 816, 1199. *See also* small RNAs, Northern blot detection
 - aptamer detection 1199
 - laboratory protocols 98–102, 744–747, 808
 - mRNA targeted by miRNA 835
 - miRNA detection 1091–1092, 1094
 - ncRNA verification 789, 808, 816
 - of RNA fragments crosslinked to protein 1030–1031
 - probe with digoxigenin label 89, 95, 101–102
 - radioactively labeled probe 744–747, 1094
 - sRNA expression profile 729, 816
- nuclear extract depletion, with biotinylated RNA 951–952
- nucleation zone, RNA crystallization 485–486, 489
- nucleic acid libraries 1148–1150
 - design 1148–1149
 - pool preparation 1149–1150
- nucleic acid library preparation 1108–1109
- nucleoside analog. *See also* modified RNAs
 - photoreactive 900
 - 4-thiouridine 7, 151–166, 231–250, 878–897, 999
 - 6-thioguanosine 7–8, 231–250, 333–334, 878–882
 - thiotriphosphate 329, 333–334
- nucleotide analog interference mapping (NAIM) 207, 232, 285–286, 329–330, 369–371
 - experimental protocols 333–355, 380–386
- denaturing PAGE of analog-modified RNA pools 336, 381–383
 - nucleoside analog thiotriphosphates 333
 - incorporation by transcription 335–337, 371–372, 380–381
 - iodine cleavage 354–355, 385–386
 - radioactive labeling of RNA pool 337–339, 383–384

- method description 330–332, 369–373
 - NAIM results
 - analysis and interpretation 355–358, 373–375
 - data evaluation 386–387
 - influencing factors 372–373
 - selection step/assay
 - analysis of bacterial RNase P function 369–372
 - *cis*-cleaving RNase P RNA-tRNA conjugates 375–380, 384–385
 - group II intron transesterification 339–344
 - group II intron folding, Mg²⁺-induced 344–347
 - Rho-dependent transcription termination 347–351
 - Rho helicase unwinding 351–354
 - nucleotide analog interference suppression (NAIS) 329–332
 - applications 332–333
 - elucidating tertiary contacts in group I and II ribozymes 332–333
 - to the RNase P system 375, 378–379
 - data analysis and presentation 363–364
 - experimental protocols 358–364
 - functional analysis of RNA mutants 362
 - RNA mutants, design and construction 358–362
 - RNA molecules, preparation with single-atom substitutions 359, 361–362
 - selection step 362–363
 - selection assay, *trans*-branching 339–341
 - nucleotide blast. *See* BLASTN
- O**
- OligoCalc (oligonucleotide properties calculator) 60
 - oligonucleotide, DNA
 - biotinylated (bait) oligonucleotide 178–179, 735
 - biotinylated oligonucleotide for affinity purification of RNA-protein complexes
 - complementary biotinylated DNA oligonucleotide 948
 - chimeric 2'-*O*-methyl RNA/DNA oligonucleotide 940
 - displacement oligonucleotides 937, 940, 945–948
 - circularization of linear RNA 75
 - complementary DNA oligonucleotide for Tb(III)-footprinting 259–260
 - DNA splint for RNA ligation 51, 57–70, 77–79, 293, 362, 501
 - DNAzyme 35–37
 - microarray for detection of small ncRNAs 720
 - microarray for HT-RNomics 795, 802, 804–808, 809–812
 - microarrays for gene expression profiling using RNA-seq 821, 827–828
 - Northern blot probe 745–747, 944
 - for miRNA detection 1091
 - primer extension 207, 220, 278–279, 1033, 1045–1046
 - randomized, in SELEX 16, 1103, 1108–1109, 1148, 1172–1173, 1186, 1208–1209, 1215
 - RNA ligase substrate specificity 72
 - RNase H cleavage 54–56, 967, 1166
 - RNase H mapping of crosslinked RNA 248
 - splint labeling 183
 - template for T7 transcription 4–5, 10, 12
 - unwinding of RNA-DNA hybrids by Rho helicase 351–354
 - oligonucleotide LNA, LNA/DNA, LNA/RNA 153, 729, 1088, 1098, 1247–1254, 1258–1265
 - LNA-ATP/UTP, for T7 transcripts 7
 - LNA/DNA mixmers as probes for Northern blotting 92–94, 96–97, 100–101
 - 5'-digoxigenin (DIG)-labeled 89, 91–97
 - oligonucleotide, RNA
 - adaptor RNA 724
 - analysis of synthetic RNA 143
 - atomic force microscopy (AFM) 528
 - biotinylated antisense oligonucleotides for affinity purification of RNP complexes 957, 976
 - 2'-*O*-methyl RNA oligonucleotide 936–941, 945–947, 949, 952–953
 - chimeric 2'-*O*-methyl RNA/DNA oligonucleotide 940
 - calibrator oligoribonucleotide 863–864, 866–867
 - cDNA libraries for miRNA profiling 863–864, 871
 - chemical RNA synthesis 130, 137, 139–147
 - crystallization of RNA 482, 493, 495
 - enhancing siRNA performance 1248, 1261
 - homooligonucleotide stretch (NAIM) 372
 - iCLIP 900–904, 915
 - initiator oligonucleotides for T7 transcription 8

- oligonucleotide, RNA (*contd.*)
 - labeling of alkyne-containing RNA for smFRET 509–511
 - oligo-U stretch 789
 - PAR-CLIP 878, 890, 895
 - purification of phosphorothioate stereoisomers 290–293
 - RACE 747–750, 753
 - Rho helicase interaction (NAIM) 351–354
 - RNA ligation using T4 DNA ligase 45–70
 - RNA ligation using T4 RNA ligase 75–76
 - – RNA donor oligonucleotide 81
 - RNA molecules containing single atom substitutions 359
 - RNA sequencing 192–194
 - – oligonucleotides derived from alkaline hydrolysis or nuclease P1 digestion 191–192
 - separation of oligomeric and monomeric RNA 108, 115
 - smFRET experiments 509
 - synthesis of modified oligonucleotides 151–166
 - temperature-gradient gel electrophoresis (TGGE) 427, 430, 434–435, 438
 - vector-encoded siRNA and miRNA 1228–1234
- oligoribonucleotide. *See* oligonucleotide, RNA
- Open BABEL utility, MD simulation 702
- OPLS force field, MD simulation 695–696
- optical density 446–447, 765
- osmolytes 463
- outer-sphere coordination, cation: RNA 319
- overlapSelect algorithm, next-generation sequencing 827
- p**
 - Packmol package, MD simulation 694
 - PAGE (polyacrylamide gel electrophoresis). *See* gel electrophoresis
 - paired-end reads, next-generation sequencing 825–826
 - PAR-CLIP
 - (photoactivatable-ribonucleoside-enhanced crosslinking and immunoprecipitation) 877–897
 - cDNA library construction outline 882
 - cDNA library preparation, from recovered RNA fragments 889–893
 - crosslinking and immunoprecipitation procedure 882–888
 - – UV crosslinking of 4SU-labeled cells 883
 - determination of 4SU incorporation levels 893–897
 - materials 878–881
 - protein-binding sites on RNA, transcriptome-wide identification 877–878
 - particle mesh Ewald (PME) summation method 697
 - partition function 550, 555, 566–570
 - pattern-recognition receptors (PRRs) 1256
 - Pb²⁺-induced cleavage. *See* probing RNA structure
 - P_{BAD} promoter. *See* promoter
 - PCR. *See* polymerase chain reaction
 - PDB (Protein Data Bank) 688–693
 - imprecision and errors in PDB structure files 690–692
 - refinement error 691
 - NMR structures 690
 - nucleic acid structures 689
 - PDB_REDO database/web site 688
 - – automatic refinement of PDB structures 691
 - PDB2SAX program (SAXS) 419
 - PDB2VOL program (SAXS) 417–419
 - periodate oxidation
 - 3'-terminal ribose of RNA 49, 73, 117–119, 124, 863
 - nucleosides and nucleotide-5'-phosphates 118
 - nucleotides 124
 - peripheral blood mononuclear cells (PBMCs) 1256–1257
 - Pfu* (DNA) polymerase
 - 3'-5' proofreading activity 37
 - phenol, phenol/chloroform extraction. *See* RNA extraction
 - phosphoramidites 45, 129, 131–134, 152–154, 156, 359, 509
 - phosphorothioate 156, 231, 285–286, 333, 370
 - analogs and transcripts preparation 335–337
 - incorporation techniques, into RNA 291–293
 - metal ion rescue of phosphorothioate modifications 285
 - stereoisomers purification, by RP-HPLC 290–291
 - photocrosslinking/UV crosslinking 159, 231–249. *See also* crosslinking and PAR-CLIP
 - probing RNA solution structure 231–232
 - – techniques for modified nucleotide incorporation 232–233

- transcription priming with GMPS 233–234
- aryl azide photocrosslinking agent, attachment to 5'-terminal phosphorothioate 236–238
- 3'-addition of aryl azide photocrosslinking agent 238–240
- 4-thiouridine (4SU) and 6-thioguanosine (6SG) 240–243
- photocrosslinking of RNase P RNA and 6SG-modified precursor tRNA 244–247
- primer extension mapping of crosslinked nucleotides 247–249
- plasmid DNA sonication 1172
- PLEXY (prediction of box C/D snoRNAs) 811
- Pme* I restriction enzyme/site 864–865
 - digestion 871–872
- poly(A) polymerase 398, 401
 - 3'-³²P-end-labeling of RNA using [α -³²P]-Cordycepin-triphosphate 182–183, 337–339
 - C-tailing of ncRNAs for cDNA library construction in experimental RNomics and RNPomics 803–806, 812, 814–815
- poly(A)-specific ribonuclease (PARN) 397–403
- polyacrylamide gel electrophoresis (PAGE). *See* gel electrophoresis
- polymerase chain reaction (PCR). *See also* RACE
 - melting point lowering agents
 - betaine 1150
 - dimethyl sulfoxide (DMSO) 1150
 - MasterAmp PCR Enhancer 1150
 - Qiagen Q-Solution 1150
 - qRT-PCR 89, 861, 1092–1093
 - RT-PCR 809–811, 1107, 1118, 1181–1182
 - inhibition by DTT 1119
 - double-nested 1210, 1213–1214, 1217–1218
 - kit, one-step 305, 310
- polystyrene
 - chemical RNA synthesis 131, 133, 135–136, 145
 - paramagnetic beads 938
 - tubes in flow cytometry 764
- postsynthetic RNA labeling 151–152, 155–157, 164–166
 - fluorophore 509
 - introduction of EPR nitroxide spin label 152, 157, 159–160, 164–166
 - at 2'-amino groups 156–164
 - reaction with succinimidyl esters 158
 - reaction with aromatic isothiocyanates 158–161
 - reaction with aliphatic isocyanates 159–161
 - at 4-thiouridine 164
- potato spindle tuber viroid (PSTVd) 431–440, 532–537
- prey
 - (target) RNA 179
 - protein prey plasmid 1076, 1078
- primer extension 56, 206–207, 210, 212, 217, 220–224, 226, 231, 241–243, 245, 247–249, 259, 265, 276, 278–279, 304, 308–311, 635, 644, 747, 773, 900, 944, 953, 977, 991, 993, 995–996, 998–1000, 1002, 1009, 1030, 1033–1038, 1040–1042, 1045–1047, 1050, 1108, 1166, 1213
- primer walking 740, 1200–1201
- probabilistic shape analysis 585–587
- probability dot plot 567, 570
- probing divalent metal ion binding sites of proteins by hydroxyl radicals 397–404
- probing RNA structure. *See also* RNase (ribonuclease)
 - *in vitro* with enzymes and chemicals 205, 224, 422, 435, 462, 563–564, 572, 602–603, 634–635, 740–741
 - backbone-specific chemical probes 211
 - base-specific chemical probes 210
 - CMCT 208–210, 212, 217–218, 635, 740, 991–995, 1006
 - DEPC 188–189, 208, 210–212, 218–220, 225–226
 - dimethylsulfate (DMS) 188–189, 206, 208, 210–212, 217–218, 220, 224, 226, 310, 359, 635, 642, 644, 740, 992–999, 1002, 1007
 - ENU 209–210
 - enzymes 207, 992–993, 995–1000, 1130
 - in-line probing 206, 210–211, 213, 216, 218, 225–226, 417, 1106
 - kethoxal 212, 740, 992, 994–996
 - lead(II) = Pb²⁺ 206, 225, 277–278, 740, 1130
 - Pb²⁺-induced cleavage of RNA 269–271, 1006
 - Pb²⁺-induced cleavage to probe metal ion binding sites, RNA structure and ligand interactions 271–275
 - primer extension 220, 1037
 - probing RNA solution structure by photocrosslinking 231–232, 244–249

- probing RNA structure. *See also* RNase (ribonuclease) (*contd.*)
 - – probing RNA structure and ligand binding sites on RNA by Fenton cleavage 301–314
 - – ribose acetylation 219
 - – RNA helicase, NAIM 354
 - – RNA preparation 213
 - – RNase hydrolysis 214
 - – SHAPE, 2'-hydroxyl acylation 206, 635, 644
 - – structure-specific chemical and enzymatic probes 208–210
 - – Terbium(III)-footprinting 255–267
 - – tertiary contacts, NAIS 332
 - – 2'-hydroxyls involved in tertiary contacts, NAIM 370
 - *in vivo* 226–227, 314
 - – dimethylsulfate (DMS) 206, 208, 222–223, 225, 314, 978, 999, 1002
 - – kethoxal 206, 314
 - – Pb²⁺-induced cleavage of RNA 275–279
 - RNP probing 981, 991–992, 996–1000, 1005–1007
 - promoter 726–731, 761, 1072, 1074, 1081, 1147, 1212, 1228
 - *ADH1* 1073
 - *HIS3* 1069
 - *istR* sRNA 729
 - *lacZ* 1068–1069
 - λ P_L 733
 - *lexA* operator 1068–1069
 - miRNA vector 1228
 - *ompC* 731
 - P_{BAD} (*araC*) 731, 733, 735, 738–739, 755
 - P_{Lac} 733
 - P_{LacO}–1 731–732, 755–758
 - P_{LtetO}–1 755–757
 - P_{tac} 733
 - *RPR1* 1071
 - shRNA vector 1223–1225, 1232
 - SP6 4, 1167–1169, 1173
 - *sraC* 748
 - T7 promoter(s) 4–9. *See also* T7 RNA polymerase
 - T3 promoter 3–4
 - proteinase K digestion 888
 - PURESYS[®] 742
 - *in vitro* translation 766, 768–772
 - PURExpress[®]. *See* PURESYS[®]
 - PyMOL program 418, 657–659, 688, 693, 702
 - pyridyl disulfide-activated matrix/support 1144, 1146
- q**
- QPALMA sequence read alignment program 823, 826
 - QRNA software (comparative genome sequence analysis to detect conserved RNA secondary structures) 726, 788–789
 - qRT-PCR. *See* polymerase chain reaction.
 - quantification plots 311
 - QuB software package, smFRET analysis 507
- r**
- RACE (rapid amplification of cDNA ends) 75, 184, 731, 738, 759, 761
 - 5'-RACE protocol 747–755
 - radioactive labeling (radiolabeling) methods
 - 5'-³²P-end-labeling of RNA 64, 180–181, 198–199, 213, 337–338, 350, 383, 866–867, 883, 885–886, 909–910, 980
 - 5'-³²P-labeling of nucleoside 3'-phosphates 195–196
 - 3'-³²P-end-labeling of RNA 73, 181–183, 213, 337–339, 383–384
 - – with Klenow polymerase, [α -³²P]dNTP and DNA splint 337–338, 341
 - – with poly(A) polymerase and [α -³²P]-Cordycepin-triphosphate 182–183, 337–339
 - – with T7 DNA polymerase, [α -³²P]dATP and DNA splint 182–183
 - DNA primer/probe 5'-³²P-end-labeling 774, 1045–1046, 1091
 - iCLIP RNA 5'-labeling 909–910
 - internal ³²P labeling of RNA by *in vitro* transcription 14, 16, 980–981, 1003, 1112, 1153, 1179, 1190
 - NAIM 369
 - 5'-³²P-end-labeling of analog-modified pool RNA 383
 - 3'-³²P-end-labeling of analog-modified pool RNA 383–384
 - NAIM/NAIS 329
 - ³²P-labeling of an RNA pool 337–339
 - ³²P-labeling of recombinant polypeptide 403
 - probing RNA structure *in vitro* with enzymes and chemicals 213
 - radiolabeling of RNA segments crosslinked to immunoprecipitated proteins 885–886
 - radiolabeling of RNA size markers 866–867, 883

- radius of gyration 407, 416–417
- RALEE, RNA alignment editor 572, 621, 625
- randomized exon cassette
 - design 1210–1211
 - ligation 1216
 - preparation 1215
- Rapidshapes heuristics 586
- parameters 592–593
- reads. *See* RNA-Seq
- reads per kilobase per million (RPKM) units 828
- recombinant polypeptide, radioactive labeling 403
- renaturation 224, 312, 429, 431, 436, 980–981, 1005
- reporter gene 738–739, 769, 1067–1069, 1075–1076, 1078, 1080–1081
- respiratory syncytial virus 1212, 1261
- reversed-phase HPLC (RP-HPLC) 132
 - analysis of purified RNA 143–147
 - analysis of ribonucleosides 893–896
 - click reaction 512
 - ion-pair reversed-phase liquid chromatography (IP-RPLC) 142–143, 213
 - postsynthetic labeling of 2'-amino-modified RNA 162–165
 - purification of phosphorothioate stereoisomers 290–291
 - purification of trityl-on RNA 138, 140–142
- reverse transcription 211, 280, 313, 753, 772–773, 804–807, 809–811, 813–816, 869–870, 891–892, 900–904, 913–914, 1017, 1024, 1099, 1118, 1141, 1151–1152, 1189, 1192–1193, 1208–1210
 - SuperScript™ plasmid system 802–804, 806–807
 - SuperScript II reverse transcriptase 279, 753, 773, 775, 806, 811, 813–815, 1118, 1120, 1151
 - SuperScript III reverse transcriptase 224, 260–261, 305, 310, 750, 753, 865, 869, 879, 881, 892, 913, 1120
- Rfam database 583, 619, 629, 667, 789
- Rho
 - Rho helicase 351–354
 - Rho-independent transcription termination 347–351, 595, 604, 726–727, 789–791
- rhodamine 159
 - tetramethylrhodamine 504
 - sulforhodamine 1139
- ribonuclease. *See* RNase
- ribonucleoprotein (RNP). *See also* RNA binding proteins and RNA-protein interaction
 - analysis of RNA-protein interaction using iCLIP 899
 - affinity purification 957
 - – immunoaffinity purification 958–962
 - – RNA aptamer-based affinity purification 963–971
 - biotin-based affinity purification of RNA-protein complexes 935
 - experimental identification of microRNA targets 1087
 - identification and characterization through three-hybrid analysis 1067
 - immunopurification of endogenous RNAs associated with RNA binding proteins (RBPs) *in vivo* 1017
 - isolation of RNA binding proteins using RNA affinity matrices 921
 - PAR-CLIP 877
 - protein-RNA crosslinking in native ribonucleoprotein particles 1029
 - – UV crosslinking 1031–1033
 - – identification by primer extension analysis 1033–1037
 - – identification of crosslinked proteins 1037–1040
 - – protocols 1042–1050
 - RNA-protein interactions and RNA structure 975
 - – RNP reconstitution 978–981
 - – EMSA 981–986
 - – purification of RNPs reconstituted in complex cellular extracts 986–987
 - – tobramycin-sepharose or MS2-MBP affinity chromatography 987–991
 - – RNA structure probing 991–999
 - – UV crosslinking and immunoselection 999–1005
 - RNPomics 801–802, 812–817
 - sedimentation analysis of RNP complexes 1055
 - – glycerol gradient centrifugation 1056–1060
 - – fractionation of RNPs by cesium chloride density gradient centrifugation 1061–1064
 - specialized SELEX method to identify antisense and protein target sites in RNA or hnRNPs 1165–1166
 - generation of RNA 20-mer library using *Mme* I 1166–1169

- ribonucleoprotein (RNP). *See also* RNA binding proteins and RNA-protein interaction (*cont'd.*)
 - – natural RNA substrates for proteins, identification 1171
 - – procedure and protocols 1171–1182
- riboregulation, riboregulator 96, 724, 743
- ribosomal RNA (rRNA) 93, 95, 118, 175, 206, 220, 232, 302, 304, 308, 310, 481, 493, 501, 549, 677–678, 719–722, 724, 787, 791, 802, 807, 816, 823, 921, 975, 999, 1002, 1029, 1258
 - competitor RNA 924
 - rRNA extraction protocol 309
 - 5S rRNA 95, 269, 551, 571, 746, 1060
 - – temperature-gradient gel electrophoresis 432–435
 - – endogenous standard 89, 94–95, 732
 - – abstract shapes 583
 - – structure prediction 642–644
 - – glycerol gradient centrifugation 1059–1060
 - 16S rRNA 90, 95, 269, 278, 302, 304, 746, 791–794, 1057
 - – Fe²⁺-mediated hydroxyl radical cleavage 310, 311
 - – crystallization 494
 - – high-throughput sequencing 722
 - 18S rRNA 90, 95
 - 23S rRNA 90, 95, 278, 302, 746, 1057
 - – high-throughput sequencing 722
 - 28S rRNA 90, 95, 637, 640
 - – structure prediction 646, 657, 659
- ribozyme 6, 53–54, 75, 78–79, 106–107, 124, 129, 152, 158, 176, 223–224, 241, 244, 248, 416, 442, 492, 500–501, 527, 535, 549, 595, 1098, 1140, 1166
 - folding 245–246
- NAIM/NAIS: application to the RNase P system 372, 375, 378, 380–381, 384,
- NAIM/NAIS: group I and group II ribozymes 332, 339, 344
- primer extension analysis of crosslinks between photoagent-modified precursor tRNA and RNase P 248–249
- probing RNA structure and ligand binding sites by Fenton cleavage 301
- protocol for RNase P cleavage 40–41
- ribozyme cassettes 30
 - – *cis*-cleaving 5'-cassette 30
 - – *cis*-cleaving 3'-cassette 30–31
 - – *trans*-cleaving for generation of homogeneous 3'-ends 33–35
 - – construction 36–39
 - – terbium (III) footprinting of HDV ribozyme 261–264
 - – thiophilic metal ion rescue of RNA phosphorothioate modifications 286
- RISC. *See* RNA-induced silencing complex
- rmprune utility program 607
- RNA Affinity in Tandem (RAT) method 935
- RNA backbone correction (RNABC) program 688, 691
- RNA binding proteins (RBPs) 244, 373, 440, 527, 579, 842, 852, 976, 1041, 1166, 1171, 1188, 1207, 1222. *See also* ribonucleoprotein and RNA-protein interaction
- analysis of RNA-protein interaction using iCLIP 899
- biotin-based affinity purification of RNA-protein complexes 935
- RNPomics 801–802, 812–817
- identification and characterization of small ncRNAs in bacteria 722
 - – RNASeq 721
 - – Hfq coimmunoprecipitation 724–726
 - – genomic SELEX 728
 - – finding sRNA-associated proteins 735–736
- identification and characterization through three-hybrid analysis 1067
 - – basic strategies 1068–1070
 - – components 1070–1075
 - – protocols 1075–1079
 - – additional applications 1081–1082
- immunopurification of endogenous RNAs associated with RBPs *in vivo* 1017
- – RNA coimmunoprecipitation 1020–1025
- isolation of RNA binding proteins using RNA affinity matrices 921
 - – method 921–927
 - – applications 927–931
- PAR-CLIP 877
- RNABOB, fast RNA motif/pattern search software 93, 621, 626, 811
- RNA extraction 222, 225, 280, 724, 744, 901, 911, 1037, 1044–1045
- phenol, chloroform, phenol/chloroform 5, 15, 18–19, 20, 31, 35, 61–66, 69, 80, 90, 94, 98–99, 105, 107, 112, 121, 125, 183, 213–214, 219, 223, 238, 247, 278, 305, 309, 338, 350, 381, 482, 537, 744–745, 750, 752–753, 768–770, 774–775, 804–806, 809–810, 812–813, 815–816, 866, 880–881, 888, 912, 939, 944, 962, 971, 995, 998–999, 1018, 1021, 1026, 1044, 1057, 1059–1062, 1064–1065,

- 1090–1091, 1094, 1112–1114, 1116, 1130, 1157–1158, 1181, 1192, 1216–1217, 1231
- (diethyl)ether 69, 1116, 1130
- elution from gels 62–63, 806, 1180–1181
- hot phenol method 90, 94
- *in vivo* SELEX 1216–1217
- kits 90
- recovery/extraction from membranes 901, 910–913, 1192
- rRNA extraction 309
- TRIzol reagent/method 90, 744, 750–751, 803, 806, 808, 812, 866, 894
- RNAfold 42, 84, 92, 438, 579, 620, 673, 675, 681, 740, 811, 1073, 1106
- secondary structure prediction 549
- – formal background 552
- – internet addresses 557
- – RNAfold 565
- – output from the RNAfold Server 570
- RNAforester, structure alignment program 572, 592, 624–625
- RNA fragments, simultaneous splint ligation, for FRET experiments 66–67
- construct design 68–69
- RNAhybrid algorithm, RNA:RNA interaction prediction 737, 759, 767
- RNA interference (RNAi) 845, 1166, 1210.
- See also small interfering RNA
- gene silencing using vector-encoded siRNAs or miRNAs 1221
- – background information 1221–1223
- – construction of shRNA vectors 1223–1228
- – construction of miRNA vectors 1228–1229
- – construction of extended shRNAs (e-shRNAs) and lhRNAs 1229–1230
- – production of lentiviral vectors encoding anti-HIV-1 shRNAs or e-shRNAs 1230–1234
- RNA-induced silencing complex (RISC) 833, 841–842, 843, 1087, 1221–1223, 1232
- – immunoprecipitation of RISC components 838–839
- using chemical modification to enhance siRNA performance 1243
- RNA isolation. See also RNA extraction, RNA preparation and RNA purification
- 5'-RACE 751–752
- crosslinked RNA 246–247
- iCLIP 911–913
- miRNA profiling by next-generation sequencing 866
- Northern blot detection of small RNAs 89
- PAR-CLIP 883–888, 894
- Pb²⁺-induced cleavage *in vivo* 277–279
- pure RNA species from biological material 176–180
- – single RNA species with partially known sequence 178–179
- – single unknown RNA species following a functional assay 176–178
- – size-fractionated RNA 176
- RNAshapes program 580, 583, 587
- parameters 588, 593
- RNAMOTIF, RNA secondary structure definition and search algorithm 596–599, 606–609, 615, 811
- program versions and download site 616
- RNA preparation. See also RNA extraction, RNA isolation, RNA purification and T7 RNA polymerase
- Atomic Force Microscopy (AFM) 537–538
- *in vitro* transcription and purification of milligram quantities of RNA 12–13, 111–113
- – weak anion-exchange chromatographic purification 113–115
- Northern blot detection of small RNAs 89
- preparation from gradient fractions 1059–1060, 1064
- preparation for RT-PCR 1114
- RNA ligation 80–81
- RNA molecules containing single-atom substitutions 359–362
- RNA-protein interaction and RNA structure 980–981
- RNA sample enriched through immunoprecipitation 809
- RNA structure probing 213, 223
- RNomics and RNPomics 801
- – cDNA library construction 804–808
- – specialized cDNA library construction 808–811
- – RNP-derived cDNA library construction 814–816
- simultaneous preparation of RNA and protein 1060
- transcripts containing phosphorothioate analogs 335–336, 378–384
- RNA-protein crosslinking 975–978, 999–1005, 1008, 1009, 1191, 1199
- aptamer target selection 1100–1101
- iCLIP 900–902, 905–906
- *in vivo* 1021–1022
- native RNP particles 1031–1033, 1042–1044
- PAR-CLIP 877–878, 882–884

- RNA-protein interaction. *See also*
 - ribonucleoprotein and RNA binding proteins
 - affinity purification 935, 945, 949, 953
 - aptamer-tagged sRNA 736
 - atomic force microscopy (AFM) imaging and force spectroscopy 527, 536, 541
 - coimmunoprecipitation 724
 - deep sequencing 742
 - fluorescence techniques 8, 117, 501
 - genomic SELEX 1199–1200
 - NAIM/NAIS 333, 347–350, 351–354
 - RNA ligation methods 50
 - study by crosslinking 232
 - study in RNPs 975
 - temperature-gradient gel electrophoresis (TGGE) 439–441
 - three-hybrid analysis 1067–1068, 1070–1071, 1074–1076, 1080–1083
- RNA purification. *See also* RNA extraction, RNA isolation and RNA preparation
 - anion-exchange chromatography 111–115
 - HPLC purification 482–483
 - polyacrylamide gel electrophoresis (PAGE) 483–484
 - – denaturing (PAGE) 18, 61–63, 83–84, 381–383, 770
 - – diffusion elution 19, 484
 - – electroelution 19–20
 - RNA downstream purifications 122
 - – gel chromatography 122
 - – purification on denaturing polyacrylamide (PAA) gels 122–123
 - chemical RNA synthesis 138–139
 - – anion-exchange HPLC purification 139–140
 - – reversed-phase HPLC purification of trityl-on RNA 140–142
 - – detritylation of trityl-on RNA 142
 - – desalting by HPLC 142–143
 - isolation of pure RNA species from biological material 176
 - – preparation of size-fractionated RNA 176
 - – isolation of a single unknown RNA species following a functional assay 176
 - – comments on electrophoretic purification 178
 - – purification of a single RNA species from 1 mg crude small RNAs 179–180
 - purification of phosphorothioate stereoisomers by RP-HPLC 290–291
 - immunopurification of endogenous RNAs associated with RBPs *in vivo* 1017
 - – RNA coimmunoprecipitation 1020–1022
- RNA recognition motif (RRM) 958, 975–976
- RNase (ribonuclease) 210, 224–225, 325, 921, 945, 984, 1008–1009, 1025, 1280
 - activity of T7 RNAP 9
 - cleavage susceptibility to derive base-pairing constraints (MC-Tools) 635
 - complete hydrolysis of RNA for UV spectroscopy 1281
 - – 3'-exonuclease ERI-1 1253
 - from plants 434
 - nuclease P1 190–192, 195–196, 198–199, 1281
 - – identification of terminal nucleotides 184, 190
 - poly(A)-specific ribonuclease (PARN) 397–402
 - ribonuclease recognition sites 738
 - RNase I
 - – iCLIP 900, 902, 906–909, 911–912
 - RNase III 975, 1087
 - – Dicer 833, 852, 1087, 1222–1223, 1227–1228, 1232, 1235, 1243–1244, 1249, 1263
 - – Drosha 833, 1087, 1222–1223, 1228, 1232, 1235
 - probing RNA structure 740
 - RNase A 1000, 1003, 1253, 1281
 - – analysis of nucleotide content 195–196, 198
 - – probing RNA structure 635, 740
 - – RNA 3'-ends 124
 - – tissue activity 906
 - RNase CL3
 - – enzymatic sequencing 185–187
 - RNase contamination 20, 59, 70, 84, 146, 175, 226, 280, 324, 327, 389, 495, 520, 536, 897, 921, 979, 987, 1003, 1025–1026, 1041, 1093, 1107, 1113
 - RNase D 397
 - RNase E 742–743
 - RNase H 750
 - – cDNA library construction 804, 807
 - – engogenous activity 59
 - – generation of homogeneous ends 35, 54
 - – generation of RNA ligation substrates 51–52, 57
 - site-directed cleavage 51–56, 64, 967, 970
 - – site-directed cleavage protocol 63
 - mapping 248, 937, 939, 1030, 1166
 - reverse transcription 753, 1151

- RNase inhibitor 10, 13–14, 17–18, 20, 60–61, 63–65, 70, 81–82, 234, 293, 305, 310, 336, 362, 536, 750, 752–753, 812, 815, 906, 938, 952–953, 1019, 1020–1021, 1026, 1113, 1153, 1174, 1179, 1181–1182, 1190
- RNase P (RNA) 80, 922, 924, 927, 1055, 1280
- bioinformatic studies 595, 619, 622, 628, 678
- cleavage protocol 40–41, 293–294
- crystallization 481, 492
- deletion of gene for RNA subunit 730
- generation of homogeneous 3'-ends 33–35
- *in vitro* T7 transcription 8, 13–14
- mascRNA processing 622
- menRNA processing 628
- metal ion coordination interactions 287–290, 292
- Mg²⁺-probing 277
- NAIM 369–393
- Pb²⁺-probing 271–273, 275–276
- RNA ligation 77
- SAXS 411, 417, 422
- single molecule FRET 500
- stoichiometry of Mg²⁺ ions bound to RNA 326
- structure probing by photoaffinity crosslinking 231–232, 241–242, 244–246, 248–249
- terbium (III) footprinting 264
- three-hybrid analysis 1068, 1071, 1075, 1082
- RNase protection assay 1199, 1213
- RNase S1
 - probing RNA structure 635, 1130
- RNase T1 1003–1004, 1281
- RNA 3'-ends 124
- enzymatic sequencing 185–187, 192, 1108–1109
- probing RNA structure 207–208, 212, 214–217, 224, 226, 635, 740, 992–993, 995, 998–1001, 1130
- terbium (III) footprinting 258, 263
- sequencing ladders 306–307
- PAR-CLIP cDNA library construction 879, 882, 884–887
- RNase T2 1281
- identification of terminal nucleotides 184, 190–191
- analysis of nucleotide content 195–196, 198
- probing RNA structure 207–208, 212, 214–215, 635, 740, 992–993, 995–998
- RNase U2
 - enzymatic sequencing 185–187
- probing RNA structure 635
- RNase V1
 - probing RNA structure 207–208, 210, 212, 214–215, 217, 224, 635, 740, 992–993, 995–998, 1130
- RNase Z (tRNase Z)
 - menRNA processing 628
- purification using RNA affinity matrices 922, 924–925, 927–929
- siRNA stability 1248, 1253–1255, 1261
- *Staphylococcus aureus* nuclease 185
- RNase-free water preparation 8, 59, 175, 280, 389, 512–513, 744, 803, 1143, 1173
- RNA-Seq (high-throughput sequencing)
 - aptamer selection 1099, 1106, 1109, 1122
 - barcoded cDNA libraries for miRNAs 861–872
- cDNA reads
 - iCLIP 901
 - microRNAs 861–863
 - PAR-CLIP 878
- computational methods for gene expression profiling 821–826, 829
- expression-based discovery of bacterial sRNAs 720–726
- differential RNA sequencing (dRNA-seq) approach 722–724, 732
- genomic SELEX 1186, 1194–1195, 1197
- iCLIP 899–900, 901, 916
- immunopurification of endogenous RNAs associated with RBPs 1018, 1025–1026
- PAR-CLIP 877, 886–887, 889, 892–893
- protein-RNA crosslinking in native RNP particles 1029
- RNA sequencing
 - chemical 187–189
 - enzymatic 184–187
 - identification of modified nucleotides 194–201
- terminal RNA sequence determination 190–194
- RNashapes program 580–581, 584–586, 588, 590, 624
- parameters 585, 592–593
- RNA tester sequence (RTS) 1069, 1071–1073
- RNAz program, conserved secondary structure identification 620, 624, 788–790
- RNomics 794, 801
- library construction and analysis protocols 804–811

RNomics (*contd.*)

- materials 802–804
- ncRNA sequence, computational analysis 811

RNPomics

- library construction protocols 813–816
- materials 812–813
- RPR1 promoter 1071

S

S2S 572, 667–675, 677–681

- interactive RNA alignment viewer and editor 668–671
- Assemble2 667–668, 671–677, 680–684
- – interactive RNA 3D modeler 671–672
- – installation of S2S and Assemble2 673–674

SAM (S-adenosylmethionine) 1174, 1178

- SAM riboswitch 501

SAM format (computational tool for RNA-Seq) 823, 826, 1197

SAMs (self-assembled monolayers) of alkanethiols on gold surfaces 543

SAXS. *See* small-angle X-ray scattering

S-domain (RNase P RNA) 272, 388–389, 408, 411–416, 418–419, 421

- folding intermediate
- – global conformation 412–414
- – stable and extended conformation 414–416

SDS-PAGE. *See* SDS-polyacrylamide gel electrophoresis

SDS-polyacrylamide gel electrophoresis

- (SDS-PAGE) 23, 114, 399–404, 729–732, 881–882, 886–887, 897, 901, 908, 910–911, 928–931, 942–943, 949–950, 961, 968–969, 971, 977, 980, 984–986, 990–992, 1003–1005, 1042, 1057, 1060, 1090, 1100–1101, 1110–1111

- NuPAGE 880, 910, 968–969, 971

secondary structure prediction, of RNA 549

- mfold (UNAFold web server) 92, 549, 555–565

- – energy dot plot 560–561

- – p-num 561–564

- – ss-count 561–564

- thermodynamics 550–552

- RNAfold (Vienna RNA server) 92, 549, 565–571

- – mountain plot 571

- – probability dot plot 569–570

sedimentation analysis, of ribonucleoprotein complexes

- cesium chloride density gradient 1055–1056, 1061–1064

- glycerol gradient centrifugation 802, 813–816, 940, 948, 960–962, 966–968, 970–971, 976, 986, 1056–1059, 1061

- sucrose gradient 839, 1061

SELEX (Systematic Evolution of Ligands by EXponential enrichment)

- database 1098

- genomic 720, 728, 1165–1166, 1171, 1185–1186

- identification of antisense and protein target sites in RNA or hnRNP complexes 1165–1182

- *in vitro* selection 16, 442, 1097, 1099–1101, 1104–1107, 1116

- – against proteins and carbohydrates 1097–1131

- – against small targets 1139–1162

- – selection cycle 1099, 1104–1106, 1141

- *in vivo* strategies 1207–1218

- – minigene design 1212–1213

- – monitoring for enrichment of exon sequences that function as splicing enhancers 1213–1214

- – procedure overview 1208–1210

- – protocols 1215–1218

- – randomized exon cassette design 1210–1211

- – RT-PCR amplification 1213

- subtractive 1098

Semi-Automated Footprinting Analysis (SAFA) software 221

sense strand (SS)

- DNA cleavage by *Bse* RI and *Bsg* I 761

- genome

- – coverage in microarrays 720

- – ncRNA-coding 789

- lhRNA 1226–1227, 1230

- shRNA 1225–1227

- siRNA 1243–1245, 1249–1251, 1253, 1255, 1257, 1263–1265

Sephadex, Sepharose. *See* chromatography sequence reads. *See* RNA-seqsequencing. *See also* next-generation

sequencing (NGS), RNA-seq and RNA sequencing

- direct RNA sequencing 175, 183–184, 1029

- – enzymatically 184–187, 725

- – chemically 187–189

- – determination of terminal RNA sequences by two-dimensional mobility shift 190–194

- identification of modified nucleotides 194–201
- phosphorothioate-modified RNA by iodine 341, 350
- DNA sequencing reaction/ladder, dideoxy 217, 221, 223, 226, 245, 248, 259–261, 311, 741, 774, 807, 995–997, 1000–1002, 1034, 1046–1047, 1108
- additive betaine 1234
- DNA sequencing kit 772, 804, 1224, 1227
- of PCR products 754–755
- of 5'-RACE products 748–749
- Sanger method 720–721, 802, 816, 833
- SEQUEST, tandem mass spectrometry data analysis program 929–930
- ShapeFinder software, analysis of nucleotide reactivity 221, 265
- short hairpin RNA (shRNA) 1222, 1243–1244. *See also* extended shRNA (e-shRNA)
- construction of vectors 1223–1228
- Dicer-dependent short shRNA 1244
- lentiviral vectors encoding anti-HIV-1 shRNAs 1230–1235
- long shRNA 1244
- shreps (shape representative structures) 581, 586–591
- computing 584–585
- shRNA. *See* short hairpin RNA
- shuffle program 599–600, 608, 616
- Simulaid package, MD simulation 694, 704
- single-end reads, next-generation sequencing 825
- single molecule fluorescence resonance energy transfer (smFRET) 499–500
- antifading agents 510–511, 515–516
- “click reaction” between RNA-alkyne and dye azide 509–510
- confocal microscopy 500
- “burst mode” vs. measurements with immobilized RNA molecules 500
- donor-only populations 521
- dye labeling of RNA construct 508–509
- dye pairs 504
- choice of pair 507–508
- Förster theory of FRET 502–503
- instrumental setup 505–507
- protocols 511–521
- bulk FRET measurements 513–515
- measurement chambers and sample preparation 516–520
- removal of unbound fluorophore 520–521
- RNA construct design 503–505
- RNA dynamics addressed by smFRET 500–502
- RNA immobilization 519
- biotin-streptavidin-biotin interaction 519
- surface coverage 521
- surface passivation 519
- RNase contamination 520
- sample drying 521
- single-stranded DNA-binding protein (SSB), in T7 transcription 21
- SIPHT web interface, prediction of bacterial intergenic loci 790–791
- siRNA. *See* small interfering RNA
- SITUS suite for SAXS analysis 417–419
- Sma* I restriction enzyme/site 532–533, 537, 1071–1072
- 7SK RNA 628, 827
- 7SK polymerase II promoter 1225
- small-angle X-ray scattering (SAXS) 407–409
- application example bacterial RNase P RNA 411
- folding intermediate of *Bacillus subtilis* RNase P RNA S-domain 412–416
- experimental setup 410–412
- RNA modeling, low-resolution real-space reconstruction 416–421
- small interfering RNA (siRNA) 129–130, 142, 301, 842, 976, 1221–1223, 1227–1228, 1230, 1232, 1235, 1243. *See also* gene silencing methods
- architectures, siRNA designs 1243–1244
- helix stability asymmetry 1245
- aiRNA, asiRNA, bulge-siRNA, blunt fork-siRNA, blunt siRNA, 27 bp-siRNA, Dicer-independent short shRNA, dumbbell siRNA, long shRNA, sisiRNA, ss-siRNA 1244
- chemical modification 1244–1245
- backbone modification 1246–1248
- base modifications 1252
- ribose 2'-OH substitutions 1248–1251
- ribose backbone alteration 1251–1252
- guide for *in vivo* studies 1261–1265
- guide strand = antisense strand (AS) 1243
- passenger strand = sense strand (SS) 1243
- potency improvement 1252–1253
- stability against ribonucleases 1253
- stabilization strategies 1254–1255
- silencing duration enhancement 1255–1256
- immunogenicity
- abrogation by chemical modification 1257–1258

- small interfering RNA (siRNA) (*contd.*)
 - cellular responses 1256–1257
 - off-target effects, reduction 1258–1259
 - pharmacokinetics, improvement by chemical modification 1259
 - biodistribution, altering by conjugation 1261
 - cellular delivery enhancement by conjugation 1260
 - tolerance toward modification 1244–1245
- small non-coding RNA (sRNA), bacterial 89, 301, 600, 719–720
 - biological role deciphering 728–729
 - associated proteins, identification 735–736
 - biocomputational approaches to find targets 736–737
 - deletion of sRNA gene (knockout strain) 729–731
 - expression profile 729
 - libraries (collections) of sRNA deletion strains or overexpression plasmids 734–735
 - overexpression 731–732
 - pulse expression combined with transcriptome analysis 733–734
 - experimental target validation 737–738
 - *in vitro* characterization of function 741–742
 - *in vitro* RNA–RNA footprinting 739–741
 - reporter gene fusions and sRNA chimera 738–739
 - expression-based discovery 720
 - Hfq coimmunoprecipitation 724–726
 - high-throughput sequencing and RNA-seq 721–724
 - microarray 720–721
 - expression-independent searches 726
 - biocomputational prediction, of sRNAs 726–728
 - biocomputational prediction, of sRNA targets 736–737
 - genomic SELEX 728
 - protocols
 - 30S ribosome toeprinting assay 772–776
 - flow cytometry 764–767
 - green-fluorescent protein (GFP)-based reporter system for target validation 755–764
 - Northern blot protocol 744–747
 - PURESYSTEM *in vitro* translation 766, 768–772
 - 5'-RACE protocol 747–755
- small RNAs, Northern blot detection 89
 - application example 6S RNA-derived pRNAs 96
 - DIG-labeled probes for detection 95, 101–102
 - hybridization with complementary probe 92
 - DNA/LNA mixmer probe design 92–94
 - method limitations 96–97
 - native versus denaturing gels 90–91
 - Northern hybridization protocols 98–102
 - probe generation by T7 transcription using DIG-11-UTP 89, 92, 100–101
 - RNA isolation 89–90
 - RNA transfer and fixation to membranes 91
 - 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide (EDC) crosslinking 89, 91–92, 95–97, 100
- smFRET. *See* single molecule fluorescence resonance energy transfer
- SnoReport (computational identification of snoRNAs with unknown targets) 811
- snRNP (small nuclear ribonucleoprotein). *See* spliceosomal small nuclear ribonucleoproteins
- SOAP sequence read alignment program 823, 826
- solid-phase synthesis of RNA 132–136, 151–152
 - automated RNA synthesis 136
 - chemistry variants
 - 2'-ACE method 129
 - 2'-O-TBDMS method 129
 - TOM method 129
 - deprotection protocol 136–137
 - desilylation protocols 137–138
 - detritylation of trityl-on RNA 142
 - HPLC purification 138–142
 - anion-exchange 139–140
 - desalting 142–143
 - reversed-phase 140–142
 - manual RNA synthesis 134–136
 - modified nucleoside phosphoramidites 117, 152–155, 359, 509
 - quality control of modified RNA 164–166
 - product quality and purity analysis 143–147
- SP6 linker 1169, 1173, 1180
 - dimer 1170
 - ligation 1167, 1170, 1172
 - preparation and ligation protocol 1178–1179
- spatial resolution of AFM 528, 530–531

- Sph* I restriction enzyme/site 1071–1072
- spliceosomal and small nuclear ribonucleoprotein complexes, affinity purification 957
- anti-m₃G cap antibodies 986
 - biotin-based 935–937
 - immunoaffinity purification using antipeptide antibodies 958–962
 - RNA aptamer-based affinity purification 986–993
 - isolation of native spliceosomal complexes 963–971
- spliceosomal small nuclear ribonucleoproteins (snRNPs) 957, 986, 996, 1029, 1040–1043, 1055–1056, 1061, 1063–1064
- affinity purification using biotinylated antisense 2'-O-methyl RNA oligonucleotides and streptavidin agarose 940–948
 - enrichment by DEAE chromatography 938
 - immunoaffinity purification using antipeptide antibodies 958–962
 - reconstitution 981
 - U1 type 945–948, 960–961, 963, 986, 1031, 1034, 1036, 1038, 1040, 1042
 - U2 type 957, 960–963, 986, 1043, 1064
 - U4 type 963
 - U4/U6 941–942, 986, 1031, 1063–1064
 - U4/U6.U5 tri-snRNP 963, 986, 996–997, 1031, 1043, 1055
 - U5 type 961, 963, 986, 1041–1043, 1064
 - RNA aptamer-based affinity purification 963–971
- splint labeling, of RNA 3'-end 182–183
- spring constant, of cantilever in AFM 528, 535, 538, 541
- sRNA. *See* small non-coding RNA
- sRNATarget web server (bacterial sRNA target prediction) 737
- ss-count (mfold) 561–564
- staining
- cells with Trypan blue 1021
 - DNA 1121
 - ethidium bromide (EtBr) 807, 1215
 - proteins
 - Coomassie (Brilliant Blue) 114, 176, 943, 946, 950, 959, 961–962, 968–969, 971
 - silver 399–400, 402, 930–931, 942–943, 953, 1043, 1048, 1110
 - RNA 430, 768, 770
 - toluidine blue 15, 112, 114, 177–178
 - ethidium bromide (EtBr) 15, 24, 41, 59, 90, 95, 471, 1060
 - silver 178, 180, 427, 433–434, 437, 439, 942, 944, 946, 962, 967–968, 971, 1059, 1063
 - SYBR Gold 41, 768
- statistical RNA hydrolysate 190–192
- stem-loop RNA structure 595, 992–993
- binding to MS2 coat protein 992–993, 1069, 1071, 1073
 - binding to PP7 coat protein 935
 - 3'-extension recognized by *trans*-acting VS ribozymes 33
 - miRNA 1227–1228
 - qRT-PCR of miRNAs 89
 - Rho-independent transcription termination 347, 350–351, 588, 604
 - stem-loop binding protein (SLBP) 951, 1078
 - tRNA 76, 692
 - U1 snRNA 1034, 1036, 1039
 - U4 snRNA 1031
 - yeast U3 snoRNA 1002
 - yeast U5 snRNA 996–997
- streptavidin 117, 947
- alkaline phosphatase conjugate 118
 - aptamer 735, 935, 1108
 - gel retardation/shift of biotinylated RNA 123, 125
 - (agarose) beads/resin/matrix 178–179, 372, 936–937, 940–941, 949–950, 1147, 1154, 1157–1158
 - in smFRET 519–521
 - magnetic beads 937–939, 1147
 - RNP affinity selection/chromatography 936, 976, 1088
 - sensor/cantilever functionalized with, in AFM
 - neutravidin agarose 947
- structural RNA alignment 598–599, 667, 669
- INFERNAL 614
 - LOCARNA 602
 - S2S 668–669, 673, 678–680
 - visualization 573
- 2D and 3D structure prediction, over the internet 633–634
- equipment and input 634
 - MC-Tools 633–634
 - MC-Cons 636, 651–654
 - MC-Fold 635, 637–651
 - MC-Sym 636, 654, 657–661, 663
- 4SU, 6SG. *See* 4-thiouridine and nucleoside analog
- succinimidyl ester 156, 158. *See also* N-hydroxysuccinimide (NHS)

- SuperScript. *See also* reverse transcription
- SuperScriptTM plasmid system 802–804, 806–807
 - surface and sensor functionalization, AFM 542–543
 - surface passivation, AFM 519
 - surface plasmon resonance (SPR) 1106, 1126, 1128–1130
 - Survival of Motor Neuron (SMN) protein 935
 - synchrotron 409
 - beamline 411
 - generation of hydroxyl radicals 304, 314
 - SAXS experiments 409–411, 421–422
 - X-ray beams 301
 - Systematic Evolution of Ligands by EXponential enrichment (SELEX). *See* SELEX
- t**
- T4 DNA ligase (T4 Dnl) 293, 361–362, 757, 1174, 1224
- advantages 49
 - DNA ligation 110, 759–760, 807, 1121, 1178, 1216
 - protocols 60–61, 65–66, 68–69
 - reaction mechanism 46, 48–49, 54
 - RNA ligation method 50–52, 57–58, 78
 - – simultaneous splint ligation of five RNA fragments 66–69
 - substrate specificity 47
- T4 polynucleotide kinase (PNK) 70, 180, 182, 772, 803, 865, 879, 979, 1094, 1174
- 5'-end-labeling with [γ -³²P]ATP 32–33, 57, 64–65, 68–69, 80–81, 181, 195, 198–199, 213, 337–338, 350, 383, 404, 774, 808, 810, 866–867, 883, 885–886, 902, 909–910, 923, 980, 1045–1046, 1091, 1130, 1176–1177
 - 5'-end-labeling by phosphate exchange 181
 - 5'-³²P-labeling of nucleoside monophosphates 196
 - 3'-dephosphorylation, removal of 2',3' cyclic phosphate 33, 39–40, 121
- T4 polynucleotide ligase 46, 51. *See also* T4 DNA ligase and T4 RNA ligase
- T4 RNA ligase(s) 182, 212, 750, 757, 803, 813, 865, 879, 979
- adapter ligation 752–753, 804–805, 809–811, 814–815, 862–872, 886–887, 889–891, 901, 903–904, 909–910, 1037
 - applications 73–76, 79–80
 - comparison to T4 DNA ligase 49
 - 3'-labeling by ligation of 5'-³²P-pCp 181–183, 213, 337–338, 383–384, 980, 1130
 - ligation of large RNA molecules 76–79
 - mechanism and substrate specificity 46–49, 71–73
 - properties 46
 - protocols 80–84, 817
 - site-specific 3'-terminal modification 233
 - T4 RNA ligase 1 (T4 Rnl1, RnlA) 46–50, 70–73, 75–76, 78–82, 862, 865, 869, 879
 - T4 RNA ligase 2 (T4 Rnl2) 46–47, 49, 68
 - T4 Rnl2 “Truncated K227Q” 46–47, 862–863, 865, 868, 872, 879, 889
- T7 DNA Polymerase
- 3'-³²P-labeling with [α -³²P]dATP 182–183
- T7 RNA Polymerase (T7 RNAP)
- bacteriophage promoters (T3, T7, SP6)
 - conformational heterogeneity
 - – analyzed by TGGE 436–437
 - – RNA renaturation to generate homogeneous conformers 213, 224, 412, 485, 980–981, 989–990, 1005, 1008, 1124
 - endogenous DNase and RNase activity 9
 - *in vitro* transcription
 - – annealed oligonucleotides as template 12
 - – for AFM studies 537
 - – for RNA probing 213
 - – for SAXS studies 412
 - – for SELEX studies 1099, 1108, 1112–1113, 1153, 1179
 - – homogeneous 5'-/3'-ends and 5'-end groups 5–6, 29–42
 - – homogeneous 5'-/3'-ends, RNase H approach 54–56, 63
 - – homogeneous 3'-ends by 2'-O-methyl residues at 5'-end of template strand 53–55
 - – incorporation of phosphorothioate analogs 335–337, 369–372, 380–381
 - – internally digoxigenin-labeled (DIG-11-UTP) 100–101
 - – internally ³²P-labeled 14, 16, 747, 980, 1179
 - – large RNA amounts for structural studies 12–13, 111–113
 - – low or no product yield 20–21, 1107
 - – nucleotide analogs for internal and 5'-terminal incorporation 6–8
 - – purification of transcripts 18–20, 61–63, 336, 381–382, 1179–1180

- RNAs with biased nucleotide composition 14–15
- transcripts initiated with guanosine, 5'-GMP or 5'-ApG 11–12
- with 5'-cap structures 17–18, 51–52, 61
- with 5'-terminal GMPS modification 233–234, 240
- with 2'-fluoro-modified nucleotides 16–17
- with unmodified nucleotides 9–15, 60–61, 980
- mutant T7 RNAP
 - Y639F 7–8, 16–17, 21, 335–336, 348–349, 351, 371
 - Y639F/H784A 7–8, 349, 371
- NAIM analysis of T7 RNAP termination 347–351
- rapid T7 RNAP preparation 21–25
 - procedure 22–24
- template types 4–5
- plasmid template preparation 108–111
- TagIt*-affinity purification 935
- Taq* (DNA) Polymerase 37, 750, 753–754, 768, 772, 803, 811, 813, 816–817, 865, 870, 879, 892, 1118, 1120–1121, 1149, 1152, 1173–1174, 1179, 1218
- TargetRNA2 algorithm (bacterial sRNA target identification) 737
- TBDMS (2'-O-) RNA synthesis chemistry 129–132, 136
- temperature-gradient gel electrophoresis (TGGE) 427, 462
 - application examples 433–443
 - analysis of RNA mixtures in a single TGGE experiment 432–435
 - mutant RNA analysis 438–440
 - protein-RNA complex detection 439–441
 - structure transition analysis 435–438
 - experimental conditions optimization 430–431
 - electric field 430
 - gel matrix pore size 430
 - ionic strength and urea 431
 - handling 429–430
 - instruments 429
 - interpretation rules 431, 433
 - principle 428–429
- template-stripped gold (TSG) method (force spectroscopy AFM) 543
- terbium(III) footprinting 255–256
 - application example HDV ribozyme 261–264
 - of long RNAs by primer extension 259–261
 - of short RNAs 256–259
 - 3'-terminal attachment of fluorescent dyes and biotin 117–125
 - background and chemistry 118–120
 - protocols
 - biotin attachment 120–121
 - fluorescence labeling 121
 - labeled RNA purification and quality control 122–123
 - RNA 3'-dephosphorylation 121
 - terminal (deoxynucleotidyl) transferase (TdT) 76
 - tertiary (3D) structures. *See also* crystallization of RNA, NAIM, NAIS and SAXS
 - probing, RNA
 - enzymatic 205–206
 - Tb(III) 255–266
 - Pb²⁺ 269–280
 - Fenton cleavage 301–314
 - NAIM/NAIS 333, 375
 - temperature-gradient gel electrophoresis 427–442
 - UV melting 462, 466
 - RNA crystallization 492
 - FRET 501, 507
 - AFM 531
 - prediction/modeling 633, 667. *See also* 3D architectures, Assemble2, S2S and molecular dynamics (MD) simulations of RNA systems
 - molecular dynamics 687
 - RNA-protein interaction 975–976, 978, 981
 - aptamers, *in vitro* selection 1130, 1148, 1150, 1155, 1171
 - proteins, Fe(II)-mediated cleavage 400
 - tetracycline, mapping of binding sites on RNA by drug-directed Fenton cleavage 310–312
 - thin-layer chromatography (TLC) 62, 175, 184, 190, 192, 194–199, 201, 236
 - thiophilic metal ion rescue
 - kinetic analysis 293–294
 - RNA phosphorothioate modifications 286–290
 - 6-thioguanosine. *See* nucleoside analog
 - 4-thiouridine. *See also* nucleoside analog, PAR-CLIP and photocrosslinking/UV crosslinking
 - postsynthetic labeling of modified RNA 152, 157, 160, 164

- 4-thiouridine. *See also* nucleoside analog, PAR-CLIP and photocrosslinking/UV crosslinking (*contd.*)
- UV crosslinking of 4SU-labeled cells 883–884
- three-way ligation 51–52, 58
- tiling (micro)arrays 720–721, 725, 787, 795–796
- T-jump technique 468
- 2'-thionocarbamate (TC) chemistry method 130
- TLC. *See* thin-layer chromatography
- tobramycin 957, 963, 987–989, 991–992, 1006, 1117
- aptamer 735, 935, 957, 963, 987–989, 1006, 1008
- tobramycin-sepharose 987–989, 991–992, 1006, 1008
- toeprint assay, 30S ribosomes 741–742, 772–776
- toll-like receptor (TLR) 1257–1258, 1265
- TOPHAT sequence read alignment program 823, 826
- total internal reflection fluorescence microscope, prism-based (prism-TIRF) 500, 511, 517
- transcription elongation complex (TEC) 3, 347, 349
- transcription priming 17, 61, 231, 233–234
- priming of reverse transcription 901, 903, 995
- transcriptome 789, 794, 853
- biocomputational analysis 727, 821, 823, 826, 830
- (d)RNA-seq 724, 821, 823, 826, 830, 1025
- experimental RNomics/RNPomics 801, 814
- genomic SELEX 1202
- meta-transcriptome data 727
- microarray 720, 796
- miRNA targeting 853
- sRNA pulse expression and transcriptome analysis 733–734
- transcriptome-wide identification of protein binding sites on RNA (PAR-CLIP) 877
- transduction 730, 1233–1235
- transfection 837, 844–845, 849, 1089, 1208, 1211–1214, 1216, 1221, 1223, 1232–1234, 1255–1256, 1260
- TransTermHP algorithm (prediction of bacterial transcription terminators) 791
- trimethylamine oxide (TMAO) 463
- tris-(hydroxypropyltriazolylmethyl)amine (THPTA) 509–510, 512
- trityl-off RNA 145, 361
- desilylation 137–138
- failure sequences 141
- trityl-on RNA 137, 139, 145–146
- desilylation 138
- detritylation 142
- reversed-phase HPLC purification 140–142, 144
- transfer RNA (tRNA) 355
- abstract shape analysis 579, 584, 588
- affinity purification of RNPs 1050
- AFM 532–533, 536–537
- 3'-biotinylation/tagging with fluorescent dyes 118, 123
- crystallization 492
- homogenous 5'/3'-ends 32–35, 38, 40–42
- homology search for small ncRNAs 619, 622, 627–628
- identification and characterization of small ncRNAs in bacteria 719
- – microarray 720
- – RNA-seq 721–722, 724
- – cell-free translation 741–742
- – toeprinting assay 741, 772–775
- immunopurification of RNPs 1021, 1047
- isolation of RNA binding proteins 921–932, 946
- metal ion coordination 287–290
- – incorporation of phosphorothioates 291–293
- – thiophilic metal ion rescue 293–294
- miRNA targets 1092
- molecular dynamics (MD) simulation 692, 700
- NAIM 355, 372–375, 377–381, 383–384, 387, 389–392
- quality control of total cellular RNA, Northern blotting 95
- RNA ligation, *in vitro* production of full-length tRNA 50, 70–71, 74–81, 84
- RNA-protein interactions and RNA structure in RNPs 981–982, 984, 989–999, 1007, 1009
- RNA sequence and modification determination by radiolabeling 175–176, 182, 185, 187–188, 200
- – alkaline hydrolysis 191
- – chemical sequencing 188–189
- – end group analysis 190–191
- – enzymatic hydrolysis 191–192
- – production of labeled random RNA fragments 199
- – radiolabeling 180–181
- – TLC 192, 196

- RNomics/RNPomics 802
 - screening of genomes for known RNA genes or motifs 595–596
 - sedimentation analysis of RNPs 1059, 1063
 - shRNA vectors 1223
 - smFRET 501, 504–505, 515
 - – dye labeling 508
 - structure and folding analysis by SAXS 409, 411, 416
 - structure prediction
 - 2D 551
 - 2D and 3D 644–645, 651–652, 655
 - structure probing by crosslinking 242, 244–246, 248–249
 - 5'-end-labeling by transcription priming 233–234
 - structure probing by Fenton cleavage 302
 - structure probing by Pb²⁺ cleavage 269
 - – probing of metal ion binding sites 271, 273
 - structure probing by Tb³⁺ footprinting 264
 - structure probing *in vitro* 205, 211, 214, 216, 218–219
 - temperature-gradient gel electrophoresis 434, 439, 441
 - UV melting 446, 449, 461–462
 - tRNA processing. *See* RNase (RNase P, RNase Z), tRNase Z and tRNA nucleotidyltransferase
 - tRNA nucleotidyltransferase 182, 922
 - tRNA synthetase 1082
 - aminoacylation of tRNA 29, 33, 79, 176, 504–505, 922
 - aminoacyl-tRNA 501
 - aminoacyl-transferase ribozyme 264
 - tRNA splicing 922–923, 927, 930
 - adenylyltransferase 923, 931
 - 2' phosphotransferase 923
 - endonuclease 923, 927
 - ligase 922–924
 - – purification 930–931
 - tRNase Z (RNase Z)
 - bioinformatic studies 628
 - purification from wheat germ using RNA affinity matrices 922, 924–925, 927–929
 - tRNAdb (transfer RNA database) 816
 - Turnip Yellow Mosaic Virus (TYMV) RNA 531, 533
- u**
- UCSC Genome Browser 620, 625, 816, 822–823, 826, 829
 - ULSTM Fluorescent Labeling Kit, for Agilent arrays 796–797
 - ultracentrifugation 427, 1041, 1055
 - cesium chloride/sulfate 1056, 1061–1064
 - glycerol gradient 814–816, 1056, 1058–1059
 - sucrose gradient 1061
 - – polysome profiling 839
 - ultra performance liquid chromatography (UPLC) 143–144
 - ultraviolet (UV) crosslinking/irradiation
 - RNPs *in vivo* 877–878, 883–884, 899–901, 905–906
 - RNP structure analysis 977, 999, 1003–1005, 1008–1009, 1026, 1030–1035, 1042–1044
 - – naked RNA control 1045
 - RNA structure probing 240–247
 - RNA to nylon membrane 91, 746, 808
 - ultraviolet (UV) melting studies of RNA 439, 445
 - definitions and nomenclature 446–447
 - equilibrium and non-equilibrium 459–460
 - experimental solution to sloping baseline 468
 - – experimental data processing 472–473
 - – heating rate and data sampling 471
 - – pH variation and buffers 468–470
 - – RNA degradation 470–471
 - – softwares 473
 - experiments for thermodynamic studies 449
 - – bimolecular transitions 452–459
 - – melting temperature definition 450
 - – unimolecular transitions 451–452
 - melting curves of large RNAs, thermodynamic information 461–462
 - parameters influencing melting temperature 462–463
 - practical problems
 - – evaporation during heating 463–464
 - – sloping baseline 464, 466–467
 - self-complementary sequence pitfalls 460–461
 - two-state approximation 459
 - UV absorption
 - – characteristics of nucleic acid bases 447–449
 - – physical basis 445–446
 - ultraviolet (UV) shadowing 13, 19, 41, 53, 62–63, 84, 112, 163, 336, 382, 484, 1045, 1123, 1214
 - ultraviolet (UV) spectroscopy, for RNA
 - chemical reaction monitoring 164

- ultraviolet (UV) spectroscopy, for RNA (*contd.*)
 - duplex melting (thermal denaturation) 159, 164, 166
 - GMPS 236–237
 - RNA quantification 60, 1108, 1279–1281
- UNAFold web server (UNAFold program package) 549, 555–557
- unlocked nucleic acid (UNA) 1246–1249, 1251–1254, 1259, 1262, 1264
- URA3* gene (*ura3*) 1070–1071, 1081
- v**
- Varkud satellite (VS) ribozyme 33, 500
- vbFRET software package, smFRET analysis 507
- Vienna RNA package/Web Services/Websuite 549, 620–621, 673, 811
- w**
- Watson-Crick base pair 66, 91, 301, 463, 596, 606, 633, 650, 671, 684, 689, 1100
 - disruption of H bonds, UV melting 449
 - NAIM 371
 - non-Watson-Crick base pair 206, 255, 493, 551, 596
 - reverse transcription 1034
 - RNA probing 210–211, 217, 220
 - stacking 549
- Weblogo program 620
- wiggle track file format 826, 829
- Wincott deprotection protocol 359
- wobble base pair 549, 560, 633, 645, 671, 684, 692
- worm-like chain (WLC) model 534
- x**
- X-ray
 - crystallography/structure of RNA
 - – crystallization of RNA 481–495
 - – 2D and 3D RNA structure prediction 633, 642, 644–645, 652, 656
 - – ligand-aptamer complex 1188
 - – molecular dynamics (MD) simulations 688–690, 692–693
 - NAIM 374–375
 - – RNA-metal ion interaction 285, 287, 301, 308
 - – RNA structure probing 206, 211, 241, 264, 269, 301
 - – RNA synthesis 132, 143, 152
 - – RNP 978
 - – SAXS 416
 - – smFRET 502–503
 - film 63, 95, 194, 197, 201, 403, 437, 867, 911, 978, 993, 996, 1000, 1037, 1047, 1092, 1180–1181
 - generation of hydroxyl radicals 301, 421–422
 - small-angle X-ray scattering (SAXS) 407–423
 - – RNA damage 421–422
- y**
- yeast (*Saccharomyces cerevisiae*)
 - mitochondrial ai5 γ group II intron 344
 - poly(A) polymerase 182, 339
 - protein A system 1030
 - pyrophosphatase 10, 234
 - small nuclear ribonucleoproteins (snRNPs), spliceosomal complexes 957
 - – isolation of native complexes 963–971
 - Snu13 protein 982
 - three-hybrid system 899, 1067–1083
 - total RNA 279, 1157–1158
 - tRNA 38, 78, 123, 181, 185, 188–191, 199, 269, 271, 409, 644–645, 655, 700, 939, 979, 981–982, 984, 989–992, 994–996, 998, 1007, 1019, 1021
 - – intron-containing 923
 - U3 snoRNA/snoRNP 983, 999, 1002
 - U5 snRNA 996
 - [U4/U6.U5] tri-snRNP 996
- z**
- zero-length crosslink 1030–1031