

Index

a

- ABC (acid binding capacity) 57, 587
 abiotic factor 279
 abundance
 – affected by hydrological chain 578
 – affected by thermal chain 578
 – and discharge pattern 514
 – bacterial 134ff.
 – monthly 558ff.
 – prediction 561f.
 – yearly 557ff.
 β -N-acetylglucosaminidase 166, 170ff.
Acidobacteria 151ff.
Actinobacteria 145f., 148ff., 181f.
 activity trap 206
Adephaga 309
 adult
 – feeding 331
 – upstream movements 390
Agabus spp. 310
Agapetus 231
 – *fuscipes* 329ff., 344ff., 484, 594
 – – case construction 432
 – – egg mass abundance in the stream 430
 – – egg number per female
 – – larval mass and case length 424
 – – number of larval instars 423
 – – molt increments 426
 – – oviposition behavior 429
Agriotypus armatus 226
 air temperature 5, 30ff., 52, 279, 529
 – Erlenbach 545
 algae 100
 – variation in algal biomass 108
 algal assemblage 100ff., 122
 algal communities 100f.
 algal drift 99
 alkalinity 58ff.
 allometric growth 228, 253, 287,300
Alphaproteobacteria 147ff.
Ameletus celer 298
 7-amido-4-methyl-coumarin (MCA) 166
 ammonium 67ff.
Amphinemura standfussi 264, 579
Anacaena spp. 315
Ancylus fluviatilis 215f.
 animal–microbial interactions 418
 Annelida 211f.
Anomalopterygella chauviniana 231, 356
Apatania
 – *fimbriata* 230, 291, 330ff., 350ff., 484, 562, 594
Aphelocheirus aestivalis 224
 aquatic Diptera 369, 500ff., 526ff., 548, 595
 – Brachystomatidae 394
 – Ceratopogonidae 377
 – Chironomidae 373ff
 – Dixidae 380
 – Dolichopodidae 396
 – Empididae 395
 – Ephydriidae, Muscidae, Faniidae 403
 – Limoniidae and Peditiidae 371
 – Lonchopteridae 403
 – Psychodidae 392
 – Ptychopteridae 394
 – Rhagionidae 400
 – Simuliidae 381ff.
 – Stratiomyidae 400
 – Syrphidae 402
 – Tabanidae 401
 – Thaumaleidae 381
 – Tipulidae 370
Argyroneta aquatica 217
 Arthropoda 216
 artificial neural network (ANN) 554
 – modeling discharge 554
 ash free dry mass (AFDM) 75

- assemblages 100ff.
 - algal assemblages in streams 100
- Aulodrilus pluriseta* 212
- b**
- bacteria 131ff., 589
 - food web system of the Breitenbach 567
 - role in streams 131
 - taxonomic distribution of strains 146
- bacterial abundance 134ff.
 - different habitats 134
 - effects of temperature and discharge 139ff.
- bacterial biomass 141
- bacterial community composition 142ff.
 - cultivation 144
 - genetic diversity 154
 - methodological approaches 142
 - sediment 152ff., 161
 - taxonomic composition 151
 - water 152ff., 161
- bacterial production 182ff.
 - annual 193
 - importance of bacterial carbon production in streams 182
 - influence of temperature and discharge 190
 - methodological approach 183
 - seasonal fluctuation 184
 - seasonal pattern 187
 - stream 182
- bacterial species
 - Breitenbach 160
- Bacteroidetes* 145ff., 182
- Baetis* 244ff.
 - *rhodani* 227ff., 241ff., 560, 572ff.
 - *vernus* 227ff., 241ff., 305ff., 495, 572
- bank-full 42
- base flow 36
- base flow index (BFI) 36
- behavior 244, 405ff.
 - competitive 471
- behavioral constraints 244
- benthic diatom assemblages 112
- benthos 228
 - mayfly larvae 228ff.
 - Plecoptera larvae 258ff.
 - Trichoptera 348ff.
- Betaproteobacteria* 145ff., 182
- biofilm structure 135
- biomass 122, 134
 - algal 108
 - bacterial 141
 - meiofaunal 569
- bionomics 318
- Biozönotische Gliederung der Fließgewässer 3
- biting midges 377
- blackfly 207, 233, 246, 381ff.
 - species distribution in space 384
 - species distribution in time and life cycle 385
 - species of the Breitenbach 384
- body size and weight variation
 - seasonal 299, 304f., 344, 583
 - spatial 340
 - temperature dependence 476
- Brachycera 369, 401
- Brachyptera* spp. 256, 258, 286, 296
- Breitenbach 5ff.
 - ammonium 67ff.
 - annual irradiance 23
 - average annual precipitation 25
 - bacteria in the food web system 567
 - bacterial species 160
 - chloride 71
 - climate and weather 21ff., 577
 - comparison with the Erlenbach 543
 - conductivity 55, 62, 545
 - diatom occurrences 101ff.
 - discharge 31ff., 489
 - dissolved organic matter 84ff.
 - ecosystem 487ff.
 - EPT community 522
 - fauna 195ff.
 - flood event 40ff.
 - geographical and physical characteristics 6
 - global change 570
 - high water 41
 - hydrological measures 37ff.
 - irradiance 25
 - life cycle study 358
 - low water 44
 - mayfly faunas 229
 - microbial metabolic activity 163
 - monthly precipitation 26
 - nitrate 66f.
 - nitrogen 64
 - nutrient 67
 - organic matter budget 564
 - outputs of POM 83
 - oxygen 60ff.
 - particulate organic matter 74ff.
 - pH 59f.
 - phosphate 69f., 179f.
 - precipitation–discharge model 555
 - primary producer 99, 588

- sampling sites 18
- sandy sediments as habitat 495
- sandy streambed area 487
- scraper 596
- settlement history 15ff.
- silicate 70
- soluble inorganic element 64
- species distribution 529
- species list 227, 254
- standing stocks of POM 83
- sulfate 71
- water temperature 476
- yearly precipitation 27
- Breitenbach valley
 - air temperature 32ff.
 - catchment 14
 - discharge 34ff.
 - flora and fauna 14ff.
 - global irradiance 23
 - precipitation 24, 38
 - runoff 36ff.
 - vegetation profile 17
 - water saturation 44
 - wind 27ff.
- Brook's rule 253, 298
- Brychius elevatus* 311
- buffering capacity 56
- Bunter sandstone 11
- Bythinella compressa* 215
- C**
- caddis fly 197ff., 327ff., 530, 562
 - adults in traps along the Breitenbach 340
 - scraper 483
- Calopteryx splendens* 223
- carbohydrate 74, 91ff., 175f., 180ff.
- carbon budget 559
- carbon flow 133, 564, 590
 - stream 133
- carbonate system 59
- CARD-FISH (catalyzed reporter deposition FISH) 144ff.
- Carychium minimum* 216
- catchment
 - Breitenbach 7ff.
- Centropilum luteolum* 227f., 241, 502
- Ceratopogonidae 220ff., 377ff.
- Chaetopteryx villosa* 330ff., 361ff., 478ff., 583
- Chelicerata 216
- chemistry 56
 - buffering capacity (ABC) 55
 - pH 60
 - oxygen 62
 - soluble inorganic elements 64
 - chemoautotrophic production 99
- Chironomidae 217ff., 312, 372ff., 500ff.
- Chordata 404
- Chironominae 374ff.
- chloride 71
- Chloroperlidae 270
- chitinase 154, 173, 180ff.
- ciliates 210
- Ciliophora 209
- Clavatospora longibrachiata* 163
- climate 5, 80, 585
- coarse particulate organic matter (CPOM)
 - 74ff., 339, 480
 - amount of inputs 78
 - composition of inputs 81
 - seasonal distribution of inputs 76
- Cocconeis placentula* 123
- Cochlicopa lubrica* 216
- coexistence, competitive 471
- cohort splitting 246, 259, 266, 275ff., 295, 325
- Coleoptera 309, 595
- Collembola 219
- colonial web 368
- community 505ff., 529, 547
 - data treatment 505
 - Erlenbach 546
 - statistics 527
- community diversity 503, 519, 530
- community measures 537ff.
- community respiration (CR) 127ff.
- community state 514ff., 598
 - multiple 517
- competition 308, 348, 388, 390
 - exploitative 472, 475, 484f.
 - fights 395
 - for food 456f., 459ff.
 - for space 411
 - interspecific 340, 346, 445, 455, 462, 483
 - intraspecific 305, 340, 346, 405, 410f., 474, 483
- competition avoidance 272, 297
- competitive exclusion 266
- congener 295
- Cordulegaster boltoni* 223
- correlation coefficient 52f.
- correspondence analysis 534
- crane fly 370
- cross-section measurement 46
- Crustacea 218
- Cyphon* 322ff.
 - *coarctatus* 323
- Cytophaga* 145, 149

d

dance fly 395
 data treatment 505
Deltaproteobacteria 151ff.
 denaturing gradient gel electrophoresis (DGGE) 181, 144
 detrital energy sources 73ff., 587
 detritus 73
Dianous coeruleus 312
 diapause
 – egg development 296f.
 diatom occurrences 101ff.
 – spatial and temporal patterns in the Breitenbach 101
 diatom species 109
Diatoma mesodon 103f.
Dictyogenus fontium 228ff., 300f.
 dimorphism
 – sexual 300
 Diptera
 – aquatic – see aquatic Diptera
 – life cycle characteristics 532
 discharge 34, 282, 489, 528
 – bank-full 42
 – base flow 36
 – base flow index (BFI) 36
 – falling limb 34
 – flood event 40ff.
 – flow duration curve (FDC) 39
 – high water 42
 – low water 44
 – rising limb 37
 – Erlenbach 545
 – modeling 554
 discharge pattern 503ff., 525ff.
 – discriminant analysis 517
 – EPTD community 526
 – monthly abundances 560
 – species 531
 discriminant analysis 517, 539
 dispersion 390
 dissolved organic carbon (DOC) 84ff., 494, 558
 dissolved organic matter (DOM) 73, 84ff.
 – composition 84, 91
 – concentrations and dynamics in the Breitenbach 84
 – groundwater 86ff.
 – inputs and outputs to and from the Breitenbach 89
 – interstitial water 88f., 484
 – stream water 84ff.
 diurnal foraging 367, 461ff.
 Dixidae 220ff., 380

Dolichopodidae 220ff., 370, 396
 drift 246, 250, 310, 317ff., 369, 377, 382ff., 404, 410, 438
Drusus annulatus 329ff., 353, 484, 572ff., 594
 Dryopidae 320
Dugesia gonocephala 211f.
 Dyar's rule 253, 298
 Dytiscidae 310

e

ecological inferences 115
 ecological segregation of grazers 471
 ecophysiology 405ff.
 ecosystem 487ff., 598
 – Breitenbach 487ff.
 egg
 – numbers 334
 – origin of biomass 288
 – size 288
 – volume 335f.
 egg development 250, 292, 335
 – diapause 250, 296f.
 – temperature and photoperiod 292
 Elmidae 315
Elmis 315ff.
 – *maugettii* 315ff., 595
Elodes 322ff., 596
 emergence
 – cues 272
 – diurnal emergence periodicity 390
 – shift along the stream 284
 – short-term control by abiotic factors 279
 emergence dates 573ff.
 – accuracy and possible sources of error 573
 – shift 573
 emergence pattern 239ff.
 – aberrant 241, 244
 – standard 239
 – unimodal 276
 – plurimodal 276
 emergence time
 – shift 284, 575
 emergence trap 206ff., 591
 – function 203
 – silted 498
 Enchytraeidae 212
Enochrus 313
 environment 585
 environmental variable 508, 528
 – Erlenbach 545
Ephemerella 249f., 499
 – *ignita* 233ff., 304, 579
 – *mucronata* 227ff., 241, 304

- Ephemeroptera 226ff., 499, 513, 526ff., 592
 – fertility 248
 – instar 252
 – life cycle characteristics 532
 – life histories 231
 – mean annual specimen numbers 518
 – pairwise correlation 515
 – shift in emergence time 575
 Ephemeroptera, Plecoptera, and Trichoptera (EPT) 226, 228, 506ff.
 – community of the Breitenbach and environmental variation 522
 – mean annual specimen numbers 518
 – pairwise correlation 515
 Ephemeroptera, Plecoptera, Trichoptera, and Coleoptera (EPTC) 226
 Ephemeroptera, Plecoptera, Trichoptera, and Diptera (EPTD) community 526ff.
 – life cycle characteristics 532
 Erlenbach 543f., 599
 – air and water temperature 545
 – comparison with the Breitenbach 543
 – conductivity and pH 545
 – discharge 545
 – environmental variables 545
 – Ephemeroptera, Plecoptera, Trichoptera, Neuroterioidea, and Diptera species 548
 extracellular enzyme 165
 extracellular enzyme activity 164ff.
 – environmental control and regulation 175
 – function in streams 164
 – stream habitat 168
- f**
 false crane fly 394
 fauna 591
 – Breitenbach 195ff.
 – sampling method 196
 fecundity 248, 287, 333
 feeding strategies of trichopteran grazers 471
 female longevity 289
 filterer 227, 328, 356, 382ff., 404
 fine particulate organic matter (FPOM) 74, 339, 480
Firmicutes 148ff., 182
 first instar 299
 – larva 251
Flagellospora curvula 163
Flavobacterium 145, 155, 160
 flood event 40ff.
 flow duration curve (FDC) 39
 fluorescence *in situ* hybridization (FISH) 145
- food 461, 480
 food chain 564
 – in aquatic ecosystems 132
 food choice 462ff.
 food quality 453
 food web system 131
 – bacteria 567
 Fulda
 – crustacea 218
 – mayfly fauna 229
 – species exchange with Breitenbach 201, 223, 228, 244ff., 329, 348, 384ff., 404
 functional feeding group 256, 477
 fungal community 162ff.
 fungal production 194
 fungi 131ff., 162ff.
 – role in streams 131
- g**
Gammaproteobacteria 145ff., 182
Gammarus 218, 481
 Gause's principle 272
 global change 570, 600
 – shift in emergence time 568ff.
 – emergence dates and thermal regime 570
 global irradiance 21
 Glossosomatidae 339
 β -glucosidase 165ff.
 grain size
 – analysis 112ff.
 – distribution 489
 grazers and biofilms 467
 gross primary production (GPP) 127ff., 557
 Gyrinidae 311
- h**
 habitat 134, 168
 – epiphytic 136f.
 – planktonic 168
 – sediment 134ff., 161, 168ff., 184ff., 484
 – suspended 137ff., 168ff.
 habitat templet concept 504, 563
 Haliplidae 311
Haliplus 311
 harsh–benign concept 504, 563
 hatching 295
 Helophoridae 313
 Heteroptera 224
 heterotrophic production 189ff., 198, 560
 high water 41
 Hydrachnidia 217
 hydrological chain 586
 hydrological measure 46
 Hydrophilidae 314f.

- Hydrophorinae 398
Hydropsyche spp. 290, 333, 348, 356
Hymenoptera 225
hyporheic habitat 487
- i**
- incubation period 250ff., 292ff., 336ff., 369
input average 42
input average 42
insect
– abundance 554
– aquatic 220ff., 496ff., 539
– collecting 202
– errors during trap operation 205
– evaluation 208
– minor taxa 219
– seasonal size decline 304ff.
– trap 202f.
insect emergence
– timing 570
Insecta 219
insecticide poisoning 201
instar 251, 299ff.
– number 252, 303
interference 471
intermediate disturbance hypothesis 504, 563
interstitial water 88ff.
– chemical characteristics 490
– dissolved organic carbon 88f., 484
irradiance
– global 21ff., 283
Isoperla goertzii 258ff., 267ff., 296
- I**
- Laccobius* 314
– *striatulus* 314
larval abundance 406
larval age structure 258, 405ff.
larval biology 323
larval development 338, 360, 405ff.
larval drift 246
larval functional feeding groups 227
larval population structure
– benthos 324
larval predator 360f.
larval retreats 405ff.
Lepidoptera 327
Lesteva 312
– *longoelytrata* 312, 327
Leuctra 266ff.
– *braueri* 579
– *digitata* 303ff.
– *geniculata* 303
– *inermis* 255, 266
– *nigra* 269ff., 291, 303, 502, 581
– *prima* 254ff., 301, 581
Leuctridae 266
life cycle 324
– analysis 201, 358ff.
– *Agapetus fuscipes* 421ff.
– *Apatania fimbriata* 358ff.
– *Chaetopteryx villosa* 361ff.
– loss rate 246, 361
– *Sericostoma personatum* 365ff.
Limnephilidae 333
Limnodrilus hoffmeisteri 212
Limnologische Flusstation Schlitz
– history 1ff.
Limoniidae 220ff., 370ff.
longevity
– female 289
low water 44
- m**
- macrophyte 16f., 74ff., 99, 124
macrozoobenthos 196
– collecting 196
mate finding 246, 285
mating 246, 285
mayfly 227ff.
– development 246ff.
– emergence 235
– growth 251
– larvae 228
– reproduction 246
Megaloptera 224
meniscus midges 380
Meridion circulare 117f.
metamorphosis 253, 300
– early 254
metazoan size distribution 196
4-methylumbelliferone (MUF) 166, 185
Micrasema longulum 353
microhabitats 115
microbial food chain 132
microbial loop 132, 183
microbial metabolic activity 163
– measurements in the Breitenbach 163
Microcara testacea 323
Microsporus acaroides 309
model
– seasonal effects of variables on species 559
modeling 599
modeling discharge 554
– artificial neural network (ANN) 554, 599
– yearly abundance patterns 554
– yearly species abundance 550

molecular biology perspective 179
 Mollusca 214ff.
 Monard's principle 272
 morphological measure 46
 mosses 125f.
 moth fly 392f.
 MUF- β -glucoside 167ff.
 Myxophaga 309

n

Naididae 212
 Navicula 103
 – *capitata* 105ff.
 – *gregaria* 106ff.
 – *hungarica* 106f.
 – *protractoides* 106ff.
 – *rhyngocephala* 105f.
Nebrioporus elegans 310
 Nematocera 369, 392
 Nematoda 212ff.
Nemoura 259
 – *cinerea* 302
 – *flexuosa* 256
 – *marginata* 286, 295
 Nemouridae 259
Nemurella pictetii 199ff., 246, 259ff., 275ff.,
 291ff., 307, 572
Neomys fodiens 359
Nesovitrea hammonis 216
 net primary production (NPP) 126
 Neuroptera 225
 Neuropteroidea 540ff.
 nitrate 66f.
 nitrogen 64
 nonreactive phosphorus (NRP) 180
 nutrient 67

o

Odeles 322ff.
 – *marginata* 322ff., 596
 Odonata 221
Odontocerum albicorne 339, 358
Oligoneuriella rhenana 227
 ordination 517, 598
Orectochilus villosus 312
Oreodytes sanmarkii 310
 organic carbon (OC) 75
 – input 564
 – output 565
 organic matter (OM) 75
 – balance 590
 – Breitenbach 564
 – budget 557
 – stream 73

Orthoclaadiinae 374ff.
 Orthoclaadiini 217, 377ff.
 osmoregulation 220ff., 324
Osmylus fulvicephalus 225, 359
Oulimnius tuberculatus 315f., 595
 oviposition 249, 290, 332
 – modes 291, 432
 – substrata 244
 – uncoupling from emergence 244
 – upstream oviposition flights 290
 oxygen 60ff.

p

Paederus 312
Palingenia longicauda 247
Paracladopelma camptolabis 377
Paraleptophlebia submarginata 228, 502
 parthenogenesis 248, 292, 336
 particulate organic carbon to total fine
 particles (POC/TFP) 496
 particulate organic matter (POM) 73ff.,
 189, 558
 – composition 81
 – content 188
 – inputs to the Breitenbach 75, 124f., 557
 – outputs from the Breitenbach 83, 558
 – standing stocks 83
 – stream 74
 – total inputs 82
 patch dynamics concept 563
 Pediciidae 371f.
 peptidase 165ff.
 periphyton (*Aufwuchs*) 99
 periphyton primary production 127
Perla marginata 256
Perlodes
 – *dispar* 255
 – *microcephalus* 255
 Perlotidae 258ff.
 Perloidea 258
 pH 59, 283
 – Erlenbach 545
 – flowing current 59
 pheromone 332
 phosphatase 165ff.
 phosphate 69f., 180
 photoautotrophic production 99
 phytoplankton 99, 127
Picea spp. 581
Pinnularia borealis 115
Pinus 581
Pisidium 215
Placoneis clementis 103ff.
 planktonic primary production 127

- Planothidium lanceolatum* 102ff.
 - plant biomass
 - average composition 165
 - Platambus maculatus* 310
 - Platyhelminthes 211
 - Plea minutissima* 224
 - Plecoptera 254, 272f., 293ff., 499, 513, 526ff., 593
 - development 297, 593
 - egg size 287
 - emergence 272, 593
 - Erlenbach 548
 - fecundity 287
 - growth 297
 - life cycle characteristics 532
 - mean annual specimen numbers 518
 - morphological changes during last instar 302
 - pairwise correlation 515
 - reproduction 284, 593
 - sex ratio 284
 - shift in emergence time 575
 - Plectrocnemia conspersa* 333, 356, 368, 583
 - Polycentropodidae* 339
 - Polynucleobacter* 162
 - Polyphaga 312
 - polysaccharides 74
 - dissolved 82f.
 - enzymatic degradation 164
 - Pomatinus substriatus* 322
 - Potamophylax* 332ff.
 - *cingulatus* 342, 578ff.
 - *luctuosus* 578
 - *nigricornis* 333
 - pre-metamorphic instar 300
 - precipitation 24ff., 39, 528
 - average annual 25
 - monthly 26
 - yearly 27
 - precipitation–discharge model
 - Breitenbach 555
 - primary producer 99ff., 588
 - stream 99
 - primary production 126
 - Prionocyphon* 327
 - *discoideus* 324
 - Prosimulium tomosvaryi* 384ff.
 - protandry 285
 - Protonemura* 261
 - *auberti* 256ff., 276ff., 304, 581
 - *intricata* 263, 579
 - *meyeri* 263ff., 580
 - *nitida* 255
 - Protozoa 209
 - Protzia eximia* 217
 - Psychodidae 392
 - Pteronarcys* species 303
 - Pterothecae 301
 - Pterygota 221, 591
 - Ptychoptera*
 - *albimana* 394
 - *paludosa* 367, 394, 496ff., 595
 - *townesi* 496
 - Ptychopteridae 394
- r**
- rainy day 529
 - reproduction 331
 - resource limitation 484
 - resource overlap 453
 - respiration 193, 220, 313ff., 381, 393ff., 412ff., 436
 - Rhithron–Potamon concept (RPC) 526
 - Rhyacodrilus coccineus* 212
 - Rhyacophila* 332
 - *fasciata* 212, 329ff., 356ff., 572ff., 594
 - Rhyacophilidae* 339
 - Riolus subviolaceus* 317
 - River Continuum Concept (RCC) 526
- s**
- saccharide 94
 - sand trap 498
 - sandy sediment
 - habitat 495
 - sandy streambed area
 - Breitenbach 487
 - Scirtidae 322ff.
 - scraper 483ff.
 - Breitenbach community 596
 - crawling velocity 438
 - dietary composition of epilithic Trichoptera 445ff.
 - interactions with periphyton 442ff.
 - – comparison between species 446f.
 - – gut content and periphyton composition throughout the year 447
 - – variability along the stream 451
 - secondary production 198
 - estimates 207, 210
 - sediment
 - core 185
 - stream 11, 487
 - selection criteria 572
 - Sericostoma* 320, 335
 - *personatum* 329ff., 365ff., 502, 583

- Serratella ignita* 233
 sex ratio 237, 284, 331
 sexual
 – dimorphism 297ff., 349, 423, 430
 – size difference 478
 Shannon–Wiener diversity 519
Sialis fuliginosa 224
 silicate 70f.
Silo pallipes 578
 Simuliidae 207ff., 270, 370ff., 454
Simulium
 – *costatum* 385ff.
 – *cryophilum* 384, 595
 – *ornatum* 384ff., 595
 – *trifasciatum* 388ff.
 – *vernum* 384, 595
Simulium (*Nevermannia*) *vernum* group 388
Simulium (*Simulium*) *ornatum* group 388
Siphonoperla 258, 270ff., 291ff., 580
 small-scale variability 188
 soluble inorganic element 64
 soluble reactive phosphorus (SRP) 180, 493
 species
 – abundance 503
 – discharge pattern 531
 – distribution 529
 – list 227, 254
 – spatial and temporal variability 534
 specimen 197
 Staphylinidae 312
 statistics
 – community 527
Staurosira elliptica 103f.
 stonefly 256ff., 276ff., 530, 593
 – emergence 276
 stream
 – area 82
 – carbon flow 133
 – shift of emergence 282
 – structural change 253
 stream bed
 – sediment structure 46, 477ff.
 stream bottom 598
 – artificially changed 496
 – sandy stream bottom 484
 stream flow 8ff.
 stream habitat
 – extracellular enzyme activity 168
 – spatial distribution 168
 study sites 10
 substratum dynamics 44

Succinea putris 216
 sulfate 71
 surber samples 197, 394, 459, 592
Surirella brebissonii 103f.
Synedra ulna 117f.

t
Tabellaria flocculosa 117
 temperature 292, 480
 temperature effect
 – episodic 280
 temperature gradient gel electrophoresis (TGGE) 151ff.
 temperature regime 573
 terrestrial by-catch 327
Tetrachaetum elegans 163
 thermal chain 586
 thermal demand 235
 thermal equilibrium hypothesis 476
 thermal regime 577
 threshold emergence temperatures 292
Tinodes 328
 – feeding strategy 419
 – retreat construction 348, 405ff., 444ff.
 – *rostocki* 331, 346ff., 355, 405ff.
 Tipulidae 370
 Trichoptera 327ff., 513, 526ff., 594
 – benthos 348
 – Erlenbach 548
 – life cycle characteristics 532
 – mean annual specimen numbers 518
 – shift of emergence time 576
 trophic relation 256
Tubifex tubifex 212
Tumularia aquatica 163

u
 upstream
 – flights 249, 289, 390
 – migration 246
 – movement 473

v
 valley
 – saturation 44
 vicariance 256
 – temporal 272, 295, 297

w
 water
 – interstitial 88ff., 490
 water beetles 221
 water chemistry 587

water temperature 47ff., 279, 476ff.,
529
– body size 476
– Erlenbach 545
– in the sediment 47
– in the stream 47
wet stream area 46

wind
– systems 27 ff.
wing pads 230, 253, 266, 301f., 307
Wormaldia occipitalis 329, 347, 581

x

Xanthochlorinae 397