

## Index

### a

- acetoxylation 109–111
- alkenes
  - arylation 107–109
  - C–H borylation 112–114
- allyl acetates arylation 108
- amides synthesis 15
  - from alcohols and amines 15–18
  - from esters and amines 18–20
  - polyamides synthesis from diols and diamines 20–24
- ammonia activation 217–218
- appended functionality 71
  - coplanar with pincer chelate
  - pyridine-2,6-dicarboxamide systems 84–85
  - systems incorporating 2,2':6',2''-terpyridine 75–84
- design criteria 72
  - supramolecular architectures 73–75
  - transition-metal catalysis 72–73
- not coplanar to pincer chelate
  - ENE pincer systems 86–87
  - PCP pincer systems 88
  - PEP pincer systems 88–90
  - pyridine-2,6-diimine systems 90–91
- atom-transfer radical addition (ATRA) and polymerization reactions (ATRP) 46–48
- carbon–heteroatom coupling reactions 62–65
- C–C cross-coupling reactions 54–62
- Heck reaction 48–54

### b

- benzyl amine synthesis 103, 105–107
- bond activation by metal–ligand cooperation 3–4

### c

- carbometallated PC(sp<sup>3</sup>) P complexes
  - synthesis and coordination behavior 215–216
- acyclic PC(sp<sup>3</sup>) P complexes coordination flexibility 216–217
- carbon–carbon bond-forming reactions 141–142
  - Kumada–Corriu–Tamao coupling 118–129
  - Mizoroki–Heck reaction 140
  - Negishi coupling 132–139
  - Sonogashira coupling 139
  - Suzuki–Miyaura coupling 129–132
- carbon–carbon coupling reactions 221–223
- carbondioxide
  - activation 225
  - reduction with trialkylsilanes 174
- carbon–heteroatom
  - bond-forming reactions 143–144
  - coupling reactions 62–65
- catalysis, by pincer complexes 1–2
  - amides synthesis 15
  - from alcohols and amines 15–18
  - from esters and amines 18–20
  - polyamides synthesis from diols and diamines 20
  - bond activation by metal–ligand cooperation 3–4
  - esters synthesis 4
  - cross-esters synthesis from primary and secondary alcohols 9–11
  - polyesters synthesis from diols 11, 13–15
  - from primary alcohols 4–9
  - by secondary alcohols acylation 9, 11–13
  - peptides synthesis from  $\beta$ -amino alcohols 24–26

- C–C, C–O, and C–B bond formation 95–96
  - C–H functionalization of organonitriles
    - benzyl amine synthesis 103, 105–107
    - imines allylation 102–105
  - hypervalent iodines reactions 107
    - acetoxylation 109–111
    - alkenes arylation 107–109
    - alkenes C–H borylation 112–114
  - imines and isocyanoacetates reactions
    - chiral pincer complexes application 99–100
    - imidazolines stereoselective synthesis 96–99
    - mechanistic considerations 101–102
- C–C cross-coupling reactions 54–62
- cyclic peptides 24–25
- d**
  - dehydrogenative borylation 237
    - of alkenes and 1,3-dienes 237–239
    - mechanistic considerations 239–241
  - dendritic effect 48
  - density function theory (DFT) studies 159, 167, 172, 259–261, 264, 269, 272
- e**
  - ENE pincer systems 86–87
  - esters synthesis 4
    - cross-esters synthesis from primary and secondary alcohols 9–11
    - from primary alcohols 4–9
    - polyesters synthesis from diols 11, 13–15
    - by secondary alcohols acylation 9, 11–13
- f**
  - frustrated Lewis pairs 77
- g**
  - Grignard reagents 54–61, 118–120, 122–129, 132, 133, 151, 152, 201, 222
- h**
  - Heck reactions 32, 34, 41, 48–54, 253–256
    - computational investigations on thermal feasibility of cycles 262
    - cycles and nanoparticle formation 274–275
    - phenyl bromide oxidative addition and investigations on mechanisms 263–265
    - possible initial reaction steps 263
    - proposed cycles 270–274
    - styrene coordination and chloride dissociation and investigations on mechanisms 265–270
    - intermediates and nanoparticle formation promoted by single system 256–262
  - hydrocarboxylation 231–232
  - of 1,3-dienes 234–235
  - of allenes 232–234
  - hydrogenation and dehydrogenation 223–225
  - hydrogen atom abstraction (HAA) 285, 286
  - hydrogenolysis of late-transition-metal hydroxide and alkoxide complexes 289–294
  - hypervalent iodines reactions 107
    - acetoxylation 109–111
  - alkenes arylation 107–109
  - alkenes C–H borylation 112–114
- i**
  - imidazolines stereoselective synthesis 96–99
  - imines allylation 102–105
  - imines and isocyanoacetates reactions
    - chiral pincer complexes application 99–100
  - imidazolines stereoselective synthesis 96–99
  - mechanistic considerations 101–102
  - isotopic labeling 219
- k**
  - ketones transfer hydrogenation 157
  - Kharasch reaction 46–48
  - Kumada synthesis 54–59, 61
  - Kumada–Corriu–Tamao coupling 118–129
- l**
  - LXL tridentate pincer ligation 150
- m**
  - Mizoroki–Heck reaction 140
  - Mulliken spin density 203
- n**
  - Negishi coupling 54–58, 132–139
  - Negishi cross-coupling reaction 275–277
  - Ni, Pd, and Pt pincer complexes in oxidation states 34
    - higher oxidation states complexes (III, IV) 34–43
    - reduced complexes with pincer ligands 43–46
  - nickel-catalyzed cross-coupling reactions 117
    - carbon–carbon bond-forming reactions 141–142
    - Kumada–Corriu–Tamao coupling 118–129

- Mizoroki–Heck reaction 140
- Negishi coupling 132–139
- Sonogashira coupling 139
- Suzuki–Miyaura coupling 129–132
- carbon–heteroatom bond-forming reactions 143–144

## **o**

- olefins reactions 219–221
- organonitriles C–H functionalization
  - benzyl amine synthesis 103–107
  - imines allylation 102–105

## **p**

- palladium complexes synthesis bearing
  - PXP-pincer ligands (X=Si, Ge, Sn)
  - structural analyses 230–231
  - synthesis 230
- palladium pincer-catalyzed C–C cross coupling reactions 250, 252–253
- complex organic molecules synthesis 249–251
- computational investigations on thermal feasibility of cycles of pincer-catalyzed Heck reactions 262
- cycles and nanoparticle formation 274–275
- phenyl bromide oxidative addition and investigations on mechanisms 263–265
- possible initial reaction steps 263
- proposed cycles 270–274
- styrene coordination and chloride dissociation and investigations on mechanisms 265–270
- Heck reactions 253–256
- intermediates and nanoparticle formation promoted by single system 256–262
- theoretical investigations on
  - pincer-catalyzed Negishi cross-coupling reaction 275–277
- PCP pincer systems 88
- PC(sp<sup>3</sup>) P complexes reactivity and catalytic applications
  - ammonia activation 217–218
  - carbon–carbon coupling reactions 221–223
  - CO<sub>2</sub> activation 225
  - coordinated olefins reactions 219–221
  - hydrogenation and dehydrogenation 223–225
  - isotopic labeling 219
- PEP pincer systems 88–90

- peptides synthesis from  $\beta$ -amino alcohols 24–26
- phenyl bromide oxidative addition and investigations on mechanisms 263–265
- PNP-pincer ligand 118
- polyamides synthesis from diols and diamines 20–24
- polyesters synthesis from diols 11, 13–15
- PSiP chemistry
  - group 8 metal 153–160
  - group 9 metal 161–169
  - group 10 metal 169–179
  - group 11 metal 179
- pyridine-2,6-diimine systems 90–91

## **r**

- redox-active bis(imino)pyridine pincer ligands 189
  - reduced manganese, iron, and cobalt complexes 189–190
  - reduced bis(imino)pyridine cobalt chemistry 200–209
  - reduced bis(imino)pyridine iron chemistry 193–200
  - reduced bis(imino)pyridine manganese chemistry 190–193
- redox process 31–34
  - atom-transfer radical addition (ATRA) and polymerization reactions (ATRP) 46–48
  - carbon–heteroatom coupling reactions 62–65
  - C–C cross-coupling reactions 54–62
  - Heck reaction 48–54
  - Ni, Pd, and Pt pincer complexes in oxidation states 34
  - higher oxidation states complexes (III, IV) 34–43
  - Ni, Pd, and Pt reduced complexes with pincer ligands 43–46
- reductive aldol type reaction 235–237

## **s**

- $\eta^2$ -(Si–H)Pd(0) complex synthesis and reaction as equivalent to PSiP-palladium hydride complexes 241
- reaction with allene 242–243
- reaction with diboron 244–245
- synthesis and structure 241–242
- saturated frameworks and pincer complexes 213
  - carbometallated PC(sp<sup>3</sup>) P complexes synthesis and coordination behavior 215–216

- saturated frameworks and pincer complexes (*contd.*)
    - – acyclic PC(sp<sup>3</sup>) P complexes coordination flexibility 216–217
    - ligands synthesis 213–214
    - PC(sp<sup>3</sup>) P complexes reactivity and catalytic applications
      - – ammonia activation 217–218
      - – carbon–carbon coupling reactions 221–223
      - – CO<sub>2</sub> activation 225
      - – coordinated olefins reactions 219–221
      - – hydrogenation and dehydrogenation 223–225
      - – isotopic labeling 219
  - silyl-based pincer-like bis(phosphino)silyl (PSiP) transition-metal pincer complexes 149–150
    - alternative silyl pincers 180–182
    - group 10 metal PSiP chemistry 169–179
    - group 11 metal PSiP chemistry 179
    - group 8 metal PSiP chemistry 153–160
    - group 9 metal PSiP chemistry 161–169
    - syntheses 151–153
  - Sonogashira coupling 54, 139
  - square-planar d<sup>8</sup> pincer complexes with oxygen and hydrogen 281–284
  - hydrogenolysis of late-transition-metal hydroxide and alkoxide complexes 289–294
  - molecular oxygen insertion into-late-transition-metal hydride bonds 284–289
  - Stille coupling 32, 54, 57
  - styrene coordination and chloride dissociation and investigations on mechanisms 265–270
  - sulfonimines
    - asymmetric benzylation 107
    - catalytic benzylation 106
  - Suzuki coupling 31, 54, 55, 57
  - Suzuki–Miyaura coupling 129–132
- t**
- 2,2':6',2''-terpyridine incorporating systems 75
  - appended hydrogen-bond acceptor/donors 79–84
  - appended Lewis acid/bases 77–79
  - synthetic strategies 75–77
  - transesterification reaction 9
- u**
- unsaturated hydrocarbons catalytic synthetic transformations 229–230
  - $\eta^2$ -(Si–H)Pd(0) complex synthesis and reaction as equivalent to PSiP–palladium hydride complexes 241
  - – reaction with allene 242–243
  - – reaction with diboron 244–245
  - – synthesis and structure 241–242
  - dehydrogenative borylation 237
  - – mechanistic considerations 239–241
  - – of alkenes and 1,3-dienes 237–239
  - hydrocarboxylation 231–232
  - – of 1,3-dienes 234–235
  - – of allenes 232–234
  - palladium complexes synthesis bearing PXP-pincer ligands (X=Si, Ge, Sn)
  - – structural analyses 230–231
  - – synthesis 230
  - reductive aldol type reaction 235–237