

Index

a

- acetophenone 50–53, 293, 344, 348, 443
- acetoxycyclization, of 1,6-enyne 76
- acetylacetone 48
- A₃ coupling reactions 231, 232
- acrylonitriles 69, 211, 212, 310, 348
- activation period 123–125
- active species 123
- acyclic alkane 62
- acyclic aminocarbenes 499
 - ligands 503
 - metalation routes 500
- acyclic aminocarbene species 499
- acyclic carbene chemistry 500, 516–520
- acyclic carbene complexes
 - in Suzuki–Miyaura crosscoupling 505
- acyclic carbene–metal complexes 505
- acyclic carbenes
 - characteristic feature of 503
 - donor abilities 502
 - ligands 502
 - – decomposition routes 504
 - – donor ability 502, 503
 - – metalation routes of 500
 - – structural properties 503
 - stabilized, by lateral enamines 518
- acyclic carbene ligand 519
 - in gold-catalyzed rearrangements 520
- acyclic diaminocarbenes (ADCs) 4, 5, 499
 - Alder-type 510
 - complexes 501
 - ligands, conformational flexibility of 503
- acyclic ketones 69
- ADCs. *See* acyclic diaminocarbenes (ADCs)
- aerosolized water 155
- Ag–NHC species 207
- AgNO₃, bactericidal activity 153
- Albrecht's group 347
- alcohol oxidation 384
- alcohol racemization 356, 357
- alkali metals 398
- alkaline earth metals 398
- N*-alkenyl-substituted *N,S*-HC ligands 349
- 3-alkyl-3-aryloxindoles 58
- alkyl bis(trimethylsilyloxy) methyl silanes 122
 - Tamao–Kumada oxidation of 122
- 2-alkylpyrrolidyl-derived formamidineum precursors 516
- alkyl silyl-fluorides 209
- alkyl-substituted esters 210
- N*-alkyl substituted NHC class 119
- alkynes
 - boration of 225
 - borocarboxylation 233, 234
 - hydrocarboxylation 234, 235
 - metal-catalyzed hydrosilylation of 132
 - semihydrogenation 232, 233
- allenes 77
 - synthesis, mechanisms 203
- 3-allyl-3-aryl oxindoles 60
- allylbenzene 345
 - cross-metathesis (CM) reactions of 510
- allylic alkylations 509
- allylic benzimidate
 - aza-Claisen rearrangement of 514
- allylic substitution 220
 - reactions 361–363
 - by *in situ* generated Ru–NHC complexes 362
- 3-allyl oxindole 60
- aluminum reagents
 - conjugated addition using 209
- amide formation 355
 - proposed mechanism 354
- amidine ligand 504
- amidinium-derived chiral ADC ligands
 - enantioselective catalysis with 517

- amidinium precursors
 - chiral ADC ligands derived from 516
 - amido NHC complex 400
 - amino acid complexes 184
 - aminocarbenes 4
 - ampicilin 175
 - anisotropies 4
 - antibacterial activity 154
 - antibacterial agents, cationic complexes
 - study 174
 - anticancer NHC–Pd Complexes 106
 - antifungal agents, cationic complexes study 174
 - antiselective substitution, mechanism 204
 - antitumor reagents
 - copper complexes 195, 196
 - metals 178–195
 - – auranofin/cisplatin, structure of 178
 - – gold–NHC systems 178
 - apoptotic pathway 181
 - Arrhenius law 130
 - arylaldehydes 49
 - aryl alkyl ketones 54
 - aryl/alkyl-substituted (NHC)Pt(dvtms)
 - complexes 134
 - arylamine alkylation 353
 - arylation 357–359
 - C–H activation/deprotonation pathway 358
 - DFT calculations 357
 - of 2-phenylpyridine using 357
 - arylboronic acids 49
 - aryl bromides 95, 96
 - aryl chlorides 90
 - β -arylcycloalkanones 66
 - aryl ether reduction 384
 - 3-aryl-3-fluoro-oxindole 60
 - aryl iodide, double carbonylation 214
 - aryl magnesium bromides 68
 - 3-aryl-3-methyloxindole 59
 - Aspergillus niger* 151
 - asymmetric allylic substitution 219
 - asymmetric catalysis 97–100
 - chiral bis-NHC ligand precursor and bis-NHC–Pd complex, synthesis 98
 - molecular design of iminoalkyl imidazolium ligand 98
 - asymmetric catalysis using NHC–Pd 63–65
 - asymmetric allylic alkylation using NHC–Pd 65
 - asymmetric intermolecular α -arylation
 - – using NHC as chiral modifiers of nanoparticles 63
 - asymmetric intramolecular oxidative amination 65
 - superparamagnetic Fe₃O₄-supported NHC-based catalyst
 - – for enantioselective allylation 64
 - asymmetric conjugated addition 207
 - asymmetric cross-metathesis (ACM)
 - processes 42
 - asymmetric diamination 62
 - of alkenes 63
 - asymmetric hydrogenation 44
 - aromatic/heteroaromatic compounds 45
 - asymmetric olefin metathesis 40
 - asymmetric ring closing metathesis (ARCM) 40, 41, 335
 - using C₁-symmetric NHCs 43
 - asymmetric ringopening cross-metathesis (AROCM) 40, 42, 335
 - *E*-selectivity in 44
 - *E/Z* selectivity 44
 - metathesis 44
 - Au–ADC catalysts 512
 - [Au(Br)(NHC)] 191
 - [Au(Cl)(IMes)] 192
 - [Au(Cl)(IPr)] 188, 189
 - [Au(Cl)(NHC)] species 176
 - Au^I-bound binaphthyl isocyanides 515
 - Au(III) bis-NHC complexes 194
 - [Au(NHC)(PPh₃)]PF₆ complexes 192
 - [Au(OH)(IPr)], 188, 189
 - auranofin structure 178
 - azabicycles 48
 - aziridination 221
 - azolium salt 408
 - deprotonation 408
 - azomethine 247
- b**
- Balz–Schiemann reactions 230
 - Be–IPr complex 435
 - bent allene resonance 519
 - benzimidazole-derived carbenes 94
 - benzimidazoles 273
 - benzimidazolin-2-ylidenes 94
 - ligands 94
 - benzimidazolium ligands 96
 - benzofurans 46, 349
 - benzophenone 345
 - benzothiazolium salts 346
 - benzothiophenes 47, 349
 - benzyl alcohol 353
 - benzylamine 354
 - N*-benzylideneaniline 343
 - N*-benzylidenebenzylamine 354
 - N*-benzyl 2-phenylacetamide 354

- benzylpropargyl ether 140, 143
beryllium–NHC complex 434
– ionic 434
bidentate (NHC)–Pd complexes 92, 94
bidentate trans-chelating NHC-ligands 95
bifunctional catalysis 280
binaphthyl-2-2'-diamine (BINAM) 86
BIneoPent compound 119
BIN ligand 344
bioxazoline-derived NHC complex 27
bioxazolines 56
bis-amido NHC ligands 408
bis(benzimidazolyldiene) zirconium complex 408
bis(bis(trimethylsilyl)amino) magnesium(II) complex 437
bis-NHC CNC-type pincer ligands 410
bis-NHC complexes, macrocyclic 166
bis-NHC–copper complexes 201, 214
bis(NHC)–lithium complex 432
bis(NHC)–Pd complexes 94
bis-phenoxide–NHC ligand 405
trans bis-silyl ethylene species 140
bis-silyl platinum carbene species 144
– synthesis of 144
bis(trimethylsilyloxy)methylsilane 122, 125, 144
BMOL (bipotent murine oval liver) 181
[BnN(CH₂CH₂CH₂RIm)₂]₂PdCl₂ 96
N-Boc-7-azanorbornene 49
bond dissociation energy 26
“boomerang” effect 311
borocarboxylation
– of alkyne 233, 234
– proposed mechanism 234
boron 479–481
boronic acids 48–50
boronium salt 480
boron reagents
– conjugated addition 210
borrowing hydrogen 351–356
– reaction with C–H activated complex 351
boryl–copper complex [Cu(Bpin)(SImes)] 234
borylene 480
β-boryl ketones 71
B₂pin₂, allylic substitution 220
breast cancer line MB157
– *in vivo* xenograft model 164
N-(2-bromoaryl)-*N*-alkyl-2-arylpropanamides 57
4-bromo-2,5-bis(hexyloxy)phenyl)magnesium chloride 105
Buchwald–Hartwig aminations 88, 89, 509
Burkholderia cepacia complex 154, 156
c
CAACs. *See* cyclic alkyl amino carbenes (CAACs)
cancer cell lines 193
– IC₅₀, comparison of 185, 187, 190, 191, 196
carbene 1,3-dimethyl-4,5-dimethylimidazol-2-ylidene 398
carbenes 1, 69
– abnormal 5, 6
– derived from imidazolium salts 19
– dihydrocarbene 5
– electron distribution mapping 4
– isodesmic calculations 5
– mesoionic 273
– singlet 2, 7
– soft 282
– stable 1, 3, 5
– triplet 1, 2
carbenoids 458
carboboration reactions 226
– mechanism for 227
carbodicarbene 519
carbon–nitrogen bond-forming reactions 382
carbon–sulfur bond-forming reactions 382, 383
carbonyl compounds, boration 223
carbonyl functions, transformation of 226
carbonyls, olefination
– [Cu(Cl)(NHC)], using 227
carboxylation mechanism 214
caspase-3, 181
catalyst activation 143, 144
catalyst deactivation pathways 125–127
– (IPr)Pt(AE) by benzylpropargyl ether 144
catalyst screening 134–137
catalysts derived from nickel(0) and nickel(II) sources 373
catalytic asymmetric allylic alkylation (AAA)
– with vinylaluminum reagents 219
catalytic efficiency 31
cationic complexes studied
– as antibacterial/antifungal agents 174
cationic gold(I/III) complexes 177
cationic NHC–copper complexes catalyzed cycloaddition reaction 216
cationic transfer hydrogenation systems 344
chalcones 71
Chalk–Harrod mechanism 113, 123, 145
Chalk mechanism 113, 123
charge distribution 4

- Chauvin metallacyclobutane 307
- chelated bis(acyclic diaminocarbene) complex 499
- chelated diallyl ether 144
- chelates 93
- chiral acyclic carbene ligands 513
- chiral Au^I-ADC complexes.
 - enantioselective alkynylbenzaldehyde cyclization/acetalization catalyzed by 515, 516
- chiral bidentate NHC-bearing ruthenium complexes 334
- chiral bidentate phosphine–oxazoline ligands 39
- chiral 6,6'-dimethoxybiphenyl-2,2'-diamine 76
- chiral imidazolium tetrafluoroborates 57
- chiral NHC–CuCl catalyst 68
- chiral NHC–palladium complexes 91
- chiral NHC–Ru complex 40
- chiral palladium bis(ADC) catalysts
 - enantioselective aza-Claisen rearrangement with 514
- chiral Ru complexes, bearing
 - binaphtholate and biphenolate substituted NHCs 335
 - 4,5-di-tert-butyl-substituted unsymmetrical NHCs 333
 - monosubstituted backbone bearing NHCs 334
- chiral Ru metathesis catalysts 332
- chiral ruthenium catalysts 42
- 2-chloroamidinium 501
- chlorobenzene 104
- chloroiminium ions 501
- chromium (0) carbonyl 398
- chromium (II) complexes 414
- chromium (III) complexes 414
- cinammyl-type substrates 68, 69
- ciprofloxacin 158, 176
- cis*-arrangement 348
- cisplatin 164, 196
 - IC₅₀ (μM) measured for 195
 - structure 178
- Cl–Ti–Cl angle 403
- CO detector 288
- colloidal platinum species 113
- CoLo 320 DM 181
- 1,4-conjugated borylation 230
- conjugation 4
- copolymerization
 - ethylene/1-hexene 399
 - 1-octene 399
- copper–bifluoride complex 230
- copper hydride species 200
- copper-mediated carboboration 226
- copper–NHC complexes 211
 - hydrothiolation, hydroalkoxylation and hydroamination catalyzed 211
- Cr NHC complexes
 - in combination with MAO 414
 - indenyl-functionalized NHC complexes of 414
 - Schulz–Flory distribution 414
 - supported by a chelating bis-(*N*-heterocyclic carbene) ligand 414
- cross-coupling reactions 85–93
 - Buchwald–Hartwig aminations 88, 89
 - Heck reaction 92, 93
 - Hiyama coupling 89, 90
 - Kumada coupling 90
 - Negishi reactions 89
 - Sonogashira coupling 90–92
 - Suzuki–Miyaura Coupling 85–88
- C–S/C–N bonds, formation 210
- ¹³C shielding tensor 4
- C–Si bonds formation, from addition of halosilane to Sc–NHC complexes 399
- Cu-catalyzed allylic alkylation
 - using *in situ* generated ADC ligand 509
- [Cu(Cl)(IMes)], imidazole derivatives 220
- [Cu(Cl)(IPr)] 200
- [Cu(Cl)(NHC)]
 - NHC–copper systems of 223
- olefination of carbonyls 227
- [Cu(IPr)][HF₂] complex 231
- [Cu(O^tBu)(IPr)], σ-bond metathesis 206
- Cu(OTf)₂, 206
- [Cu(R)(IPr)] complexes
 - formation of 213
- cyclic alkyl amino carbenes (CAACs) 428, 478, 479
 - boron 479
 - borylene 480
 - CAAC-stabilized borylene 480
 - coordination with 1-bromo-2,3,4,5-tetraphenylborole 481
 - pinacolborane 479
 - stable borylene, boronium salt, and radical cation, formation 480
 - carbon 481, 482
 - [4+1] cycloaddition product with bis(cyclooctatriene)iron(0) 482
 - stable amino ketene from reaction of CAAC B with 481
 - ligands 246, 468, 479

- nitrogen 483
- activation of ammonia using CAACs resulting in oxidative addition 483
- phosphorus 483
- CAAC/NHC-stabilized phosphorus mononitride 486
- formation of a phosphinyl radical cation 486
- P-H bond activation 484
- ^{31}P NMR chemical shifts of carbene–phosphinidine adducts 487
- reactivity of CAAC with white phosphorus and [4+2] cycloadditions 485
- structure of CAAC adduct of 2,3,4,5-tetraphosphatriene 484
- white phosphorus activation and fragmentation 485
- silicon 482
- activation of Si-H bonds 482
- cyclic carbenes stabilized, by lateral enamines 518
- cyclic π -delocalization 4
- cyclic RGD 167
- [2+2+2] cycloaddition 386
- carbocycles from aryne intermediates 388
- diynes and aldehydes, 387
- diynes and carbon dioxide 386
- enynes and aldehydes/ketones 387
- heterocycles from 387
- yielding pyridines 388
- yielding substituted pyridones 388
- [3+2] cycloaddition 388
- enoate/alkyne cycloaddition 389
- enone/cyclopropane cycloadditions 389
- [4+2+2] cycloaddition 389, 390
- yielding eight-membered polycycles 389
- 2-cycloheptanone 66
- cyclohexylallene 55
- cyclometallated-NHC complexes 280
- cyclopalladate ferrocenylimine NHC–Pd complex 95
- cyclophane moieties 180
- cyclopropanation 221
- cyclopropenones 458
- cytotoxicity, neutral/cationic complexes 188
- d**
- deactivation pathways 143, 144
- (IPr)Pt(AE) by benzylpropargyl ether 144
- dehalogenation, of aryl halides 383
- dehydrogenative silylation 350
- density functional theory (DFT) 1, 203, 342
- calculations 349
- studies of a model carbene 3
- deoxygenation, of aryl ethers 385
- deprotonation, of imidazolium ions 428
- desymmetrization, meso-trienes 43
- 1,3-dialkyl-4,5-dimethylimidazol-2-ylidene 402
- N,N'*-dialkyl-substituted NHC–Ru complexes 314, 315
- N,N'*-dialkyl substituted *N*-heterocyclic carbene complexes 313, 314
- N,N'*-diamidocarbenes 458, 474
- N,N'*-diamidoketenimines 458
- diamination, alkenes asymmetric 63
- (1*R*,2*R*)-1,2-diaminocyclohexane 40
- 1,4-diazabicyclo[2.2.2]octane (DABCO) 353
- diazo compounds 221
- dibenzoylmethane (DBM) 222
- dicarbene Pd(II) complex 97
- dicarbonyl complex 288
- 1,2-dichloroethane 97
- 4,5-dichloroimidazole 157
- dichlorosilamines 461
- dienoates 69
- dienones 69, 70, 387
- 1,2-dihapto ligand 284
- 2,3-dihydrobenzofuran 46
- dihydrobenzothiophenes 349
- 2,6-diisopropylphenyl 2,4,6-triisopropylphenyl group 97
- 1,3-diisopropyl-3,4,5,6-tetrahydropyrimid-2-ylidene 433
- dimer, initiation 126
- dimerize thermodynamic propensity 504
- 2,2-dimethyl-1-arylpropane-1-amine 61
- dimethylphenylsilane 140
- dinuclear CCC pincer Ir(III) complexes 282
- dinuclear gold(I) complexes
- cyclophane and noncyclophane moieties 179
- diones 458
- diphenylberyllium 434
- diphenylberyllium–NHC complex 435
- (1*R*,2*R*)-diphenylethyldiamine 40
- N,N*-diphenyl-substituted NHC bearing complexes 327
- 1,3-dipolar cycloaddition 215, 247
- discrete Ni(II)–NHC catalysts 374–376
- discrete Ni(I)–NHC catalysts 374
- discrete Ni(0)–NHC catalysts 373
- 1,4-disubstituted-1,2,3-triazole 215
- 5,7-di-tert-butyl-3-phenyl benzoxazol-4-ylidene 273
- ditz ligand 287

- 1,3-divinyltetramethyldisiloxane (dvtms) 111
 – with hexachloroplatinic acid 111

e

- electron-deficient vinylarenes 212
 π electron density 4
 electron density mapping 4
 electronic characteristics 25
 electronic effects 25
 electronic properties, of NHC ligand 31, 32
 electron-withdrawing groups (EWG) 212
 – aryl derivatives 226
 enantioselective hydroboration 224
 enhanced permeability and retention (EPR)
 effect 167
 enoates 205, 389
 – 1,4-reduction of 205
 enones
 – addition of dialkylzinc reagents to 66
 – 1,4-conjugate addition of diethylzinc to 65
 – cyclic 48, 209, 294
 – 1,4-reduction of 205
 enthalpy 7
 1,3-enyne, hydrosilylation of 140
 enzyme thioredoxin reductase 182
 epoxidation
 – of cyclooctene catalysed by 416, 417
 – olefin 415
 – terminal alkenes with PhI (OAc)₂, 363
 esters 69–71
 ethylbenzene 347
 ethyl diazoacetate (EDA) 364
 ethyl moiety 186
 ethyl propiolate 97
 ethyl *trans*- β -methylcinnamate 205

f

- five-member ring NHC-bearing complexes 93
 fluconazole 176
 flucytosine, antibacterial/antifungal agents 175
 fluorenyl-NHC scandium complex 399
 fluorinated alkenes 310
o-fluorinated aryl substituted NHC
 complexes 318
 Friedel–Crafts/cyclization 76, 77
 Friedel–Crafts cycloisomerization
 – of arenyne 455
 Fujiwara hydroarylation, of acetylenes 97
 functional group tolerance 120–122, 139, 140, 142
 – and substrate scope 139–142

- functional magnetic nanoparticles 63
 fungicidal activity 153

g

- geometric structure 4
 glass transition temperatures 104
 gold antimicrobial agents 173
 gold catalysis 510, 511
 – alkyne functionalizations 512, 513
 – allene/alkene hydrofunctionalizations 512
 – enantioselective catalysis, with chiral acyclic
 carbenes 513–516
 – enyne cyclizations 511, 512
 gold complexes
 – IC₅₀, comparison of 185
 – series of 184
 gold(I)–ADC catalyst
 – methoxycyclization of an enyne 512
 gold(I)-catalyzed enyne cyclization
 – ligand effects on selectivity 511
 gold(I)–ferrocenyl complex 182
 gold–NHC systems 178, 183
 – antimicrobial activity of 177
 – chloride and bromide species 183
 – series of cationic complexes 180
 Grignard reagents 66, 67, 208, 218, 437
 – allylic substitution 219
 – conjugated addition 208
 group 4 N-heterocyclic carbenes complexes
 – molecular structure 407
 – NHC-incorporating chelating ligands
 with 405
 – obtained by THF substitution 402
 – ring-opening polymerisation, of lactide 406
 Grubbs olefin metathesis catalysts 39
 Grubbs second-generation catalyst 309

h

- hafnium derivatives 408
 Halex reactions 230
 4'-haloarylketones 105
 1-(4-halophenyl)ethanol 359
 Harrod mechanism 113, 123
 Heck coupling reactions 508, 509
 Heck olefination 95
 Heck reaction 92, 93, 381
 heteroatom containing NHC-bearing
 complexes 329
 N-heterocyclic carbene–lithium complex 432
 – homoleptic 432
 N-heterocyclic carbene–metal complexes 235
 N-heterocyclic carbene platinum
 complexes 138

- N*-heterocyclic carbenes (NHCs) 1, 112, 211, 499
 - bonding to metal centers 8–12
 - carbenic and ylidic resonance forms 4
 - in context of other stable carbenes 16–18
 - coordination chemistry 398
 - electron-donating ability 18
 - properties 1
 - quantifying properties 13
 - electronic properties 14–16
 - steric impact 13, 14
 - salts and adducts of 20–22
 - species with reduced heteroatom stabilization 6
 - stability 6–8
 - bonding arrangements 7
 - structure 1
 - synthesis 19, 20
- N,S*-heterocyclic carbenes 346
- heterodimetallic complexes 290
- heteroleptic bis-NHC–copper complexes 215
- hexachloroplatinic acid 111
- 1,1,1,3,3,3-hexamethyldisilazide (HMDS) 437
- 1-hexene 347
 - reduction 348
- histidine-derived NHC analog 345
- Hiyama coupling 89, 90
- Hiyama–Denmark cross-coupling reaction 142
- hole-transporting materials (HTMs) 104
- homo-dicarbene Pd(II) complexes 94
- homoleptic yttrium complexes 401
- homopolymerization
 - of functionalized norbornene monomers 104
 - for phenylene- and thiophene-based monomers 105
- H₂O-soluble silver–NHC
 - synthetic route 154
- Hoveyda–Grubbs catalyst 355
- Huisgen reaction 215
- hybrid NHC ligands 280
- hydroalkoxylation 212
- hydroboration 112, 224
 - and diboration catalysts 112
- hydrocarboxylation 235, 236
- hydrogenation
 - of both pinacolone and styrene oxide 346
 - of esters by amino-functionalized NHC complex 350
 - proposed mechanism 350
- hydrogen-bond supported heterocyclic carbenes (HBHCs) 511
- hydrophobic silver–NHC complexes
 - for anticancer activity 165
- hydrosilylation 111, 349, 385
 - alkenes 111–114
 - activation period 123–125
 - catalyst deactivation pathways 125–127
 - functional group tolerance/substrate scope 120–122
 - general scheme 112
 - *N*-heterocyclic carbene platinum(0) complexes 134
 - Karstedt's catalyst 113
 - NHC platinum(0) complexes 111–114
 - phosphine ligands 114, 115
 - quantitative kinetic modeling 129–133
 - regioselectivity of 136
 - semiquantitative kinetic studies 127–129
- alkynes 111, 112, 133, 134, 385
 - catalyst activation 143, 144
 - catalyzed by (ICy)Pt(dvtms) 121
 - deactivation pathways 143, 144
 - functional group tolerance/substrate scope 139–142
 - general scheme 112
 - proposed mechanism 145, 146
 - qualitative kinetic studies 142, 143
 - regioselectivity
 - catalyst screening and NHCs impact 134–137
 - influence of silane 137, 138
 - second-generation catalyst 138, 139
- benchmark reaction for catalysts 113
- carbonyls 385, 386
- diene 124
- 1,3-enyne during total synthesis of lactimidomycin 140
- of internal alkynes catalyzed by complex (IPr)Pt(AE) 140
- 1-octene 112
- silane conversion curves 115, 119, 120
- of phenylacetylene by triethylsilane 136
- platinum(0) catalysts for 112
- proposed mechanism for alkynes catalyzed by (IPr)Pt(AE) 145
- using cationic copper(I) complexes 201
- hydrosoluble imidazolium-chelated palladium catalyst 103
- hydrothiolation 212
 - mechanism for 212
- hydrotris(3-ethylimidazolium)borate tetrafluoroborate
 - deprotonation 432

i

IAd synthesis 3
 IBiox7 HOTf
 – allene, formation 202
 ICy (1,3-dicyclohexylimidazol-2-ylidene) 200
 (ICy)Pt(dvtms) complex 127
 – alkenes catalyzed, hydrosilylation 131
 – functionalized alkenes catalyzed, hydrosilylation of 121
 – rate constants 132
 – treatment of 125
 (ICy)Pt(η^2 -diene) derivatives 124, 125
 – chelating moiety of 124
 – modification of the chelating moiety 124
 (ICy)Pt(1-octene)₂, 132
 IMes (1,3-bis(2,4,6-trimethylphenyl)imidazol-2-ylidene) 203
 IMes chloride salt 175
 IMes ligand 235
 (IMes)Pt(dvtms)
 – polymerizations by hydrosilylation 122
 imidazol- and imidazolin-2-ylidene ligands
 – modifications in 313
 imidazole-based silver–NHC complexes 156
 imidazoles 94, 273
 – ring 52, 186, 209
 imidazole-2-ylidene congeners 86
 imidazolidin-2-ylidenes 4
 imidazolium diphosphine ([PCP]H)PF₆ salt 93
 imidazoli(ni)um salt 216
 imidazolin-2-ylidenes 94
 imidazolium-based NHCs 4
 imidazolium chloride 3
 imidazolium cyclophane gem-diol dichloride 153
 imidazolium ring 3
 imidazolium salts 58, 169
 1,2-imidazol-3-ylidenes 6
 1,2-imidazol-4-ylidenes 6
 imidazol-2-ylidenes 4
 imido Ti complex 410
 imine reduction 383, 384
 incubation test 188
 indenylidene-based ruthenium metathesis catalysts 311
 indenylidene complexes 311
 indium–NHC chemistry 456
 indole addition
 – Au^I-ADC vs. Au^I-NHC catalysts 512
 inverse addition protocol 121
in vivo xenograft model 164
 iodines 95

4-iodoacetophenone 344
 IPent–Pd–PEPPSI complex 85
 iPr₂-bimyl ligands 194
 – IC₅₀ (μM) measured for 195
 IPr ligand 214
 IPr–Li(THF) adduct, chain structure 431
 IPr–Mg(*n*-Bu)(NR₂) adducts 437
 – structure 438
 [(IPr)PdCl₂]₂(μ-pyrazine) complex 90
 IPr–Pd–PEPPSI complex 88, 104
 – synthesis of π -conjugated polymers 105
 (IPr)Pt(AE) complex 140
 – alkynes catalyzed, hydrosilylation of 141, 145
 – deactivation of 144
 – 1-octyne, hydrosilylation of 142
 – synthesis of 138
 (IPr)Pt(dvtms) 137
 – catalyst 138
 – platinum derivative 138
 (IpTol)Pt(dvtms) 136
 – crystal structure of 135
 [IrCl(CO)₂(L)] complexes 5
 iridium pincer complexes, normal/
 abnormal 282
 Ir(III)–Ru(II) heterodimetallic complex 287
 Ir(I)/Rh(III) heterodimetallic complex 284
 isocyanide-derived Pd-bis(ADC)
 complexes 514
 isocyanide–Pd^{II} synthons
 – ADC ligands 502
 isocyanides
 – amines, metal-promoted addition of 501
 – chiral ADC ligands derived from 514–516
 – complexes containing chiral bis(ADC)
 ligands 514
 isodesmic calculations 4
 isomerization 390
 – of C=C bonds 360, 361
 – products 345
 – Ru–NHC catalyzed isomerization of allylic
 alcohols 361
 – by *in situ* generated [Ru(SIMes)(PCy₃)(CO)
 HCl] 360
 isopropoxy benzyldiene 314
 isopropoxystyrene 310

j

Janus-Head-type NHC ligands 287, 352
 Jurkat leukemia 181

k

Karstedt's catalysts 122, 133
 Karstedt's complex 111–114, 121

- structure of 112
- ketones 72, 73
 - diboration of 223
 - hydrosilylation of 201
 - Rh^I-catalyzed enantioselective hydrosilylations of 516
- Kinetic assays
 - synthesis of 115–120
- Kirby–Bauer disk-diffusion method 152, 157
- Kumada–Corriu coupling reaction 376–378
- Kumada coupling 90

I

- lactimidomycin, synthesis
 - 1,3-enyne, hydrosilylation of 140
- lactones 69
- lanthanides 398
- Lewis acids 121, 278, 427, 438, 453, 479
- Lewis bases 397, 438, 440, 459, 481, 518
- ligand angle profile 28
- Li[N₂ with incorporated LiCl 432
- Lindlar's catalyst 232
- Li–NHC complexes 431
- [LiOCR1R2CH2(1-C{NCHCHNR3})], 432
 - with incorporated diethyl ether and lithium iodide 432
- lipophilicity 180, 181
- Li₄ tetrahedron, stabilized by alkynyl and NHC ligands 430
- lithium alkoxide NHCs 431
- lithium–cyclopentadienide NHC complex 430
- lithium–NHC complexes 433
- lithium *tert*-butoxide 204
- liver cell lines 181
- [LNi(CO)₃] system 273
- L-tyrosine phosphate (LTP) 160
 - molecular structure of 161

m

- macrocyclization
 - through intramolecular reductive coupling 392
- MCF-7 (breast cancer) 181
- mesityl *N*-substituents 3
- meso*-chloro-substituted tricarboindocyanine dyes
 - Suzuki arylations of 507
- mesoionic bis-NHCs 284
- mesoionic carbenes (MICs) 273
- metal-bound acyclic oxyalkylcarbenes 499
- metal–carbene complexes 2
- metal complexes

- group 3-7, 398–421
- metal–laaziridine 413
- metal–ligand bond 113
- metal–thiolate bonds 193
- metathesis
 - asymmetric reactions catalyzed by chiral Ru complexes 331
- catalysts
 - bearing NHC ligands 309
 - indenylidene-based ruthenium metathesis catalysts 311
 - phosphine-free 310
 - second-generation 309
 - structures 308
- complexes bearing fluorinated *N*-aryl-substituted NHCs 323
- diastereoselective ring rearrangement 326
- influence of
 - chiral bidentate *N*-heterocyclic carbenes 334, 335
 - chiral monodentate *N*-heterocyclic carbenes 330–334
 - four-, six-, and seven-membered *N*-heterocyclic carbenes 327, 328
 - heteroatom containing *N*-heterocyclic carbenes 328, 329
 - *N*-heterocyclic carbene bearing chiral Ru complexes 330
 - NHCs for metathesis in water and protic solvents 335, 336
 - 4,5-substituted *N*-heterocyclic carbenes 325–327
 - symmetrical 1,3-substituted *N*-heterocyclic carbene in 313–317
 - unsymmetrically *N,N*x00301;-substituted *N*-heterocyclic carbenes 319–325
- simplified mechanism for ruthenium-based metathesis. 308
- methanol 70
- methicillin-resistant *Staphylococcus aureus* (MRSA) 151
- 4-methoxyacetophenone 346
- p*-methoxyphenyl-substituted bis(NHC)–Pd complex 97
- methyl 2-acetamidoacrylate 54
- 3-methyl-3-aryl oxindoles 58
- N*-methylbenzylamine 97
- methylberylliumhydride–IPr dimer 435
- N*-methyl-4,5-diarylimidazolium salts 189
- 1-methylimidazole 89
- 1-(4-methylphenyl) ethanol 346
- β-methylstyrene 346
- N*-methyl substituted analog 183

- Michael acceptors 206
- microdilution methods 154
- minimum bactericidal concentration (MBC) 155
- minimum inhibitory concentration (MIC) 152, 159, 174, 175
- mitochondrial membrane permeability (MMP) 178
- Mizoroki–Heck reaction 94, 95
- Mn–NHC adduct $(\eta^2\text{-C}_5\text{H}_5)(\eta^4\text{-C}_5\text{H}_5)\text{Mn}(\text{IMes})$ complex 418
- $[\text{Mo}(\eta^3\text{-C}_4\text{H}_7)(\text{bipy})(\text{CO})_2(\text{N-RIm})]^+(\text{OTf})^-$ complexes 417
 - deprotonation 417
 - reactivity of molybdenum (II) complex 417
- Mo(II)–NHC compounds 416
 - epoxidation of cyclooctene catalysed by 416
 - molecular structure 416
- mono-I(OR)₂ complexes 348
- mononuclear systems 180
- monosubstituted alkenyl amines 409
- morphine 49
- Morris's amino-functionalized NHC complex 349
- multiple carbon–carbon bonds
 - functionalization of 223
- n**
- N*-naphthyl-substituted complexes 317
- natural bond order (NBO) calculations 4
- NCI-H460 (lung carcinoma) 185
- Negishi coupling reaction 89, 381
- neutral copper(I) complex 215
 - propargylamines, formation of 232
- neutral NHC–gold(I) complexes 190
- neutral NHC gold(I/III) complexes 187
- NHC–Ag 75
 - enantioselective diboration of styrenes 75
 - silver carbene complexes
 - act as efficient NHC transfer reagents for 75
 - styrene catalyzed by silver complex, enantioselective hydroboration of 75
- NHC–Au 75
 - asymmetric alkoxy cyclization of enynes 77
 - asymmetric Friedel–Crafts/cyclization of nitrogen-tethered 1,6-enynes 77
 - asymmetric hydrogenation of olefins 78
 - cycloisomerizations of 1,6-enynes 76
 - enantioselective cycloaddition 79
 - enantioselective cycloisomerizations 76
 - asymmetric alkoxy cyclization of enynes 78
 - enantioselective desymmetrization of diynesulfonamide 78
 - 1,6-enynes 76
 - intramolecular hydroamination of allenes 77
 - nitrogen-tethered 1,6-enynes 77
 - enantioselective desymmetrization of diynesulfonamide 77
 - enantioselective hydrogenation 78, 79
 - olefins 78
 - enantioselective intramolecular hydroamination of allenes 77
- NHC–Au(I) catalysts, applications of 253
 - alteration of reactivity of switching from phosphine to NHC ligands 258–263
 - gold-catalyzed additions to 1,6-enynes 259
 - gold-catalyzed allene diene cycloaddition 262
 - gold-catalyzed carbocyclization 260
 - gold-catalyzed cross-coupling reaction of enamines and alkynes 263
 - gold-catalyzed diyne polycyclization 262
 - gold-catalyzed enallene cycloisomerization 261
 - gold-catalyzed enyne cycloisomerization 263
 - enantioselective transformations based on chiral, enantiopure NHC-based catalysts 264–266
 - axially chiral gold complex-catalyzed enantioselective intermolecular [4+2] cycloaddition of allenamides and dienes 266
 - cyclization of alkynylbenzaldehydes catalyzed by acyclic diaminocarbene (NAC) gold complexes 266
 - enantioselective alkoxy cyclization catalyzed by chiral gold complexes 264
 - transformation of propargyl esters 265
 - improvement of catalyst
 - due to tuning the steric properties of the NHC ligands used 256, 257
 - gold-catalyzed hydrofluorination 258
 - gold-catalyzed intramolecular hydroamination 255
 - gold-catalyzed intramolecular rearrangement of sulfinyl alkynes 256
 - gold-catalyzed oxidative rearrangement 256
 - gold-catalyzed rearrangement of allyl acetates 257

- intermolecular hydroamination of allenes and alkynes 257
- optimized conditions for Echavarren's intermolecular applications of phenol synthesis 258
- stability due to use of NHC ligands 253–256
- by tuning electronic properties of NHC ligands used 257, 258
- NHC–Au(I) chlorides 243
 - activation of 248–253
 - isolation of an intermediate 249
 - by silver salts 249
 - commonly used NHC precursors 245
 - displacement of a thioether ligand from 244
 - gold-catalyzed formation of dibenzopentalenes 254
 - isonitrile complex, reacted with a secondary amine 247
 - proposed mechanism 252
 - scope of synthesis 248
 - synthesis
 - of Au(I)–alkyne complexes 255
 - [Au(IPr)(OH)] 250, 251
 - of CAACAuCl 246
 - of chiral, enantiopure, C₂-symmetric NHC–Au(I)Cl complexes 246
 - of dual-activation catalysts (DACs) based on IPr 255
 - of isolable, easy-to-handle and catalytically active complex 250
 - routes 244
- NHC–Au(I) complexes 243
 - mediated hydration of alkynes 243
- NHC-based Ru (pre)-catalysts 28
 - *ab initio* molecular dynamics 28
- NHC-borate, complexed to Li(THF)₂, 431
- NHC chromium (II) metallocene complexes 414
- NHC complexes of group 1 elements 429
 - lithium 429–432
 - potassium 433, 434
 - sodium 432, 433
- NHC complexes of group 2 elements 434
 - beryllium 434, 435
 - calcium, strontium, and barium 437, 438
 - magnesium 436, 437
- NHC complexes of group 13 elements 438
 - aluminum 452–454
 - monomeric aluminum(I) complex, hydrogen transfer reaction 454
 - NHC-transfer product, from reaction of IMes–AlCl₂H and an α -diimine 452
 - reaction of *It*Bu–AlMe₃ to generate 453
 - boron 438
 - chemistry of NHCs and charged boron compounds 446–448
 - chemistry of NHCs with boranes 439–441
 - frustrated Lewis pairs 444–446
 - NHC–boranes as hydrogen sources 441–444
 - NHC chemistry of other boron compounds 448–451
 - gallium 454–456
 - carbene-stabilized digallane complex 455
 - dibromogallane–NHC adducts 454
 - IMes adducts of tetrahalogallane 455
 - NHC-stabilized gallium hydride complexes 454
 - stable Ga6 octahedron with NHC ligands 455
 - indium and thallium 456
 - indium trihydride–NHC complexes 456
 - Me₂InCl–NHC complex 456
 - thallium trichloride–NHC complexes 456
 - triscarbene thallium(I) complex 456
- NHC complexes of group 14 elements
 - carbon 456–458
 - allenes as acceptor molecules for NHCs 457
 - bent allene, bis(*N,N*-dimethylbenzimidazolyl)allene 457
 - IPr–C60 adduct 458
 - Lewis pair, to cleave disulfide bonds 457, 458
 - NHC–Si(IV) adduct prepared 459
 - germanium 464–466
 - chloroarylgermylene 465
 - NHC–GeH₂–BH₃ adduct 464
 - NHC–Ge(*O**t*Bu)₂ and NHC–Ge(NCS)₂ 465
 - NHC–germylene 464
 - NHC-stabilized GeCl₂ complex 465
 - NHC-stabilized germanone 466
 - reactivity of NHC–GeX₂ adducts 465
 - silicon 459–463
 - ambiphilicity of silylynes 462
 - dibromosilylene–NHC complex 461
 - *N*-heterocyclic silylenes 461
 - IPr–SiBr₂ adduct 461
 - IPrSiCl₂ adduct 460
 - IPr–SiCl₂–B(C₆F₅)₃ 461
 - IPr–SiCl₂–BH₃ 461
 - NHC–silylene adduct 459
 - NHC-stabilized diaminosilylenes 461, 462
 - NHC-stabilized silaisonitrile 461

- NHC-stabilized silanones 461
- product of reaction between IPr-Si=Si-Pr and BH_3 463
- triaminosilanes 462
- tin and lead 466
 - NHC-Ge=Ge-NHC complex 466
 - NHC-stabilized tin-tin double bond 467
 - NHC-stannylene adduct 467
- NHC complexes of group 15 elements 467
 - arsenic and antimony 473
 - IPr-stabilized diarsenic complex 473
 - nitrogen 467
 - imidazolin-2-imines ligand 467
 - phosphorus 468
 - copper(I) complex of an NHC-stabilized phosphonium cation 471
 - iminophosphine-phosphazene heterocycle stabilized by IPr 472
 - NHC stabilized diphosphorous-boronium cation 468
 - NHC-stabilized parent phosphinidene 470
 - NHC-stabilized phosphorus(I) salt 469
 - P_{12} cluster generated by aggregation of P_4 with an NHC 469
 - phosphorus(0) 468, 469
 - phosphorus(I) 469–471
 - phosphorus(III) 471, 472
 - phosphorus(V) 472
- NHC complexes of group 16 elements 474
 - oxygen and sulfur 474
 - cyclopropenyl-1-ylidene-stabilized S(II), Se(II), and Te(II) mono- and dications 474
 - selenium 474
 - NHC-SeCl₂ adduct 475
 - NHC-1,2,5-selenadiazolium complex 475
 - tellurium 475
 - NHC-stabilized aryltellurenyl salt 476
 - NHC-TeI₂ complex 475
- NHC complexes of group 17 elements 476, 477
- NHC-copper boryl complexes
 - for reduction of CO_2 to CO 213
- NHC-copper catalyzed trifluoromethylation 230
- NHC-copper chemistry 195
- NHC-copper complexes 228
- NHC-copper systems 233
 - 1,4-additions 206
 - for 1,4-additions 206
 - alkyne
 - semihydrogenation 232, 233
 - allene formation 202–204
 - allylic substitution
 - aluminum reagents 219, 220
 - boron reagents 220, 221
 - Grignard reagents 217–219
 - zinc reagents 217
 - boration reaction
 - alkene 223, 224
 - alkyne 224–226
 - carboboration reactions 226
 - ketone and aldehyde 222, 223
 - carbene 221, 222
 - carbonyl derivatives, olefination of 226–228
 - carboxylation/carbonylation 213, 214
 - conjugate addition
 - aluminum reagents 209
 - boron reagents 209, 210
 - Grignard reagents 207, 208
 - zinc reagents 206, 207
 - copper-mediated cross-coupling reaction 228, 229
 - A₃ coupling reactions 231, 232
 - $[\text{Cu}(\text{Cl})(\text{IPr})]$ 205
 - [3+2] cycloaddition reaction 215–217
 - enones 1,4-reduction of 205, 206
 - fluoride chemistry 230, 231
 - history of 199, 200
 - hydrosilylation 200–202
 - hydrothiolation/hydroalkoxylation/hydroamination 210–212
 - nitrene transfer 221, 222
 - transmetalation 235–237
- NHC-Cp Mo(II) complex 416
- NHC-Cu complex
 - asymmetric addition to imines 73–75
 - enantioselective addition of allenyl boron reagents to *N*-phosphinoyl imines 75
 - enantioselective allylation of *N*-phosphinoyl aldimines 74
 - asymmetric allylic substitution 67–69
 - cinammyl-type substrates 68, 69
 - asymmetric conjugate addition 65–67
 - alkyl boron reagents 67
 - cyclic enones 66
 - dialkylzinc reagents to enones 65, 66
 - synthesis of all-carbon quaternary stereocenters 67
 - asymmetric hydrosilylation 72, 73
 - ketones 73
 - enantioselective β -boration, asymmetric 70–72
 - chalcones 71

- enantioselective hydroboration of alkenes 72
- enantioselective quaternary, boron-containing stereocenter synthesis 72
- esters 70, 71
- unsaturated amides 71
- silyl conjugate addition 69, 70
- enones 70
- NHC–gold(I) complexes 78
- NHC–gold(I) derivatives 191
- NHC halogen chemistry 476
- NHC–Ir complex 53
 - asymmetric hydrogenation
 - of methyl 2-acetamidoacrylate 54
 - asymmetric transfer hydrogenation of aryl alkyl ketones 54
- chiral ferrocenyl–NHC Ir and Rh complexes 54
- NHC ligands 192
 - *ab initio* molecular dynamics 28
 - bond dissociation energy 26
 - bonding modes occurring between NHCs and transition metals 31
 - chiral NHC ligands effective in asymmetric synthesis 30
 - electronic effects 31–35
 - average CO stretching frequency 32, 33
 - plots of electron density 34, 35
 - Tolman electronic parameter (TEP) 33
 - variation in Ir oxidation potential as a consequence of modification of 34
 - five-membered 26
 - model [IrCl(CO)₂(NHC)] complexes 27
 - on neutral gold complexes 176
 - percent of buried volume
 - calculated as crystallographic data 28
 - (NHC)Ir(CO)₂Cl complexes 27
 - phosphines 27
 - placement of tertiary phosphines 28
 - from quantum mechanically optimized structures of [IrCl(CO)₂(NHC)] 27, 28
 - sphere used to calculate 26
 - SImes and the SIPr ligands, comparison between 30
 - steric effects 26–31
 - steric map, in optimized geometry of selected Ru complex 29, 30
 - Tolman cone angle 26, 28
- NHC–mercury complex, synthesis 2
- NHC–metal-based complexes 196
- NHC Mn(I) complexes, neutral and cationic 418
- NHC Mo and W complexes
 - containing a cyclopentadienyl ligand 415
- NHC–Ni complexes 55
 - asymmetric reductive coupling
 - of aldehyde and allene 55
 - of aldehydes and alkene 56
 - of aldehydes and alkynes 55
- (NHC)PdCl₂(3-chloropyridine) complexes 85
- NHC–Pd complexes
 - asymmetric α -arylation of amides 56–59
 - containing heteroatom substituents 58
 - intramolecular using NHC ligand 56
 - preparing 3-allyl oxindole 60
 - synthesis of *Ra,Ra* diastereomer and its use in 61
 - using six-membered NHC 59
 - asymmetric C–H activation coupling
 - with cyclic alkane 61, 62
 - resolution via 62
 - asymmetric diamination 62
 - alkenes 63
 - diastereomeric catalysts in asymmetric α -arylation 60
 - other asymmetric catalysis using 63–65
 - asymmetric allylic alkylation using NHC–Pd 65
 - asymmetric intramolecular oxidative amination 65
 - tested against cancer cells 106
- (NHC)Pd(II) complexes 86–89
- NHC–Pd–PEPPSI complex 85
- NHC–phosphine palladium complexes 103
- NHC/phosphine system 200, 215
- NHC platinum(0) catalysts 120
- NHC platinum complexes
 - characteristic of 123
- NHC platinum(0) complexes 115
 - kinetic assays 115–120
 - synthesis 115–120
 - synthesis of 115–120
- (NHC)Pt(dvtms) complexes 122
 - activation of 124
 - activation period 123–125
 - catalyst deactivation pathways 125–127
 - features of 122
 - proposed mechanism for activation 124
 - quantitative kinetic modeling 129–133
 - semiquantitative kinetic studies 127–129
 - syntheses of 116
 - x-ray structures of 117, 118
- NHC reactivity, with protic reagents 477
 - IMes–TEMPO–H complex 477
- NHC–Re(VII) complex 421
- NHC–Rh complexes

- asymmetric catalysis using boronic acids as nucleophiles 48–50
 - asymmetric addition of boronic acids to aromatic aldehydes 49
 - asymmetric addition of boronic acids to arylaldehydes 49
 - asymmetric 1,4-addition of organoboron to enone 48
 - asymmetric hydroarylation of azabicycles 48
 - asymmetric hydroformylation 53
 - asymmetric hydrosilylation 50–53
 - acetophenone/pyruvates 52, 53
 - acetophenone using chiral 1,3,4-triazolium ligand 51
 - carbene complex with axial chirality and use in 51
 - synthesis of diastereomeric complexes 50
 - NHC–rhenium (VII) complexes 420
 - NHC–Ru complexes 40
 - asymmetric hydrogenation 44–47
 - quinoxalines/benzofurans 46
 - regioselectivity in NHC–Ru catalyzed hydrogenation 45
 - synthesis of ent-corsifuran A 46
 - thiophenes/benzothiophenes 47
 - asymmetric hydrosilylation 47
 - ketones 47
 - asymmetric metathesis 40, 42–44
 - NHC–silver complexes 220
 - NHC–silver species 236
 - NHC-stabilized magnesium-hydride cluster 436
 - NHC transfer reagent 217
 - NHC vanadium adduct complex 411
 - nickel(0) complexes stabilized by π systems 374
 - nickel–NHC catalysts 372
 - *in situ* methods to generate Ni–NHC complexes 372
 - nickel–NHC complexes 371
 - nickel(0)–NHC complexes stabilized by π systems 373
 - Ni(II)/Ir(III) heterodimetallic complex 285
 - 4-nitrobenzaldehyde 64
 - nitrogen-stabilized acyclic carbenes 500
 - NMR spectroscopy 2
 - Nolan's A_H angle 136
 - Nolan's [Pd(IPr)(allyl)Cl] complex 62
 - nondiaminocarbene species employed as ligands 18
 - NP catalysts 63
 - nucleophilic carbon compounds 517
- o**
- 1-octene, hydrosilylation
 - bis(trimethylsilyloxy)methylsilane 138
 - kinetic profile for 142
 - kinetic reaction profile 125, 126
 - optimized conditions 138
 - silane conversion 143
 - silane conversion curves 120
 - silane conversion curves for 115, 119, 120
 - terminal alkynes catalyzed 139
 - [1-octene]/[MD'M] ratio 128
 - 1-octyne, hydrosilylation of 134
 - olefins 78, 112
 - cross-metathesis
 - low E:Z ratios 510
 - hydrogenation 113
 - isomerized 112
 - metathesis 28, 307, 510
 - standard substrates used for 312
 - polymerization 398
 - oligomerization reaction 102
 - organoberyllium–NHC complex 434
 - organoborane–NHC complex 435
 - organoboron reagents 222
 - organocatalysis 1
 - organocatalysts 63
 - organometallic chemistry 1
 - ovarian cancer line OVCAR-3, 164
 - *in vivo* xenograft model 164
 - oxazolineimines 56
 - oxazol-4-ylidene 273
 - oxidation reactions 100–102
- p**
- palladium 85, 97
 - catalysts 39
 - Chugaev-type chelated bis (ADC) complexes of 501
 - complexes 59, 60
 - Pd[(^{Bu}CC^{meth})₂Pd][BF₄]₂ 95
 - Pd(II)–diNHC complexes 94
 - Pd(II) hetero-diNHC complexes 94
 - Pd(II) ion 285
 - [Pd(L)₂](PF₆)₂ 95
 - PEG-tagged water-soluble Ru metathesis complex 336
 - Penicillin-resistant *Staphylococcus aureus* 151
 - pentamethylbenzene 97
 - pentamethylcyclopentadienyl variant 346
 - phenylacetylene 346
 - phenylboronic acid 49
 - phenyl cyclohexenone 208
 - 1-phenylethanol 344

- N*-phenylpyrazole 359
 2-phenylpyridine 359
 phenyl rings 186
 phosphine-free bisimidazolyldiene complex 313, 314
 phosphine ligands 114, 115, 307
 (phosphine)Pt(dvtms) complexes 114
 – selected syntheses 116
 – selected x-ray structures 117, 118
 – synthesis of 114
 phosphines, steric property 26
 phosphine systems 211
N-phosphinoyl imines 73, 75
 pH-responsive NHC ligand 317
 picolyl-derived *N*-methyl-substituted NHCs 343
 pinacolborane 479
 – oxidative addition of the B–H bond in 479
 Pincer ligands 93
 PIN ligand 364
Plasmodium falciparum 177
 platinum dimer, synthesis of 125
 platinum(II) carbene complexes 112
 platinum NHC complexes
 – hydrosilylation reaction 119
 platinum–NHC complexes 122, 123
 pMG101 plasmid 155
 PMHS (polymethylhydrosiloxane) 235
 polyamination, of aryl dichlorides with primary aryl amines 106
 polycondensation, of haloarylketones 106
 polylactic acid (PLA) 160
 poly(lactic-co-glycolic acid) (PLGA) 160, 167
 polymeric nanoparticles (NPs) 160
 polymeric silver–NHC compounds
 – synthetic route 152
 polymerization reaction 102
 – of ethylene upon activation by MAO 405
 – fluorene-based monomer 105
 – functionalized norbornene 104
 – 1-hexene 399
 – by hydrosilylation induced by (IMes)Pt(dvtms) 122
 – isoprene 399
 – olefin 104
 – poly(9,9-dioctyl-2,7-fluorene) 104
 – “ring-expansion polymerization” (REMP) 321
 – ring-opening metathesis polymerization (ROMP) 308
 polymer-supported (NHC)–Pd complex 88
 polynorbornenes 105
 potassium–NHC–amide complex, dimeric 433

*p*_π–*p*_π delocalization 4
 propargyl alcohols 224
 propargylamines, formation 232
 propargyl esters
 – dynamic kinetic asymmetric transformation of 515
 protein cdc2, phosphorylation 195
 protein tyrosine phosphatase (PTPs) 183
 – IC₅₀, comparison of 184
 prototypical carbodicarbenes 519
 prototypical carbene
 – resonance forms of 518
 PtCl₂(cod) 137
 Pt–H–Pt bond 126
 PTPs, complexes active on 183
 pyracenebis(imidazolyldiene) 287
 1-(pyridin-2-yl)benzimidazolium salts 97
 2-pyridylidene-based rhodium complexes 272
 pyrimidazolyldienes 6
 pyrimidine–NHC complex 343
 pyruvates 52, 53

q
 quantitative kinetic modeling 129–133, 142, 143
 quaternary chiral centers, formation 210
 quinoxalines 46

r
 RCM. *See* ring-closing metathesis (RCM)
 Re bis(carbene) complex 420
 1,4-reduction, mechanism of 205
 reductive coupling 390
 – aldehydes and alkynes 391, 392
 – aldehydes and allenes 392, 393
 – aldehydes and dienes 390, 391
 – aldehydes and norbornene 393
 regioselectivity
 – hydrogenation of aromatic carbocyclic ring of substituted quinoxalines 45, 46
 – impact of NHCs on 134–137
 – influence of silane on 137, 138
 – in NHC–Ru catalyzed hydrogenation 45
 release–return metathesis, mechanism 310
 remote *N*-heterocyclic carbenes (rNHCs) 517
 REMP. *See* ring-expansion polymerization (REMP)
 representative coordination modes
 – resonance forms of 518
 Rh– and Ir–NHC-based complexes 271
 – bridging NHCs 282–285
 – complexes with NHC ligands with facially opposed coordination abilities 285–288

- catalytic applications 288
 - arylation and borylation reactions with organoboron reagents 293–295
 - borrowing-hydrogen processes 292
 - cis-selected cyclopropanation reactions 298
 - dehydrogenation of alcohols 295
 - dehydrogenation of alkanes 295, 296
 - dehydrogenation of saturated CC and BN bonds 296, 297
 - H/D exchange reactions 296
 - hydroamination of alkynes 298
 - hydrosilylation 292, 293
 - hydrothiolation of alkynes 297, 298
 - magnetization transfer from para-hydrogen 298
 - reductions with H₂ 290–292
 - transfer hydrogenation 288–290
 - water oxidation 296
 - chelating NHCs 273
 - bidentate chelating bis-NHC complexes 273, 278
 - chelating chiral bis-NHC complexes 279
 - donor-functionalized chelating NHC complexes 280–282
 - mono-NHCs 271–277
 - rhodium (V)–NHC complexes 420
 - Rh(I)/Rh(III) homodimetallic complex 284
 - rhodium complex 48
 - ring-closing metathesis (RCM) 307, 315, 318, 324, 327, 328, 336. *See also* metathesis
 - ring-expansion polymerization (REMP) 321, 322
 - ring-opening metathesis polymerization (ROMP) 308, 309, 314, 320, 328, 336
 - ring-opening polymerization
 - of cyclic esters 398
 - of lactide 400
 - ring-opening polymerization (ROP). *See also* polymerization reaction
 - lactide in presence of Zr complex 406
 - *rac*-lactide 406
 - trimethylene carbonate (TMC) 406
 - Rochow–Müller process 111
 - ROMP. *See* ring-opening metathesis polymerization (ROMP)
 - ROP. *See* ring-opening polymerization (ROP)
 - R₃Si dimer, initiation 126, 127
 - R₃SiH
 - silicon–hydrogen bond 113
 - Ru–alkoxide complex 346
 - Ru–carbene complex 307
 - [Ru(cod)(2-methallyl)₂]/[NHCH][X]/base system 348
 - regio- and enantioselective hydrogenation by 348
 - Ru(II)–Ir(I) heterodimetallic complex 288
 - Ru-metathesis complexes bearing
 - four- and seven- membered ring NHCs 328
 - six-membered ring NHCs 327
 - Ru–NHC catalyzed addition, of carboxylic acids to terminal alkynes 359
 - Ru–NHC-catalyzed β -alkylation, of secondary alcohols with primary alcohol 352
 - [Ru(N,S-HC)(PPh₃)₂(O₂CR)X] complexes 346
 - Rupert–Prakash reagent 230
 - [Ru(py-NHC)(terpy)(OH₂)]²⁺-catalyzed alkene epoxidation 364
 - ruthenium 85
 - based alkylidene catalyst 40
 - based metathesis, simplified mechanism 308
 - catalyzed trans addition 133
 - variations of NHC in ruthenium complexes 313
 - ruthenium *N*-heterocyclic carbene (NHC) complexes 341
 - alcohol racemization 356, 357
 - allylic substitution reactions 361–363
 - arylation 357–359
 - borrowing hydrogen 351–356
 - direct hydrogenation/hydrosilylation 346–350
 - isomerization of C=C bonds 360, 361
 - miscellaneous reactions 363–365
 - reactions of alkynes 359, 360
 - transfer hydrogenation 341–346
- S**
- salicylaldiminato-functionalized imidazolydene ligands 285
 - scandium complex 399, 401, 402
 - molecular structure 401
 - scandium derivatives 400
 - second-generation catalyst
 - for hydrosilylation of alkynes 138, 139
 - limitations 139
 - semiquantitative kinetic studies 127–129
 - seven-membered NHC 92
 - silane
 - electronic properties 137
 - regioselectivity in 1-octyne hydrosilylation 137
 - substituted vinyl aluminum reagents 209

- silicon-hydrogen bond 111
- siloxane polymers 122
 - cheap silane mimicking 112
- silver-NHC complexes 157, 158
 - Ac-DEX NP formulations 163
 - acetylated dextran (Ac-DEX) 162
 - into acetylated dextran NPs 163
 - as anticancer agents 165
 - anti-infective 160
 - antimicrobial applications 160
 - antimicrobial efficacy 158
 - antiproliferative effect 164
 - coadministration of 159
 - electronically tuned 155
 - encapsulation of 166
 - formation of 154
 - hydrophilic and hydrophobic 157
 - hydrophilic PEG portion 167
 - MTT assay 168
 - into nanofibers 153
 - NP system for encapsulating 161
 - nticancer effects *in vitro* and *in vivo* 169
 - from Roland and Jolival 159
 - schematic depicting core-loaded 162
 - silver-NHC-encapsulated nanoparticles 169
 - systemic delivery of 166
 - from the Tacke group 158
- silver-NHC systems 195
- silver *N*-heterocyclic carbene (Ag-NHC) complexes 151
 - as anticancer agents 163–170
 - as antimicrobial agents 152–163
 - use of 152
- silver nitrate (AgNO₃) 152
- silyl ether protecting groups 122
- SIMes (1,3-bis(2,4,6-trimethylphenyl)imidazolidin-2-ylidene) 3, 28, 200, 202, 215
- SiO₂-NHC-Cu complex 231
- SiO₂-NHC-Cu(I) system catalyzing A³ coupling 231
- SIPr (1,3-bis(2,6-diisopropylphenyl)imidazolidin-2-ylidene) 203
- six-member ring NHCs 93
- SN2' allylic silylation 69
- sodium diethyl-2-methylmalonate 97
- “soft” carbene 282
- Sonogashira coupling 90–92
 - with *in situ* generated Pd-ADC catalyst 508
- Sonogashira couplings of aryl bromides 508
- Speier catalyst 122
- Speier's catalyst 111
- stable borylene 480
- Stenotrophomonas maltophilia* 155
- Stephan's group 347
- steric effect 25, 134
- stoichiometric reactions 235
- Stryker's reagent 200, 202
- styrene 347
- 4,5-substituted NHC-bearing complexes 332
- sulfur compounds 212
- SUNE1 (nasopharyngeal carcinoma) cancer cell lines 185
- superparamagnetic Fe₃O₄-supported NHC-based catalyst 64
- Suzuki-Miyaura cross-coupling reaction 85–88, 95, 96, 378–380, 505–508
 - acyclic carbene complexes 505
 - arylation of *meso*-chlorotricarboindocyanine dyes 508
 - of aryl bromides 96
 - palladium-ADC precatalysts 507
 - at Pd loadings 507
 - with phenyl boronic acid 507
 - with *in situ* generated Pd-ADC catalyst 506
- switchable carbenes 247
- t**
- Tamao-Kumada oxidation 122
- tantalum complexes 413
- TCPTP exhibit 183
- technetium (V) dioxo tetraNHC complex 419
- telomerization reaction 102
 - 1,3-butadiene 102
 - of butadiene with primary and secondary amines 103
 - – (NHC)Pd(dvds) complexes 104
 - Pd-catalyze, of isoprene with alcohols 103
- TEP. *See* Tolman electronic parameter (TEP)
- terminal alkyne, semihydrogenation 233
- terpenoid compounds, synthetic routes of 104
- m*-terphenyl ligand 429
- tertiary phosphines, ancillary ligand 114, 115
- 5,6,7,8-tetrahydroquinoxaline 348
- tetrahydropyranyl ethers 121
- 5,6,7,8-tetrahydroquinoxaline 45
- tetrahydrothiophenes 349
- 2,2,6,6-tetramethylpiperidide (TMP) 437
- tetra-NHC dioxo Re(V) complex 421
 - molecular structure 421
- thermal stability, of NHC-Au(I) complex 246
- thermodynamic stability 4
- thermolysis 1
 - chloroform adduct of an NHC 20
- THF mono-adduct 399
- thiazol-4-ylidene 273

- thiazol-5-ylidene 273
 - thioesters 70, 72
 - thiolato derivatives
 - IC₅₀ (μM) measured for 193
 - thiolato–gold(I) complexes 192
 - thiophenes 47
 - thioureas 20
 - titanium (III) complex 410
 - comprising annulated derivative with a pendant NHC ligand 410
 - titanium NHC complex 403
 - Cl–C_{carbene} distances 403
 - Cl–Ti–Cl angle 403
 - molecular structure 403
 - monodentate adducts 402
 - reactivity with dimethyl zinc 404
 - van der Waals radii 403
 - titanium pincer complexes 410
 - titanium (III) tris(NHC) complex 405
 - TOF value 291, 294, 315
 - Tolman cone angle 26, 28
 - Tolman electronic parameter (TEP) 6, 273
 - vs. average CO stretching frequency 33
 - tolyl-substituted NHC 316
 - o*-tolyl system 28
 - comparison between steric maps 29, 30
 - distribution of ϕ_1 angle in Ru (pre) catalysts 29
 - single Me group in *ortho* position, reducing flexibility 28
 - Tomioka complex 208
 - TON value 289, 292, 294, 315, 342
 - transfer hydrogenation 341–346
 - of alkenes 345
 - transition-metal-based homogeneous catalysis 111
 - transition metals 39
 - catalysts 63
 - complexes 25, 28
 - transmetalation 235–237
 - development 237
 - transmission electron microscopy (TEM) 86
 - 1,2,3-triazole-4,5-dilylidene 284
 - triazolediylidene ligand 352
 - 1,2,3-triazoles
 - formation of 216
 - 1,2,4-triazoles 273
 - 1,2,3-triazolyliidenes 273
 - 1,2,3-triazoly-4-idenes 6
 - tricarbonyl Re(I) NHC complexes 420
 - tricyclic monocarbenes 54
 - tricyclohexylphosphine 315
 - ligands 308
 - tridentate NHC-phenoxide ligands 405
 - trifluoroacetic acid 97
 - tri(2-furyl)phosphine derivative 114
 - trimethylsilylacetylene 140
 - trimethylsilyl azide 468
 - trimethylsilyl chloride 399
 - 1,2,4-trimethyltriazol-di-ylidene ligand 287
 - trinuclear complex
 - cross-coupling reaction 229
 - tris (pentafluorophenyl)borane 430
 - tris(2,2,6,6-tetramethylheptane-3,5-dionato) yttrium 398
 - Trx, inhibition of 182
 - TrxR, inhibition of 188
 - tumorigenic cell lines, cationic complex 181
- u**
- unsymmetrical homodinuclear ruthenium catalyst 324
- v**
- vanadium (II) and (IV) NHC complexes 411
 - vanadium (V) imido NHC complex 412
 - vanadium (V) NHC complex 411
 - containing NHC bidentate/tridentate ligand 412
 - vinyl chlorides 310
 - β-vinylcycloalkanones 66
 - vinyl phosphine oxides 310
 - vinylsilane 124, 145
 - vinylsiloxanes 124
- w**
- water-soluble Ru metathesis complexes 336
- x**
- xanphos ligand 233
 - xantphos (4,5-bis(diphenylphosphino)-9,9-dimethylxanthene) 351
 - x-ray crystallography 2, 88
 - x-ray diffraction analysis 397
 - x-ray photoelectron spectroscopy (XPS) 3
 - 1,3-xylyl-bis-imidazolium salt 284
- y**
- Y–NHC chelate complex 400
 - pseudotetrahedral geometry 400
 - yttrium alkyl complex 399
 - yttrium amido NHC complexes 400
 - yttrium–NHC adduct 400
 - yttrium NHC-phenoxide complex 401

Z

Z,E-alkenes 220

zinc reagents 206, 207

– allylic substitution 217

zirconium methyl derivatives 408

Zr chlorobenzyl derivative 406

Zr complex

– aminolysis 408

– comprising annulated derivative with a pendant NHC ligand 410

– hydroamination 409

– synthesis 408

– unexpected rearrangement 407

Zr metallocene complex 404

Zr-NHC pincer complex

– synthesis 409

Z-selective metathesis complexes

321

zwitterionic acyclic carbenes 520

