

Index

a

- ABA. *see* abscisic acid (ABA)
- abiotic stresses 28, 58, 82
 - catechol 943
 - antioxidant effect 943
 - conditions 31
 - enzymes 943
 - biosynthesis 943, 944
 - isochorismate synthase gene (*ICS1*) 943
 - *NahG* transgenic lines 943
 - promotive effect, salicylic acid 943
 - resistance to 31
 - salicylic acid treatment 943
 - results 943
 - seed germination 943, 944
 - *Arabidopsis* 943
 - inhibition 943
 - tolerance (*see* abiotic stress tolerance)
 - WRKY (*OsWRKY13*) and *OsNAC1* signaling pathways 38
- abiotic stressors 31
 - causing accumulation of ROS in plant cells 735
- abiotic stress tolerance 66
 - hybrids 821
 - plants genetically engineered for
 - through overexpressing antioxidant enzymes 261, 262
 - source of genetic materials 820, 821
 - weeds 820, 821
 - genetic studies 821
 - importance 820
 - potential but untapped 821
 - strategies to ensure survival 820, 821
 - vital ecosystem service 820
- ABRE-binding factor 424, 461, 529, 556, 621, 622, 649
- abscisic acid (ABA) 31, 326, 422, 459–461, 592, 694
 - accumulations in plants 423
 - biosynthesis and transport 83, 84
 - dependent pathway 442, 449, 615, 621, 659–662
 - important roles 555
 - independent pathway 441, 442, 448, 555, 556, 564, 595, 659, 662, 663, 765, 916
 - induced gene transcription 620
 - insensitive 5 (*ABI5*) in promoter 621
 - responsive element-binding factor 424, 438
 - responsive promoter element 438
 - signaling pathways 82, 272, 277
- Acacia saligna* 369
- acetoin 332
- acetylation 276
- ACL5* gene 43
- acrolein 257
- actual crop evapotranspiration 6
- acyl-CoA oxidase (*ACX*) 329
- adaptation 28
 - adaptive evolution, driven by environment 274–278
 - adaptive response in plants 263
 - plant adaptation to stress 66
 - strategies in the Indo-Gangetic plains 65
- ADC* gene 43
- Aeluropus littoralis* 687
- aerodynamic conductance 6, 12
- aerosols 5, 251
- AFPs. *see* antifreeze proteins (AFPs)
- Agave sisalana* 367
- Agave tequilana* 367
- agglutinins 335
- AgNHX1* gene 694
- agricultural insurance 74, 75
 - index-based insurance 76
- agricultural management 403
- agriculture 5
 - AgGDP 62

- climate change 816–820
- climate-resilient agriculture 73
- conservation 71
- expansion 346
- feedstock from 365
- increasing feminization 62
- organization 422
- production (*see* agriculture productivity)
- sustainability 16
- agriculture productivity 5, 28, 58
- factors adversely impacting 58
- and stability 58
- trend of change and impact on 59–62
- Agrobacterium tumefaciens* 865
- mediated methods 371, 575, 665, 788
- agro-biodiversity 58
- agrochemicals 365
- agro-ecological security 65
- agroecosystems 368, 396, 403
- C balance 380
- land use 381
- agroforestry 363, 394
- agronomic management, of plants 17
- Agropyron longatum*
- *AeNHX1* under salt stress conditions 693
- AHA1/OST2* mutant 85
- ahk1* mutants 88
- Aizoaceae 686
- AKR4C8 and AKR4C9 proteins 468
- albinisms 790
- aldo-keto reductases (AKRs) 467
- alfalfa 687
- algae 348
- high-lipid species 368
- Al-induced genotoxic adaptation 268
- allene oxide cyclase (AOC) 328
- allene oxide synthase (13-AOS) 328
- Allium cepa* 263
- Al-induced genotoxic adaptation 271
- Ca^{2+} channel chelator/blockers blocking genotoxic adaptation 269
- comet assay in root cells 264
- pretreatments of protein kinase inhibitors 268
- Aloe biology 863, 864
- adventitious root systems 863
- antifreeze compounds, producing 864
- bioenergy production 864
- cold stress 863
- drought resistant species 863
- high temperature tolerance 863
- important component of 863
- morphological variability 863
- plant features 863
- soil requirements 864
- succulent plants 864
- water reserves 863
- wax-coated surfaces, leaves 863
- Aloe* tissue culture, cloning and mass propagation 866–868
- aloe propagation 866
- *Aloe vera*, cultivation 866
- large-scale cultivation 866
- callus induction, 6-benzylaminopurine, (BAP) 867
- cloned plants 867
- enhanced productivity, improvement 870
- explants, stem segments 867
- *ex vitro* autotrophic cultures 867
- intersimple sequence repeat (ISSR) marker 867
- micropropagation protocol 868
- natural propagation 866, 867
- plant tissue culture technology, non-GMO biotechnology for 866
- random amplified polymorphic DNA (RAPD) marker 867
- results 872, 873
- somatic organogenesis 867
- “tailor-made” biotechnology 867
- tissue culture-raised plants 867
- vegetative propagation/propagules methods 866
- *in vitro* production, aloe-emodin 867
- Aloe vera* clones plantlets
- acclimatization 871, 872
- acclimatized cloned plants 872
- culture technique 871
- hardened cloned plants 872
- hardening 871, 872
- polycarbonate caps 872
- *in vitro* generated plantlets 871
- Aloe vera*, micropropagation. *see Aloe* tissue culture
- ALTERED MERISTEM PROGRAM1-like peptidase 533
- alternative oxidase pathway, salicylic acid regulation 956, 957
- alternative oxidases (AOX) 335
- aluminum toxicity 315
- Amaryllidaceae 366
- γ -amino butyric acid (GABA) 706
- 1-aminocyclopropane-1-carboxylate (ACC) deaminase 427
- AmMYB1 transcription factor 620
- Ananas comosus* 367

- androgenesis, for crop improvement 787, 788
 - *Agrobacterium tumefaciens*-mediated methods 788
 - combination of androgenesis and plant transformation methods
 - using abiotic stress-tolerance genes 792
 - enhancement of abiotic stress tolerance, in cereal crops through 793–800
 - factors influencing 788
 - anther walls 788
 - culture medium 789
 - genotype and other physical conditions of donor plant 788
 - ploidy status, confirmation of 789
 - pretreatment/stress factors 789
 - stage of microspore/pollen development 789
 - functional microspore transformation systems 788
 - genetic transformation and in combination with 790, 791
 - major abiotic stress-tolerant crops, development of 791, 792
 - drought 801
 - extreme temperature (cold/heat) 802
 - flood/water logging 802
 - heavy metals 801, 802
 - herbicide resistance 803
 - osmotic and oxidative stress 803, 804
 - salinity 792, 801
 - micro-projectile bombardment 788
 - problems associated, with albinisms 790
- Andropogon gerardii* 114, 118, 119, 121, 123–126, 128
- anoxia, in plants 151, 608
 - NO formation during 152
- antagonistic/synergistic interactions 1027, 1028
 - ACC deaminase 1027
 - antifungal activity, endophytic bacterial 1027
 - antimicrobial activity compounds 1027
 - bacterial endophytes activity
 - against fungal pathogens 1027
 - important influence, endophytic bacteria 1027
 - insecticidal activity compounds 1027
 - plant carbohydrates uptake 1027
 - *in situ* methane recycling 1027
 - wide-spectrum antifungal activity 1027
- anther culture method 788
- anthocyanins 714
- anthropogenic influences 858
- antifreeze proteins (AFPs) 472, 473, 710
- antioxidant/antioxidant enzymes, role during abiotic stress in plants 709
- antioxidant defense mechanism 822
- antioxidant enzymes 194, 255, 332, 444, 1049.
 - See also* antioxidant systems
 - activation 1049
 - ascorbate peroxidase 195, 443
 - catalase 195, 234, 235, 443, 968, 969
 - endophytic bacterium infection 1049
 - glutathione peroxidase 443
 - glutathione reductase 195, 443, 996
 - includes 1049
 - overexpression in genetically engineered plants
 - for abiotic stress tolerance 261, 262
 - salt stress conditions 1049
 - secondary effects stress 1049
 - superoxide dismutase 194, 195, 202, 234, 235, 443, 966, 968
- antioxidant systems 708, 709, 739, 740, 959
 - antioxidant enzymes 959
 - CAT activity 959
 - temporary reduction 959
 - drought-stressed plants 959
 - salicylic acid spray 959
 - lipid peroxidation 959
 - mitochondria 959
 - adenosine tri-phosphate (ATP) production 959
 - mitochondrial enzyme activity 959
 - reactive oxygen species (ROS) 959
 - increased level 959
 - metabolism 959
 - redox homeostasis 959
 - regulation, salicylic acid 959
 - salicylic acid 959
 - application 959
 - foliar application 959
 - treatment 959
- antioxidative defense system 42, 296, 608
- AOC oligomerization 334
- Aoe, biodiversity
 - commercial harvesting threat 864
 - genetic resources 864, 865
 - international aloe science council (IASC) 864
 - Aloe Scientific Primer 864
 - international union for conservation of nature (IUCN) red list 864
 - endangered species 864
 - threatened species, IUCN Red List 864, 865
- AP2/EREBP domain family 556
- AP2/EREBP domain proteins 562

- AP2/ERF family
 - DNA-binding domain 651, 652
 - AP2/ERF proteins, subdivided 651
 - AP2-integrase DNA-binding domain (AP2-IDBD) 651
 - Arabidopsis homeotic gene APETALA2 651
 - *Chlamydomonas reinhardtii* 651
 - dehydration-responsive element (DRE)/C-repeat (CRT) 652
 - dimerization, plant AP2 domains 651
 - DREB1/CBF, DREB2 652
 - ERF subfamily members 652
 - floral meristem 651
 - genes 651
- AP2/ERF transcription factors 328, 607
- Apocynaceae 366
- apoplastic protein WPM-1 445
- apoptosis 296, 441
- AP2 protein 562
- Arabidopsis thaliana* 142
 - ABA signaling 269
 - activation of ATM and ATR kinases 270
 - AKR4C8 and AKR4C9 proteins 468
 - AOC oligomerization in jasmonate production in 328
 - AREB1/ABF2 in transgenic 529, 530
 - bZIP transcription factors 531
 - CBF1, CBF2, and CBF3 genes 564
 - CBF/DREB1 genes 564
 - CBF/DREB regulation in 565
 - CBF genes 611
 - CBF genes 563
 - Cd-induced increases of NO 147
 - CDPK transcripts 269
 - cold-exposed 150, 151
 - COR15A gene 564
 - crystal structure of OST1 from 463
 - DREB1A gene 575
 - DREB1 genes 563
 - drought caused upregulation of miR396 in 514
 - expression of CBFs, modulated by temperature 565, 566
 - expression of miR398 513
 - gene, *eskimo1* (ESK1) 570
 - genes identified as encoding transcription factor 556
 - miR398 743
 - miR169 downregulated in 514
 - miRNA families to be upregulated by UV-B radiation 490, 514
 - NAC genes overexpression 595
 - overexpressing *AtMYB44* 617
 - overexpressing CBF3 (DREB1A) 569
 - P5CS genes 570
 - P5CS2 transcript levels 569, 570
 - proline levels 731
 - promoter regions of CBF/DREB genes 565
 - protoplast, ABA require activation of 529
 - PYR1 protein 461
 - regulation of CBF pathway 566, 567
 - regulatory pathway of DREB1A 556
 - *sfr6* mutant 569
 - to stress (UV-C) 275
 - stress-inducible promoter *rd29A* 609
 - structure of AKRs 468
 - studies on root system 307
- AREB/ABF regulon 530, 593, 621–623
- L-arginine 142
- arid environment plants 935. *See also in vitro* cultures establishment
 - *Blepharis indica* 935
 - *Haloxylon recurvum* 935
 - *Haloxylon salicornicum* 935
 - *Leptadenia pyrotechnica* 935
 - shoots 936
 - *in vitro* characterization 936
 - *in vitro* multiplication, shoots 936
- arid lands (drylands), challenges 859, 860
 - characteristics 860
 - climatic boundaries 860
 - climatic characteristics 860
 - critical importance 860
 - ecosystem multifunctionality 860
 - inhabited drylands 860
 - key biomes 860
 - largest terrestrial ecosystem 859
 - plant biodiversity, conservation 860
- ARP2/3–SCAR pathway 298
- Artemisia annua* 148
- Arthrocnemum indicum* 686
- artificial miRNAs (amiRNAs) 494
- artificial ta-siRNAs (ata-siRNAs) 494
- Arundo donax* 369
- Asclepiadaceae 366
- ascorbate–glutathione cycle 255
- ascorbate–GSH cycle, 234
- ascorbate peroxidase (APX) 141, 255, 709
- L-ascorbic acid 255
- aspartate aminotransferase (AspAT) 911
- Aspergillus niger* 263
- AtHKT1* gene 445
- ATM (ataxia telangiectasia-mutated) 270
- atmosphere 815
 - chemical composition 815, 816

- atmospheric CO₂ level 818
 - anthropogenic activities 818
 - C3 plants 818
 - vs. C4 plants 818
 - increasing level 818
 - benefits 818
 - experiments 818
 - ATM signaling kinases 272
 - AtMYB61* expression 618
 - AtMYB44* transcription factor 617
 - Atnoa1* mutant 140
 - ATP/ADP ratios 152
 - ATP-binding cassette(ABC) protein 85
 - ATP consumption 272
 - ATR (ATM-Rad3-related) 270
 - Atriplex griffithii* 686
 - Atriplex hortensis* L. *see* Garden Orache
 - Atriplex nummularia* 692
 - ATR signaling kinases 272
 - autonomous pathways
 - early-flowering phenotype 952
 - flowering
 - constans (CO), key regulator 951
 - long-day conditions 952
 - phytohormones/growth regulators 952
 - short-day conditions 952
 - flowering locus *C* (*FLC*), flowering repressor 951
 - expression 951, 952
 - gibberellin signaling pathway 952
 - integrators 951
 - flowering locus *T* (*FT*) 951
 - suppressor of overexpression of constans 1 (*SOC1*) 951
 - transcripts 951
 - late-flowering phenotype 952
 - photoperiod 951, 952
 - photoperiod pathway 952
 - salicylic acid 951
 - deficient plants 951
 - exogenous application 951
 - flowering regulation 952
 - genes transcription regulation 951
 - interaction, genetic analysis 952
 - 100 μ M application 952
 - wild-type plants 951, 952
 - autophagy, leaf senescence 961–963. *See also* senescence
 - ATG (autophagy) genes 961
 - ATG5–phosphatidylethanolamine complex 962
 - controversy 962
 - dark-induced senescence 962
 - ethylene signaling pathway 962
 - mutations 962
 - importance 961
 - induction 961
 - jasmonic acid signaling pathway 962
 - mutations 962
 - NPR1 gene 962
 - allele 962
 - monomer functions 962
 - mutant transcript levels 962
 - paralog 962
 - redox-sensitive mechanism 962
 - salicylic acid 961
 - function in plant growth and development 962, 963
 - responsive defense genes 961
 - senescence phenotype 961
 - auxin 142, 332
 - auxin indole acetic acid (IAA) 948
 - auxin response factors (ARFs) 592
 - average surface temperature 817
 - heat stress, plants exposure to 817
 - harmful effects 817
 - high-temperature regimes crops 817
 - high temperatures impact 817
 - short-term temperature extremes, changes 817
 - average yield, of crops 210
 - Avicenniaceae 686
 - Avicennia marina* 687
 - Avicennia officinalis* 189, 190
 - antioxidant enzymes, role in high salinity 203
 - artificial regeneration 190
 - effect of increasing soil salinity on total dry weight 196–199
 - osmotic adjustment 201
 - reduction in growth 202
 - salinity tolerance
 - antioxidant enzymes activities 199, 200
 - chlorophyll content 196–198, 201
 - experimental studies 189–203
 - lipid peroxidation and membrane permeability 198, 199
 - organic solutes 196
 - statistical analyses 195
 - total dry weight of plants 195, 196
 - salt tolerance mechanism 190
 - Azadirachta indica* 366
- b**
- Bacillus amyloliquefaciens* 427
 - bacterial endophytes 1016–1019

- commonly isolated and studied 1017
 - definition, endophyte 1016
 - environmental stresses 1017
 - hydrolytic enzymes 1016
 - latent pathogens 1017
 - mechanism of action 1018, 1019
 - 1-aminocyclopropane-1-carboxylate (ACC) deaminase 1018
 - ethylene-sensitive plant growth, inhibition 1019
 - indole acetic acid (IAA) 1018
 - mechanistic aspects 1019
 - plant growth promotion 1018
 - as sink, bacteria 1019
 - synergistic interaction 1019
 - multivariate analyses 1016
 - plant defence 1018
 - abiotic stress 1018
 - biologically fixed nitrogen, supply of 1018
 - biotic stress 1018
 - low-molecular-weight compounds, production 1018
 - plant defenses 1017
 - Proteobacteria phylum 1016
 - soil bacteria 1016
 - strains 1016
 - diazotrophic 1016
 - non-diazotrophic 1016
 - subset 1016
 - symbiotic interactions 1017
 - systematic biology 1017
 - Bangladesh 882
 - Bengali year seasons 882
 - high organic carbon storage, forests 883
 - barley domestication 333
 - nonbrittle rachis 656
 - six-rowed spikes 656
 - type, spikes 656
 - *HvHox2* 657
 - I HD-ZIP transcription factor, subfamily 657
 - model, spikelets 657
 - recessive allele *vrs1*. 657
 - yield 656, 657
 - base excision repair (BER) 263
 - basic helix–loop–helix (bHLH) transcription factor 43
 - basic region leucine zipper (bZIP) proteins 424, 556, 557
 - betaine aldehyde dehydrogenase gene 42
 - Beta maritima* 369
 - β -glucosidase 423
 - gene 327
 - bHLH family
 - class, plant transcription factors 649
 - amino acids 649
 - bipartite domain 649
 - E-box (CANNTG) 649
 - emblematic bHLH (MYC-type) 649
 - G-box (CACGTG) 649
 - MYC plant subfamily 649
 - biodiversity 57–59, 71, 346, 373, 380, 397, 399, 820, 864, 907, 1042
 - bioenergy 363
 - agroecosystems 347, 348, 352
 - plantations 347
 - potentials 347
 - production 366
 - sustainability of 347
 - bioenergy crops 347, 369, 394
 - biogeochemical liabilities of carbon in 381–383
 - as biological systems 347
 - breeders, need to optimize plant C allocation 372
 - carbon in 361, 362
 - first-generation 363, 364
 - traditional plants 362, 363
 - carbon management in 374, 375
 - climate change mitigation potential 355–358
 - biomass vs. bioenergy density 358–360
 - climate ethics and carbon in 391, 392
 - biofuel vs. biodiversity 397, 398
 - biofuel vs. food 392–394
 - biofuel vs. water 394–397
 - ecosystem services of carbon in 387–389
 - genetic improvement of 369, 370
 - genetic models and ideotypes 373, 374
 - lignocellulosic 396
 - recalcitrant carbon in 354, 355
 - sterility in 398
- biofertilizers 68
- biofuels 346, 352, 353, 363, 364, 379, 403
 - agroecosystems 352, 353
 - affect water cycling 395
 - C4 plants as biofuel feedstocks 384
 - crops with better GHG profiles 348
 - derived from first-generation crops 364
 - as energy carriers 347
 - on food production 347
 - production 399
 - with CO₂ sequestration 353
 - water consumption in 395
- biological control, to reduce CH₄ production from livestock 71
- biomass 42, 346, 364

- energy 352
 - biomass-to-liquid biodiesel 360
 - biorefinery 352
 - biosynthetic pathway, salicylic acid 940, 941
 - *Arabidopsis* systems 941
 - isochorismate pathway, chloroplast 941
 - methyl-salicylate (MeSA) 941
 - phenylpropanoid route, cytoplasm 940, 941
 - salicylic acid glucoside (SA2-O- β -D-glucoside) 941
 - biotechnology, Aloe 865, 866
 - abscisic acid (ABA) 865
 - exogenous application 865
 - characterization 865, 866
 - conservation 865, 866
 - genetic transformation 865
 - improvement 865, 866
 - molecular biology tools 865
 - NADP-malic enzyme (NADP-ME) 865
 - gene 865
 - phosphoenol-pyruvate carboxylase (PEPCase) 865
 - phylogenetic tree 866
 - ploidy levels 865
 - productivity enhancement 865, 866
 - somatic cells 865
 - transgenic plants 865
 - biotic stress 963–965
 - mitochondrial enzyme 964
 - activity alteration 964
 - effects on ROS levels 964
 - pathogenic attacks 963
 - defensive mechanisms 963
 - *PR* genes expression, inducing 964
 - resistant against pathogens of 964
 - salicylic acid
 - benefits 964
 - causes 964
 - exogenous application 963
 - lipid peroxidation 964
 - tobacco mosaic virus (TMV) 963
 - resistant tobacco variety 963
 - Bioversity International 934
 - black carbon 251
 - Blue Revolution 666
 - Botrytis cinerea* 327
 - Botrytis* susceptible 1 (BOS1) 327
 - β -oxidation 203
 - β -polymerase 265
 - Brachypodium distachyon* 373, 450
 - Brassicaceae 686
 - Brassica juncea* 147
 - Cd-, Cu-, or Zn-exposed roots 147
 - Brassica napus* 201, 473
 - *BN115* gene from 564
 - Brassica rapa* 511
 - heat stress, to alter miRNA expression 511
 - brassinosteroids 332, 536, 712
 - breeding crop varieties 66
 - breeding strategies
 - breeding lines 848, 849
 - breeding material 849
 - chilling-tolerant cultivars 850
 - heat tolerance 848–850
 - heat-tolerant chickpea cultivars 848
 - ICRISAT 849
 - screening method 849
 - marker-assisted selection 849
 - multilocation yield trials 849
 - pollen-selected progenies 850
 - pollen selection method 849
 - development 849
 - rapid generation turnover 849
 - Single-plant progenies 849
 - *in vitro* germination, pollen 849
 - Bruguiera gymnorhiza* 689
 - Burkholderia cepacia* 1028, 1029
 - 2,3-butanediol 332
 - bZIP family 649, 650
 - ABA-responsive element (ABRE) 649
 - ABRE-binding factors (ABFs) 649
 - ABRE-binding proteins (AREB) 649
 - abscisic acid (ABA). 649
 - amino acids 649
 - eukaryotes 650
 - modules 649
 - water-deficit and high-salt stresses 650
 - bZIP transcription factors 529, 621
 - phylogenetic tree 531
- C**
- cadmium 608
 - Caenorhabditis elegans* 482
 - Cairnsian mutation 274
 - Cakila maritime* 190
 - Cakile maritima* 692
 - calcineurin 463, 464
 - calcineurin/cyclophilin complex 464
 - calcineurin/ FKBP12 complex 464
 - calcineurin signal, inhibition by cyclophilin A 464
 - calcium 191, 269
 - calcium-dependent protein kinases (CDPKs) 86, 269, 591
 - calcium signaling, in genotoxic adaptation 267
 - calmodulin-binding protein 591

- calmodulins (CaM) 168, 269
- Calophyllum inophyllum* 366
- CAM photosynthesis pathway 367, 390
- CAM plants 127
- canopy 6
 - changes in size 13
 - conductance to water vapor transfer 6
 - energy balance 14
 - impacts on WUE 14
 - level process 12
 - morphology 13
 - warmer canopy temperatures 17
- Capsella bursa* 563
- Caragana korshinskii* 610
- carbohydrates 708
 - imbalance 30
- carbon (C)
 - assimilation 30, 367
 - in bioenergy crops 361, 362
 - budget inventories and sustainability assessment 366
 - C-neutrality 353
 - cycle 346, 350, 351, 364, 366
 - eco-physiology and carbon sequestration 389–391
 - in first-generation bioenergy crops 363
 - fixation 30
 - flux 130, 346
 - from soils 352
 - isotopes 348
 - life cycle assessment 385–387
 - management in bioenergy crops 374, 375
 - managing carbon sources and sinks 375, 376
 - managing land-use change 379–381
 - managing nutrient composition, cycling, and loss 377–379
 - in photosynthesis activity 349
 - quality in bioenergy crops 383–385
 - reserves 375, 397
 - roles 348
 - as scarce commodity 349, 350
 - in second-generation bioenergy crops 364–367
 - sequestration 69, 71, 72, 345, 356, 366, 367, 370, 373, 376, 388, 393, 399, 401, 402, 883
 - in soil 349
 - storage, forests 882
 - synthesis of research needs and priorities 398–403
 - temporal changes 360, 361
 - tracking and accounting process 385
 - uptake (*see* carbon uptake)
- carbon dioxide (CO₂) 4, 28, 251, 320, 507
 - concentrations, increase with 13
 - effects on RuBisCO 30
 - effects on stomatal conductance 14
 - effects on tolerance of photosynthesis to heat stress 121
 - elevated levels, and plant response 30
 - enrichment and nitrogen management, on grain quality 18
 - flux 366, 401
 - global increase of 251
 - increase ET from canopies 14
 - intercellular concentration 14
 - plant productivity 15
 - predictive rise 508
 - and rising temperature 345
 - sequestration 368
 - soybean exposed to 17
 - soybean grown in controlled environment chambers 14
- carbon monoxide (CO) 507
- carbon uptake
 - aspartate aminotransferase (AspAT) 911
 - carbon sequestration 910
 - climate change scenario 910
 - major concern 910
 - CO₂ uptake, efficiency 911
 - efficiency enhancement 910, 911
 - gas-exchange characteristics 910
 - results 910
 - mean annual temperature 910
 - mechanism to fix higher CO₂ 912
 - partial pressure 910
 - phosphoenolpyruvate carboxylase (PEPCase) 911
 - photosynthesis 910
 - primary products analysis 911
 - photosynthetically active radiation (PAR) 910
 - ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO) 911
 - activation state 911
 - total irradiance 910
 - water-use efficiency 910
- carboxypeptidase 140
- carotenoids 255, 707, 708
- catalase (CAT) 141, 145, 146, 148, 149, 176, 177, 195, 200, 232, 234, 235, 443, 709, 714, 732, 959, 966, 970
- catastrophes, of changing climate 589, 590
- CAT1 enzyme 442
- CBF1, CBF2, and CBF3 proteins 564
- CBF/DREB regulon 609–611
- CBF/DREB transcription factors 594, 595

- CBF expression
 - downstream regulators 569
 - COR 569
 - SFR6 569
 - negative regulators 568
 - positive regulators 567
 - FIERY 568
 - HOS1 568
 - ICE 567
 - LOS4 567, 568
 - MYB15 569
- CBF genes 447, 448
- CBF3 integrates activation of multiple
 - components, of cold response 569, 570
 - *eskimo1* (*ESK1*) gene 570
- CBF proteins 565
- CBFs/DREB proteins 564
- CCAFS programme 75
 - CCAFS CSV initiative 76
 - impact pathway approach, national and regional programmes 76
 - partnership 76
- Cd-induced genotoxic stress 267
- Cd toxicity in plants 147
 - exogenous NO, role in 147, 148
- cell death 257, 686
 - regulation 948
- cell growth regulation
 - *Arabidopsis* genome 948
 - cross-talk, phytohormones 948
 - plant defense responses 948
 - transcriptional regulator NPR1 948
- cell-to-cell signaling 268
- cellular dehydration 522
 - ABA-mediated ABRE-dependent gene expression 529, 530
 - major ABA signaling components in response to 522
 - SnRK2-AREB/ABF pathway, central role in ABA-mediated gene expression in response to 528, 529
- cellulose 96
- cellulosic feedstocks 365
- cell wall compounds 95
- Centaurea tuzgoluensis* 190
- cereal crops 438
 - abiotic stress-tolerant 785
 - agronomically important 1020
 - Al toxicity 802
 - enhancement of abiotic stress tolerance through 793–800
 - haploid production 788
 - ideotype 373
 - recalcitrant to recombinant techniques 785
 - research on polyamines in 42
 - salt tolerance 174
 - yield, likely to be reduced due to 61
- Ceriops tagal* 201
- cesA7* (*irx3-5*) mutant 96
- chaperone machinery, operates in cytosol 430
- chaperones 439, 471, 591
- chaperonin family proteins 429
- checkpoint kinases (CHKs) 270
- chemical mutagens 787
- Chenopodiaceae 686
- Chenopodium album* 687
- Chenopodium quinoa* 691
- Chernobyl nuclear disaster 274
- chickpea. *see* *Cicer arietinum* L.
- chilling stress 150, 151, 715
- chilling tolerance 315
- Chlamydomonas reinhardtii* 263, 277
- Chlorella pyrenoidosa* 263
- chlorofluorocarbons 251, 295
- choline dehydrogenase 42
- choline monooxygenase 731
- chromatin 44
- Chrysanthemum morifolium* 150
- C2H2 ZF protein 597
- Cicer arietinum* L. 840
 - growth ranges 842
 - optimal temperature 842
 - new cropping systems 841
 - production areas 841
 - production constraints 840, 841
 - reproductive phase 842
 - shift in production area, India 841
 - graph 841
 - source of 840
- cis*-elements 556
- 9-cisepoxy carotenoid dioxygenase 3 83
- Citrus aurantium* 141
- c-Jun N-terminal kinase 268
- CLEL peptides 333
- Cleome gynandra* 231, 235
- Cleome spinosa* 231, 235
- climate change, Himalaya 907, 908
 - air temperature, increase 908
 - average precipitation 908
 - biannual 908, 909
 - seasonal 908, 909
 - biodiversity, impact on 908
 - increased temperature 907
 - coupled with 907
 - IS92a future scenarios, greenhouse gas 907, 908

- location-specific data, accurate predictions 908
- long-term meteorological data analysis 908
- mean temperature 908
 - annual 908, 909
 - seasonal 908, 909
- normalized difference vegetation index 908
- permanent monitoring plots 908
- precipitation data 908
- regional climate model 907
- scenarios 907
- second-generation Hadley centre regional model (HadRM2) 907
- surface albedo associated 907
- warming 907
- climate change impacts, organic carbon storage
 - in forests 894–896
 - climate change impact, determination 894
 - hill forests 896
 - high-density trees 896
 - total biomass content 896
 - mangrove forest ecosystems 894
 - high-density mangroves 894
 - low-density mangroves 894
 - organic carbon storage 894
 - saline zone 894
 - salinity 894
 - organic carbon storage, village forests 896
 - plain land sal forests 896
 - high-density trees 896
 - sal coppices 896
 - total biomass content 896
 - Sundarbans
 - bush fires 895
 - intense tropical cyclones 895
 - organic carbon stock loss process 895
 - sea level rise impacts 895
- climate changes 706, 882, 933. *See also* climate change, Himalaya; climate change scenarios, Bangladesh
 - affects 882, 883
 - agriculture 816–820
 - and ecological destruction 786
 - effect of
 - developing countries 816
 - management 75
 - threats 882
- climate change scenarios, Bangladesh 887–889
 - anthropogenic greenhouse gasses 887
 - climate change models 888
 - climate change policies 888
 - droughts 888
 - floods 888
 - future projection 888
 - GCM projections 888
 - changes in precipitation 888
 - changes in temperature 888
 - global climate change 887
 - inter-model variations, scale of change 888
 - monsoon change 887
 - national adaptation program for action (NAPA) 888, 889
 - general circulation models (GCM)s 889
 - main objective 889
 - observed climate change phenomena 887
 - projected sea level rise 888
 - rising trend of 887
 - winters 888
- climate-resilient agricultural system 66
- climate-smart agriculture (CSA) 59, 63
 - community-based actions 73
 - Crisis Management Plan 73
 - development 73
 - FAO definition 63
 - Government of India, initiatives 73
 - location-specific, and community-based activities 73
 - South Asian Programme of CCAFS, establishing CSVs 74
 - components of system at village/farm level 74
 - technologies and strategies 63
 - adaptation 64–67
 - sustainable productivity enhancement 63, 64
- climate stresses 4
- climate warming 346
- cloning, *Aloe vera* (ghee-kanwar/gwar-patha) 868
 - *Aloe* plantation industry 868
 - *Aloe* products use 868
 - history of uses 868
 - micropropagation protocol 868
- C-neutrality 352
- C:N ratio 402
- Cnx1* gene 532
- cold/freezing stress 150, 151
- cold stress 715
 - MYBs and 618, 619
- cold stress pathway 662
- cold tolerance 447–449
- collective changes, in climate 6
- comparative genomics, of moss 277
- comparative qRT-PCR assays 335
- Compositae 366

- conditioning hormesis 263
 - consultative group on international agriculture research (CGIAR) 934
 - Convolvulaceae 366
 - COP13 Climate Change Conference 76
 - COP1* gene 298
 - copper 608
 - coral bleaching 62
 - COR gene expression 569
 - corn/soybean rotation
 - high-yielding management zones 15
 - coronatine insensitive 1 (COI1) 330
 - CO₂ sequestration 353, 368
 - cotton grown, under high temperatures 18
 - C3 plants 127, 231
 - growth 13
 - C4 plants 127, 231
 - crassulacean acid metabolism (CAM) 367
 - Craterostigma plantagineum* 473
 - C-repeat transcripts (CRTs) 441
 - Crisis Management Plan 73
 - crop diversification 71
 - crop domestication 654
 - barley domestication 656, 657
 - genetic changes 654, 655
 - maize domestication 654
 - monogenic traits 654
 - examples, genetic loci 654
 - genetic tools, crop improvement 654
 - phenotypes 654
 - rice domestication 655, 656
 - crop improvement 665–672, 666
 - blue revolution 666
 - increased yield, harvest 666
 - long and winding road 665–672
 - model 668
 - observed, drought tolerance 672
 - one trait acquisition 665
 - relationship, phylogenetic distance 668
 - timetree database 668
 - transgenic plant 669
 - crop management 403
 - cropping intensity 73
 - cropping systems 16
 - crop plants
 - genotypic variability 210
 - as sessile organisms 210
 - crop production 4, 17, 352
 - affect of climate change 830
 - crop productivity, under stressful environments 252
 - crop resistance, to insects 786
 - crop responses, to climate change 7
 - crop water demand 6
 - crop yield 29
 - cross-breeding, mutation-assisted breeding 787
 - CRT-binding factor (CBF) 438
 - Cruciferae 366
 - CSA. *see* climate-smart agriculture (CSA)
 - cultures establishment, *Aloe vera* clonal Shoots 868–871
 - additives, MS medium 871
 - antioxidant-treated shoots 871
 - axillary shoot segment, explants 871
 - botanical survey of India (BSI), Jodhpur 868
 - multiplication, clonal shoots 868–871
 - surface-sterilized explants 871
 - culture, agar-gelled semi-solid MS medium 871
 - *in vitro* produced shoots 871
 - customized fertilizers 71
 - Cu toxicity
 - exogenous SNP application 148
 - cyclophilin-like protein PiCypA 469, 470
 - cyclophilins 469
 - Cynara cardunculus* 367
 - Cynodon dactylon* 372
 - Cynodon* spp. 365
 - CYP707A3 genes 423
 - cytokinins 328, 332, 712
 - cytoskeleton-associated proteins 267
- d**
- Datura innoxia* 787
 - deforestation 16, 376, 380, 397
 - dehydration-responsive element-binding protein (DREBs) 424
 - dehydration-responsive element/C-repeat (DRE/CRT) cisacting element 256
 - dehydration-responsive elements (DREs) 441
 - dehydrins 31, 710, 711
 - dehydroascorbate reductase (DHAR) 255
 - desert crassulacean acid metabolism (CAM) succulents 121
 - desiccation tolerance 277
 - detoxification 203, 235
 - diamine oxidase 142
 - diazotrophic bacterial endophytes 1020–1022
 - agronomically important 1020
 - *Azospirillum seropedicae* 1020
 - physiological characteristics 1020
 - *Burkholderia* spp 1021
 - *Burkholderia brasiliensis* 1021
 - *Burkholderia vietnamiensis* 1021
 - DNA and rRNA hybridization analysis 1021

- graminaceous plants 1020
- *Herbaspirillum frisingense* 1020
- *Herbaspirillum rubrisubalbicans* 1020
- most widely studied diazotrophic bacterium 1020, 1021
- nitrogenase structural genes, *nif* 1020
- nitrogen fixation regulation 1020
- nitrogen-fixing bacterium 1020
- plant growth-promoting activity 1021
- *Saccharobacter nitrocapta* 1021
- saccharolytic activity 1021
- Dicer-like (DCL) proteins 277
- dicot plants 307
- dicyandiamide 71
- disordered stress proteins 473
- diversification
 - production 934
- divinyl ether synthases (DESs) 328
- DL- α -difluoromethylarginine (DFMA) 1054
- DNA damage 257, 263, 267, 270, 272
- DNA demethylation 276
- DNA hypomethylation 277
- DNA lesions
 - induced by ROS 265
- DNA ligase 271
- DNA methylation 267, 274, 275, 321, 498
 - RNA-directed 277
- DNA methyltransferases 272, 276
- DNA polymerase 257, 267, 271
- DNA repair 264, 267, 270, 271, 272, 278, 296
- DNA replication 271
- dog1* mutants 535
- doubled haploid (DHs) 787
- double-strand break (DSB) repair 263
- Δ^1 -pyrroline-5-carboxylate synthetase (P5CS) 148, 315
- DREB1A, DREB1B, and DREB1C proteins 611
- DREB2A overexpression 95
- DREB2A protein 443, 564, 565
- DREB1A*-targeted genes 571, 572
- DREB2A* transcription factor 442
- DREB2B protein 564, 565
- DREB1/CBF homologs 563
- DREB factors 595
- DREB1* gene 670
- DREB* genes in plant species
 - CBF/DREB1 genes of *Arabidopsis* and rice
 - functionally different 576, 577
 - discrepant expression in monocots and dicots 576
 - overexpression of 572
 - heterologous expression 573, 575, 576
 - in transgenic *Arabidopsis* 572, 573
- DRE-binding protein (DREB) 256, 438, 562, 563
- DREB1 proteins 564
- DREB2 proteins 564
- DREB transcription factors 609
- drought 58, 142–146, 210, 305, 326, 713, 818, 819, 917
 - ABA-dependent pathway 659–662
 - ABA-independent pathway 662, 663
 - antioxidant defense system in plants under 232
 - enzymatic components 234, 236
 - non-enzymatic components 233, 234
 - ascorbate peroxidase (APX) 917
 - chaperones 917
 - drought-downregulated genes 917
 - drought-modulated expressed sequence tags (ESTs) 916
 - drought-responsive root transcripts 917
 - drought-tolerant clones 916
 - drought-tolerant plants 819
 - early signaling events 658, 659
 - genes conferring tolerance 67
 - major tea-growing areas 916
 - molecular response 916, 917
 - and oxidative stress 229–231
 - Palmer drought severity index (PDSI) 819
 - indication 819
 - from perception to gene expression 657, 658
 - plant transcriptomes 916
 - polyethylene glycol (PEG) 917
 - possible effects in plants 211
 - *PR-5* gene 917
 - proteomic approach 917
 - desiccation-responsive proteome 917
 - reasons for 818, 819
 - resistance of ICC4958 335
 - sensitive variety 336
 - signaling 82
 - strategies, tolerate drought stress 819
 - stress induce ABA accumulation in plants 83
 - stress-responsive mechanisms 916
 - tolerance (*see* drought tolerance)
 - UPASI tea research foundation 916
 - water-deficit-induced genes 916
- drought and salinity
 - environmental changes 657
 - gene expression 657, 658
 - greenhouse gases 657
 - cellular homeostasis 657
 - global warming 657

- ion and osmotic homeostasis imbalance 657
 - roots 658
 - stomatal closure 657
 - plant families 644, 645
 - Arabidopsis 644
 - *Caenorhabditis elegans* 644
 - DNA-binding domains 645
 - *Drosophila melanogaster* 644
 - *Saccharomyces cerevisiae* 644
 - drought tolerance 81, 440–443, 823, 824
 - crop performance analysis 823
 - definition 823
 - polyamines 823
 - biosynthetic pathway 823
 - resurrection plants 823
 - transgenic approaches 823
 - transgenic rice plants 823
 - *adc* (arginine decarboxylase) gene 823
 - water stress proteins 824
 - aldose reductase 824
 - induction pattern 824
 - late embryogenesis abundant (LEA) dehydrin 824
 - weed 823
 - molecular dissection 823
 - *Oropetium thomaeum* 823, 824
 - *Physalis minima* 823
 - dry habitats plant resource, aloe 860–863
 - aloe-emodin, interferon-inducing agent 862
 - aloe products use/utilities 862
 - aloes 861
 - characteristics, aloe 861
 - chemical ingredients 862
 - classification 862
 - critical role, plants 860
 - family Xanthorrhoeaceae, aloe 861
 - field-based solutions 860
 - gel 862
 - global warming, to ameliorate 861
 - medical use, aloe 861
 - medicinal products, aloe leaves 861
 - bitter exudates 861, 862
 - nurse plant 863
 - Socotra aloe, commercial potential 861
 - therapeutic use, aloe species 861
 - world-wide use 862
 - dsm1* mutants 424
 - dust 251
- e**
- early signaling events
 - drought and high salinity stresses 658, 659
 - mutant yeast 658
 - histidine kinase 658
 - osmosensor proteins, SLN1 and SHO1 658
 - phytohormone ABA 659
 - plasma membrane 658
 - osmosensor protein, ATHK1/AHK1 658
 - osmotic stress 658
 - ecosystem 347
 - productivity 110
 - response, climate change 904
 - services 387, 859
 - constraints 859
 - fertilizers 859
 - human-induced climate changes 859
 - world's major challenges 859
 - World Water Day 859
 - effector-triggered immunity (ETI) 326
 - EF-hand motif-containing Ca^{2+} -binding proteins 269
 - Elaeis guineensis* 357
 - electric conductivity 194, 686
 - electric power 62
 - electrolyte leakage rate (ELR) 194
 - electron transport 29
 - electrostatic forces 686
 - E3 ligases 516
 - endopeptidase 140
 - endophytes 1041, 1042
 - characterizing fungal endophytic classes 1042
 - class 1 clavicipitaceous endophytes 1042
 - class 2 endophytes 1042
 - class 3 endophytes 1042
 - class 4 endophytes 1042
 - symbiotic criteria 1042
 - definition 1041
 - DNA based detection technology 1041
 - microbial associations with plants 1041
 - symbiosis 1041
 - energy, from biomass 363
 - energy impinging onto the soil 15
 - energy return on investment 354
 - engineered rice *OsWRKY13* 38
 - engineering plants 31
 - engineering superoxide dismutase
 - climate change 913, 914
 - drought stress 913
 - engineered thermostable 913, 914
 - implications 913, 914
 - thermostable PotSOD 913
 - transgenic plants 913
 - environmental stress factors 933
 - ERD1* gene 662

- EREB proteins 607
 - ERECTA* gene 742
 - ERF/AP2 proteins 564
 - ERF-associated amphiphilic repression (EAR) motif 610
 - ERF* gene 608
 - ERK (extracellular signal-regulated kinase) 267
 - Eruca sativa* 201
 - Escherichia coli* 693
 - espimo1* mutant 448
 - ethylene 142, 326, 328, 536, 608, 739
 - ethylene-responsive genes 332
 - ethyl methanesulfonate (EMS) 263
 - Eucalyptus camaldulensis* 368
 - Eucalyptus globulus* 368
 - Eucalyptus grandis* 368
 - Eucalyptus rudis* 369
 - Eucalyptus urophylla* 368
 - Euglena gracilis* 263
 - Euphorbiaceae 366
 - evaporation 5
 - Evergreen Revolution 830
 - excess water (flooding) tolerance 826, 827
 - alcohol dehydrogenase activity 827
 - anaerobic conditions 826
 - *Echinochloa crusgalli* 826
 - metabolic characteristics 827
 - studies 826, 827
 - ethanol production 826
 - Pasteur effect 827
 - pentose phosphate pathway 826
 - respiratory metabolism, study 827
 - seed germination, study 827
 - alcohol fermentation, anaerobic condition 827
 - submergence tolerance 827
- f**
- F-box protein COI interaction 330
 - F-box proteins 516
 - fermentation 364
 - Ferocactus acanthodes* 112, 119, 121
 - fertilizers 306, 787
 - firewood 363
 - Flacourtiaceae 366
 - flood-prone rice-growing areas 66
 - floods 63
 - Florin, DH wheat variety 788
 - flowering 950, 951
 - amino-oxyacetic acid, treatment 950
 - *Arabidopsis* plants 951
 - late-flowering phenotype 951
 - salicylic acid-deficient 951
 - aspirin, exogenous application 950
 - chelating agents 951
 - cumulative effect, salicylic acid 950
 - flower-inducing property, salicylic acid 950
 - fruit yield enhancement 950
 - metal-chelating property 951
 - non-photo-inductive light cycle 950
 - phloem-transmissible factor 950
 - salicylic acid 950
 - salicylic acid, exogenous foliar spray 950
 - thermogenic plants 951
 - inflorescences 951
 - transcription factor HAHB10 951
 - flu* mutant 93
 - fodder 363
 - food consumption, seasonal reduction in 62
 - food insecurity 57
 - food production 27
 - food security 59
 - forage quality 18
 - forestry 346
 - forests, Bangladesh 883–887
 - Bangladesh forest department 883, 884
 - classification, classes 884
 - based on 884
 - key ecological attributes 884
 - topographic conditions 884
 - climate change, effects of 884
 - forest area 883
 - forests-depleting factors 884
 - negative change rate, forests 884
 - unclassified state forests (USF) 884
 - UN food and agriculture organization (FAO) 883
 - forests definition 883, 884
 - report 883
 - fossil fuels 28, 251, 346, 348, 352, 360
 - energy production 352
 - Fragaria ananassa* 19
 - freezing stress 150, 151
 - freezing tolerance 277
 - fungicides 786
 - Fusarium oxysporum* 43
- g**
- galactinol 94
 - Ganges 62
 - Garden Orache 42
 - GDP 62
 - gene discovery
 - approaches, forward and reverse genetics 663

- mutant 663
- approaches, crop improvement 664
- *Escherichia coli* 664
- gene selection 663
- genomes 663
- transgenic model organisms 665
- stress responses 663–665
- genetic diversity, importance of 934
- genetic engineering 96, 787
- engineered plants, for overexpression of
 - antioxidant enzymes 261, 262
- improve abiotic stress tolerance
 - in crops 32–34
- intervention to build crop plants 30, 31
- plants enhanced to accumulate
 - osmolytes 41
- genetic models 373
- genetic network regulating systemic response
 - *miR399* 1000
 - biological significance 1000
 - expression level 1000
 - long-distance movement 1000
 - overexpressing plants 1000
- miRNAs 1000
- Pi accumulation 1000
- Pi homeostasis 1000
- Pi starvation 1000
- Pi-sufficient conditions 1000
- ubiquitin- conjugating E2 enzyme (UBC24) 1000
- Genetic Resources Information Network (GRIN) 348
- genetic variability
 - adverse climatic conditions effects 847
 - chickpea germplasm 847
 - large genotypic variability 848
 - screening 847
 - drought-tolerant genotypes 847
 - heat stress affected traits 848
 - heat tolerance 847, 848
 - heat tolerance index (HTI) 848
 - variations 848
 - heat tolerant, chickpea genotypes 847
 - kabuli types 847
 - heat-tolerant line ICCV 92944 848
 - pod-filling period 848
- genome–environment interactions 791
- genome hypermethylation 274
- genome protection
 - genes and proteins, induced by abiotic/genotoxic stress conferring 265, 266
- genomics 326
- genomics, bacterial endophytes 1029
- endophytic bacterial genomes
 - sequencing 1029
- genes encoding type 1029
- GENOPAR consortium of Parana state 1029
- quorum sensing system 1029
- genotoxic adaptation
 - role of DNA damage response in 269–272
 - role of MAPK, and calcium signaling 267–269
- genotoxic stress 260
 - adaptive response to 260, 263–265
 - signal transduction from abiotic stress
 - perception to gene activation 273
 - tolerance, epigenetics of 272–274
- germplasm 18
 - collection 402
 - and genetic resource databases 348
- GHG balances of bioenergy crops 353
- GHG emissions 353, 356, 365, 367, 387, 392, 397, 399, 401
 - and loss of biodiversity 380
 - released during plant growth 363
- GHG mitigation
 - intermittent drying 72
 - options potential and constraints 72
 - site-specific N management 72
- GHG net GHG exchange 379
- gibberellic acid 713
- gibberellic acid insensitive (GAI) 665
- gibberellins 332, 427, 487, 536, 739
- global average mean water vapor 5
- global climate changes 305, 521
 - mitigation 363
- global dimming 5
- global ecological crisis 903
 - climate change 903
- global hydrological cycle 5, 858
 - under pressure from 858
- global temperature 252
- global warming 30, 58, 62, 109, 252, 346, 396, 507, 903
 - effects on roots and plant–soil links 129
 - potentials 353, 357
 - of CH₄ and N₂O under alternate wetting and drying 70
- glucose-6-phosphate dehydrogenase 140
- glutathione 255, 330
- glutathione peroxidase (GPX) 177, 255, 709
- glutathione reductase 235, 255, 330
- glycerophosphodiesterase-like protein (NtGPDL) 267
- glycine betaine 42
- Glycine soja* 88

- glycophytes 687
- glycosylation
 - of NHX1 protein 693
- GmERF3 transcription factor 608
- Golden Rice 787
- Gossypium hirsutum* 465
- G-protein 447
- grain production 4
- greenhouse gases (GHGs) 4, 58, 109, 251, 346, 507, 818
 - elevated temperatures due to 29
 - emissions 58
- Green Revolution 28, 57, 65, 306, 320, 786
- ground water 63
 - depleted 63
- growth
 - *Brassica juncea* 945
 - bud formation inhibition 946
 - growth-promoting responses 945
 - high-salinity stress 946
 - effects 946
 - presowing seedsoaking 945
 - salicylic acid 945, 946
 - effects 946
 - exogenous application 946
 - inhibitory effect 945
 - proline accumulation regulation 946
 - spray 946
- growth-inhibitory effects 327
- growth rate 947, 948
 - *Arabidopsis NahG* transgenic plants 947
 - tumor-like growth 948
 - at lower temperatures 947
 - wild-type plants 947
- guaiacol peroxidase 714, 966
- guard mother cell (GMC) 89
- GUS signals 617
- Gypsophila oblongeolata* 202
- h**
- HadCM3 global climate model 252
- HaHB1* gene 671
- Halliwell–Asada pathway 255
- halophytes 190, 348, 368, 369, 686, 687
 - expressed sequence tag (EST) databases 687
 - salt-tolerance mechanism 687
 - tissue-specific adaptation 687
 - salt exclusion 689
 - salt-saturated organs 689
 - salt-secreting hairs 689
 - salt secretion by bladder cells 688, 689
 - salt secretion by salt glands 688
 - succulence 687, 688
- Haloxylon salicornicum* 686
- Hal2p protein 466, 467
- hard inheritance 274
- HARDY (HRD) gene 608
- hardy* (HRD) gene 595
- HD family
 - eukaryotic lineages 653, 654
 - HD-ZIP I subfamily 654
 - helixes 653
 - *Knotted-1* gene 653
 - *Medicago truncatula* 654
 - SELEX (systematic evolution of ligands by exponential enrichment) 654
 - whole-genome sequences 653
 - yeasts 654
- HD-ZIP proteins 556, 561
- heat/drought combination 29
- heat resistance 277, 278
- heat sensitivity, of labile-enzymes 29
- heat shock factors 445
- heat shock proteins (HSPs) 31, 91, 429–431, 439, 470, 471, 710, 822
- heat stress 109, 148–150
 - chronic vs. acute heat stress 121, 122
 - conceptual model of observed/predicted effects 130
 - direct vs. indirect effects on roots and shoots 122, 123
 - effects on nutrient relations 123–125
 - effects on root respiration and carbon metabolism 125, 126
 - effects (excluding nutrients), on roots 112–115
 - root growth 116, 119
 - effects, on roots, nutrient-related 117, 118
 - effects on root water relations 126
 - interactions with other global environmental-change factors on roots 126–128
 - on plants, direct vs. indirect effect 116
 - and root-soil interactions 128, 129
 - roots vs. shoots, effects of 119
 - shoot and root vs. root-only vs. shoot-only heating 119–121
- heat stress, chickpea 842, 843
 - adverse effects on 843
 - heat shock conditions 843
 - high-temperature conditions 843
 - earlier phenology 843
 - high-temperature tolerance 842
 - morphological processes, impact on 842
 - photoperiod 843
 - physiological processes, impact on 842
 - plant phenology, modification 842

- plant–water relationship 842
- pod development, effects on 843
- pollen germination 843
- related symptoms 842
- sensitive organs 843
- heat tolerance 445–447
 - defined 431
- heat tolerance, screening techniques 844–846
 - cell membrane thermostability (CMT) 845
 - correlated with 845
 - cellular membrane systems 845
 - electrolyte leakage 845
 - leaf disks 845
 - International Crops Research Institute for the Semi-Arid Tropics (ICRISAT) 844
 - long-term average temperatures 844
 - reproductive-stage heat tolerance screening 844
 - late-sown crop (ferbruary) 844
 - normal-sown crop (october) 844
 - pod set difference 844, 845
 - heat-sensitive line (ICC 4567) vs. heat-tolerant line (ICC 15614) 844, 845
 - pollen viability tests 845, 846
 - Alexander's stain 845
 - sorghum (*Sorghum bicolor* (L.) Moench), study of 845
- heat waves 4
- heavy metal
 - contamination 146, 147
 - exogenously applied NO, protection from toxicity 147
 - role of NO in stress 146
 - toxicity, SNP-induced alleviation 148
- heavy metal stress, abiotic stress 268, 965, 966
 - aluminium tolerance 965
 - antioxidant enzymes 966
 - cadmium 965
 - hazards to plants 965
 - lead toxicity 966
 - lipoxxygenase activity 965
 - salicylic acid 965
 - ameliorative role 965, 966
 - exogenous application 965
 - heavy metal toxicity alleviation 965
 - seed-soaking treatment 965, 966
 - signaling response 966
- Helianthus annuus* 44, 119, 212, 263, 661
 - *Hahsp17.6G1* expression 611
- hemicelluloses 383
- hemoglobins 152
 - detoxification of excess NO 152
- herbicides 787
- heterochromatic siRNAs (hc-siRNAs) 277
- heterologous expression, of key genes 42
 - chimeric bifunctional *TPSP* 42
 - *TPP* 42
 - *TPS* 42
- heterologous systems 31
- Hibiscus moscheutos*
 - exogenous SNP application 148
 - exposed to toxic Al^{3+} levels 146
- high altitude plants, clues 914–916
 - ABA-independent pathway 916
 - gene expression 916
 - kinetics 916
 - growing season temperature 915
 - high altitude 914
 - high altitude, evident characteristic features 915
 - jasmonate signaling pathway 915
 - Kibber, natural habitat of high altitude 915
 - late embryogenesis abundance (LEA) protein 915
 - upregulation 915
 - lipoxxygenase gene 915
 - low-temperature-inducible *QM* gene 915
 - low-temperature-responsive genes 915
 - mRNA-mediated gene expression analysis 915
 - photosynthetic acclimatory response
 - low-temperature-mediated 916
- hill forests 886
 - biological diversity 886
 - dominant trees 886
 - ecosystem importance 886
 - map 885
 - species distribution 886
 - total area 883
 - types 886
 - USFs 886
- Himalaya, mountain ranges 904, 906, 907
 - annual precipitation 907
 - biological diversity 905
 - Cenozoic era 904
 - classification 905
 - ecoregions 905
 - flower diversity 906
 - India 907
 - glaciers 904
 - land cover 904
 - major ecosystem services 906
 - major zones, India 906
 - central Himalaya 906
 - eastern Himalaya 906
 - western Himalaya 906, 907

- map 905
- physical diversity 905
- rainfall pattern, summer monsoon 907
- restricted migratory routes 906
- as sink, carbon dioxide (CO₂) 906
- snow cover 904
- sustainable development 906
- topographical features variations 906
- various ecosystem services 906
- water tower of Asia 904
- histidine 469
- histidine kinase 82
- homeostasis 620
- homologous recombination (HR) 263
- homologous systems 31
- Hordeum vulgare* 263, 608, 610
- salt tolerance in 731
- hormonal regulation, of root architecture 315–317
- hormones 326
- hos9* and *hos10* mutants 596
- 9-HPOD hydroperoxy fatty acids 337
- HsfA3* factor 611
- HT1* gene 86
- ht1* mutants 86
- human hCyp33 protein 469
- humanity 391, 403
- humidity 305, 380
- HvRAF* gene 608
- hydrogen (H) 349
- hydrogen peroxide (H₂O₂) 190, 209, 210, 317, 327, 431, 739
- hydrogen sulfide (H₂S) 142
- hydroperoxide lyases (HPLs) 328
- hydroperoxy fatty acids 336
- 12-hydroxy-JA (12-OH-JA) 330
- hydroxyl radicals 210
- hydroxyurea 271
- HY5* gene 298
- hyl1* gene 487
- hypermethylation 277
- hyperoxia 271
- hypothetical model depicting, general abiotic stress response in plants 439
- hypoxia, in plants 151, 608
- NO levels 151

- i**
- IAA. *see* indole acetic acid (IAA)
- ICE1 protein 595
- Indo-Gangetic plain 65
- agro-ecological regions 65
- potential adaptation strategies 65
- rainfall 65
- temperature 65
- indole acetic acid (IAA) 332, 426, 427, 592
- mediated signaling pathways 147
- induced systemic tolerance (IST) 426
- induced systemic tolerance, to salt and drought 332
- industrialization 73
- information and communication technology-based agro-advisories 75
- Digital Green 75
- IFFCO Kisan Sanchar Limited (IKSL) 75
- inositol 31
- insecticides 786
- interannual variations, in crop yields 6
- Intergovernmental Panel on Climate Change (IPCC) 252
- special report on emissions scenarios 252
- intergovernmental panel on climate change (IPCC) 816
- international board for plant genetic resources (IBPGR) 934
- International Crops Research Institute for the Semi-Arid Tropics (ICRISAT) 844
- International Rice Research Institute (IRRI) 66
- studies on GHG emissions on flooded rice reveal 70
- intersimple sequence repeat (ISSR) marker 867
- intracellular ion homeostasis 686
- intrinsically unstructured proteins 473
- in vitro* cultures establishment 936
- arid environment plants 936
- juvenile explants 936
- cotyledonary nodes, seeds 936
- mature explants 936
- nodal shoot segment 936
- *in vitro* mass multiplication 936
- in vitro* produced shoots rooting 871
- auxin-treated shoots 871
- culture composition 871
- *ex vitro* rooting 871
- ion toxicity 326
- ion transporters 690
- plasma membrane transporters 690
- plasma membrane H⁺-ATPase 692
- SOS1 690, 691
- vacuolar transporters 692
- H⁺-PPase (V-PPase) 695–697
- NHX1 692–694
- vacuolar H⁺-ATPase 694, 695
- IPCC Fourth Assessment Report 59

Ipomoea campanulata 508
irrigation 62

j

Jacquemontia pentantha 508
JAR amino acid synthetase 330
jasmonate
– pathway 336
jasmonate-ABA balance 91
jasmonate biosynthesis pathway 328–330
jasmonates 327, 711
jasmonate ZIM domain (JAZ) repressors 330
jasmonic acid 142, 327, 330, 336, 608
– in abiotic stress response 333
– biosynthesis pathway 329
– signal transduction in roots 333, 334
jasmonic acid signaling 960, 961
– *Arabidopsis* 961
– – ESR gene, dual function 961
– jasmonic acid-inducible protein 960
– – epithiospecifier senescence regulator (ESR) 960
– senescence regulation 960, 961
– transcription factor WRKY53 960
– – expression 960
– – salicylic acid-inducible 961
– – suppression 961
jasmonic acid signaling, regulation 336, 338
jasmonic acid synthesis 333
jasmonoyl derivatives 330
Jatropha 379, 396
Jatropha curcas 368
Jatropha curcas 366, 709
JAZ proteins 330

k

K⁺ channels 86
ketoacyl-CoA thiolase (KAT2) 329
Kosteletzkya virginica 140
Krishi Vigyan Kendras 68
K⁺ transport systems 464

l

Lactuca sativa 465
land management 346, 381
late embryogenesis abundant proteins (LEA) 31
latent heat 6
– of vaporization 12
lateral roots formation 999
– auxin perception complex 999
– – transport inhibitor response 1 (TIR1) 999
– auxin role 999

– Pi deficiency 999
Lauraceae 366
leaf area index (LAI) 13
leaf relative water content 14
leaf size 110
leaf temperature 12, 14
leaf transpiration 12
leaf turgor 190
leaf water potential 14
LEA proteins 39, 423, 591
LEA (late embryogenesis abundant) proteins 710
Legume–rhizobia symbiosis 331
Leguminosae 366
Lens culinaris 201
Leptochloa fusca 369
lignification 96
lignin 368
lignocellulose 348, 364
lignocelluloses 360
Limonium bicolor 611
Limonium latifolium 688
Limonium sinense 687
linkage-based phenomenon 31
linolenic acid (9-HPOT) 333
linolenic acid (18 : 3) 328
lipid epoxides 202
lipid peroxidation 190
lipoxygenases (LOXs) 328
livestock
– adversely impacted by climate change 62
– as emitters of CH₄ 71
Lolium temulentum 432
low altitude plants, clues 916–919
– molecular response, drought 916, 917
(See also drought)
– molecular response, temperature 917–919
(See also temperature)
low-oxygen environmental conditions 151
– plant tolerance to 152
low temperature (cold stress) 150, 151
LOX sequences 334
LRR (leucine-rich repeat) proteins 267
Lupinus albus 378

m

macrophyta adaptation
– salicylic acid 971
Macrotyloma uniflorum 687
maize domestication
– apical dominance 654, 655
– teosinte branched 1 (tb1) gene 654
– – alleles 655

- axillary meristems, branch elongation 655
- genes, cell cycle 654
- SBP (squamosa promoter-binding protein) 655
- TCP (TB1, cycloidea, PCF-domain protein) 654
- teosinte 655
- *teosinte glume architecture 1 (tga1)* gene 655
- malondialdehyde 194, 468
- Malus baccata* 610
- Malus hupehensis* 230, 233, 236
- Malus prunifolia* 230, 233, 236
- Malus pumila* 19
- Malvaceae 366
- mangrove forests 884, 885
- Bay of Bengal 884
- map 885
- dominant trees 884
- Royal Bengal Tiger 884
- Sundarbans 884
- ecosystem 884, 885
- reserve forests 884
- species distribution 885
- zones 884
- total area 883, 884
- mangroves 191
- manuring 71
- MAPK cascade 267
- MAPK cascades 272
- MAPK, in genotoxic adaptation 267
- MAPKKK-MAPKK-MAPK module 267
- MdSIMYB1* gene 44
- mechanical tillage 16
- Medicago sativa* 365, 608
- Medicago truncatula* 88, 144, 146, 147, 231, 314, 318, 333, 335, 488, 491, 513, 514, 608, 613, 654, 743, 916
- *miR169* downregulated in 514
- model legume, transcriptome analysis 335
- transient downregulation of *miR169* and *miR166* 513
- upregulation in *miR398* 514
- mediterranean climate 5
- melatonin 713
- membrane transport systems 591
- meristemoid mother cell (MMC) 89
- Mesembryanthemum crystallinum* 445, 686
- mesophyll conductance 12
- Mesorhizobium ciceri* 335
- metabolic activity 686
- metabolic adaptations, to environmental stresses in plants 707
- metabolic networks 948, 949
- apical dominance 949
- *Arabidopsis* plants 949
- *cpr* mutants 949
- transcriptomic analysis 949
- auxin-deficient/auxin-insensitive mutants 949
- auxin indole acetic acid (IAA) 949
- endogenous levels 949
- auxin-overproducing mutant yucca 949
- cell division rate, increased 949
- gibberellin pathway 949
- photosynthetic efficiency 949
- salicylic acid-accumulating mutants 949
- salicylic acid and auxin signaling 948, 949
- salicylic acid, high doses 949
- transcription factors *OBP1* 949
- salicylic acid-inducible *DOF* 949
- wild-type plants 949
- auxin sensitivity 949
- exogenous treatment, salicylic acid 949
- metabolic pathway 940, 941
- salicylic acid 940, 941
- metabolite, changes during abiotic stresses 729
- carbohydrates 733–735
- glycine betaine 729–733
- biosynthetic pathway 732
- polyamines 735, 736
- proline 729–733
- model proposed for 730
- multiple functions of 732
- metabolite modulation, stress tolerance 770–774
- ABA metabolic pathways 773
- abiotic stress-resistant transgenic plants 771
- abiotic stress-tolerant cultivars 770
- adaptation process 770
- antioxidant system 773
- *Arabidopsis* 772
- heat-tolerant transgenic 772
- betaine aldehyde dehydrogenase (BADH) gene 771
- transgenic bread wheat 771
- biotechnology advancement 770
- choline monooxygenase (CMO) gene 772
- transgenic cotton 772
- classical breeding programmes 770
- major limitations 770
- stress-tolerant plants 770
- compatible solutes 770
- accumulation 770
- biosynthesis 770
- *de novo* synthesis, choline 772

- glycine betaine synthesis 772
- $\Delta 1$ -pyrroline-5-carboxylate reductase (P5CR) gene 772
- gene pyramiding/coexpression 773
- gene(s) overexpression 770
- glycine betaine
 - accumulation constraint 772
 - overproduction 771
- heat stress-tolerant transgenic tobacco plants 773
- main abiotic stresses 770
- mannitol content, enhanced 772
- modulated main metabolites, plant stress response 771
- osmoprotectants 771
- phytochelatin synthase (PCS) 774
 - genes encoding 774
 - heavy metal hyperaccumulators 774
 - heavy metal response 774
 - polyamine levels 771
 - modification 771
- resurrection plants 772
- salinity tolerance, enhanced 773
- stress-resistant transgenic plants 770
- transgenic chickpea 772
 - P5CSF129A expression 772
- transgenic technology 770
- trehalose gene overexpression 773
- wild-type plants 772
- yeast TPS1, transgenic tomato 772, 773
- metabolome 439
 - analysis 94, 95
- metabolomics 728, 729, 755, 756, 757
 - comprehensive plant metabolic profile 756
 - different study approaches 756, 757
 - limitation 756
 - metabolites
 - ever-broadening range 756
 - structural diversity 756
 - plant metabolomics research, approaches 756, 757
 - gas chromatography (GC) 756
 - liquid chromatography (LC) 756
 - mass spectrometry (MS) 757
 - nuclear magnetic resonance (NMR) 757
 - vs. proteomics 756
 - vs. transcriptomics 756
- metagenomics, rhizospheric microbes 1029–1031
 - culture-independent analysis 1030
 - culture-independent methodologies 1030
 - culture-independent PCR-based amplification 1030
 - DNA-based methods 1030
 - DNA sequencing 1030
 - endophytic microbial communities 1030
 - environmental microorganisms 1029
 - genomic analysis 1030
 - nitrogen fixation
 - DNA-based analysis 1030
 - mRNA-based analysis 1030
 - *nifH* genes 1030
 - potential 1030
 - rice endophytic community 1030
 - ubiquitous microorganisms 1029
- metal toxicity 210
- methane (CH₄) 251, 252, 299, 507
 - emissions 59, 69
- 3-methyl-adenine DNA glycosylase (AAG) 265
- methyl glyoxal bis-guanyldrazone (MGBG) 1054
- methyl jasmonate 31
- methyl-jasmonate (Me-JA) 330
- methyl methane sulfonate (MMS) 271
- micro aggregation 71
- microalgae function 368
- microalgal bioenergy systems 353
- microorganisms
 - role in plant drought stress tolerance 426, 427
- microRNAs (miRNAs) 277, 336, 337, 432, 482, 522
 - in abiotic stress, role of 742, 743
 - artificial 494
 - biogenesis 483–485
 - and function, in plants 509–511
 - differentially regulated, in response to climate change-associated stresses 512
 - downregulation of *miR398* under ozone and UV-B stress 515
 - drought-responsive, identified in plants species 513
 - heat stress, to alter expression in 511
 - miRNA responses during ozone and UV-B stress 515
 - and nutrient homeostasis 490–492
 - in response
 - to cold and heat stress 488, 489
 - to drought stress 487, 488
 - to hypoxia and oxidative stress 489, 490
 - to salt stress 488
 - to UV-B radiation 490
 - role in plant stress responses for adapting to climate change 486
 - stress-miRNA networks for adapting to climate change 494–497

- Millennium Development Goal (MDG) 1 786
 - miR159* expression 487
 - miR156*, overexpression 513
 - miR319* signaling molecule 337, 338
 - Miscanthus* (*Miscanthus* × *giganteus*) 365, 366
 - mismatch repair (MMR) 263
 - mitigation 69
 - CH₄ emission from rice cultivation 70
 - strategies 70, 71
 - of CO₂ emissions 71
 - mitogen-activated protein (MAP) 438
 - mitogen-activated protein kinases (MAPKs) 87, 142, 257, 424, 591, 741
 - molecular biomarkers 44
 - molecular chaperones 591
 - molecular genetics 326
 - molecular reprogramming events, mitigate environmental constraints 590–592
 - monocot plants 307
 - monodehydroascorbate (MDHA) 230, 234
 - monodehydroascorbate reductase (MDHAR) 255
 - monosaccharides 94
 - Moraceae 366
 - mountains ecosystem 904
 - altitudinal dimensions 904
 - environmental variations influence 904
 - genetic diversity 904
 - Himalaya, mountain ranges 904
 - land cover 904
 - valley exit jet 904
 - weather-associated specific niches 904
 - mtlD* gene 42
 - MtNAC969, NAC transcription factor 314
 - mulching 71
 - multifunctional bioenergy agroecosystems 387
 - multigene phenomenon 31
 - mutations
 - in epigenetic regulator *SWP1* 321
 - in *OsWOX11* 307
 - MYB family
 - biotic, abiotic stress, example, transcription factors 646–648
 - classified, subfamilies 645
 - DNA binding and protein interaction 645–648
 - MYB proteins 645
 - MYB1* gene 44
 - MYB33* gene 497
 - MYB (myeloblastosis)-like proteins 556–558
 - MYB-like transcription factor (StMYB1R-1) 620
 - MYB transcription factors 327
 - and cold stress 618, 619
 - and salinity tolerance 619–621
 - MYC (myelocytomatosis)-like basic helix–loop–helix (bHLH) proteins 556
 - MYC-like bHLH proteins 558–560
 - vs. MYB proteins 560, 561
 - MYC/MYB transcription factors 593, 594, 615–618
 - MYC2-regulated gene transcription 620
 - myelocytomatosis oncogene 592, 607
 - Myraceae 366
- n**
- Na⁺ absorption ratio 686
 - NAC and ZF-HD proteins 595, 596
 - NAC family
 - C-terminal domain 650
 - DNA-binding domain 650
 - five subdomains (A–E) 650
 - plant-specific transcription factor families 650, 651
 - NaCl-induced genotoxic stress 276
 - NaCl stress 620
 - NAC transcription factors 611–614
 - NAD⁺ breakdown 272
 - NADPH oxidase 297
 - Na⁺/H⁺ antiporter 689, 692, 693
 - NahG* gene 738
 - Na⁺/K⁺ ratio 140, 141
 - NAM, ATAF, and CUC (NAC) 424
 - National Action Plan on Climate Change, 2008 73
 - National Agricultural Risk Fund 75
 - National Commission on Farmers (2006) 75
 - National Initiative on Climate Resilient Agriculture (NICRA) project 67–69
 - action research implementation 67, 68
 - climate change mitigation options 69
 - partnership 69
 - technology assessment modules 68, 69
 - National Mission for Sustainable Agriculture (NMSA), 2010 73, 74
 - National Mission on Micro Irrigation 73
 - natural products, endophytic bacteria 1025, 1026
 - common soil bacterial genera members 1025
 - secondary metabolic products 1026
 - cultured microorganisms range 1026
 - diazotrophic endophyte 1026
 - fungal-based production 1026
 - antimicrobial products 1026

- low-molecular-weight compounds 1026
- microbial bioplastics 1026
- natural products list 1026
- poly-3-hydroxyalkanoate (PHA) 1026
- poly-3-hydroxybutyrate (PHB) 1026
- NCED6* expression 533
- NCED3* genes 83, 423
- NCED3* mutant 95
- neem-coated urea 71
- net new energy, produced by bioenergy crops 354
- N fertilizers 352, 353
- N fixation 128, 129
- NHX1* gene 693
- NHX1* transmembrane domain 693
- NICRA programme 76
- nitraphyrin 71
- nitrate 251
- nitric oxide (NO) 86, 137, 138, 707, 711
 - hemoglobin/NO cycle 152
 - postharvest NO fumigation 151
 - protection against heavy metal toxicity 147
 - protection against oxidative damage caused by ROS 151
 - scavenger 152
 - signaling regulation 140
- nitric oxide (NO)-mediated cysteine nitrosylation 327
- nitrification inhibitors 71
- nitrogen (N) 14, 191, 346
 - assimilation 30
 - as critical agricultural crop nutrient 18
 - cycles 350, 351
 - fertilization 18
 - status in plants and grain quality, interaction between 18
- nitrogen dioxide 138
- nitrogen fixation 957, 958, 1019
 - associative nitrogen fixation
 - limitation 1019
 - BNF 1019
 - diazotrophic bacterial endophytes 1019
 - lives in 1019
 - exogenous application, salicylic acid 957, 958
 - free-living
 - vs. endophytic bacteria 1019
 - metabolism 958
 - nitrate/nitrogen metabolism 958
 - nitrate reductase activity 958
 - enhanced 958
 - lead-induced decline 958
 - nodulation inhibition 957
 - nodule development 958
 - non-symbiotic nitrogen-fixing forms 1019
 - limitation 1019
 - significance 1019
 - obligate symbionts, leguminous plants 1019
 - *Rhizobia* 958
 - rhizobial strain 957
 - *Rhizobium*, legume symbiosis 958
 - endogenous salicylic acid 958
 - exogenous salicylic acid 958
 - NOD factors 958
 - *Rhizobia*, incompatible strain 958
 - salicylic acid 958
 - endogenous synthesis 958
 - foliar application 958
 - inhibitory effect 958
 - symbiotic nitrogen-fixing forms 1019
 - symbiotic relationship 958
 - tissue-specific metabolism 958
- nitrosative stress 138
- nitrosylation, of cysteines 328
- nitrous oxide (NO_x) 251, 252, 507
- N leaching 365, 394
- N mineralization potential 382
- nod* gene 333
- nodulation 128
- N₂O emissions 71, 353
- NO-induced osmotic stress-activated protein kinase 142
- NO₃ leakage 353
- NO-mediated detoxification 146
- non-legumes 1022–1025
 - Actinobacteria, representative 1024
 - BNF benefits 1022
 - cereals 1022
 - diazotrophic bacterial endophytes 1022
 - endophytic bacteria 1024
 - endophytic bacteria, diverse array 1024
 - enterococcal populations screening 1025
 - grasses 1022, 1024
 - bacterial diversity, nitrogen fixation ability 1024
 - fast-growing C4 grass 1024
 - *Microbacterium* 1024
 - nitrogen-fixing bacteria 1024
 - plant growth-promoting bacteria 1024
 - maize/sorghum 1023
 - nitrogen-fixing bacteria 1023
 - non-mycorrhizal fungal endophytes 1023
 - novel bacterium 1025
 - *Pennisetum glaucum* 1023
 - root-associated bacteria 1023
 - rice (*Oryza sativa*) 1022, 1023

- endophytic bacteria kinds 1022, 1023
 - methanogens 1023
 - rice paddy soil 1023
 - sugarcane (*Saccharum officinalis*) 1022
 - inoculation 1022
 - main products 1022
 - most frequent endophyte, sugarcane stems 1022
 - surface-sterilized tissue 1025
 - non-nitrogen-fixing plants 18
 - non-reducing sugars 42
 - NOS-like activity 142
 - NPR1 proteins 327
 - nuclear factor (NF)-κF 296
 - nuclear factor Y (NF-Y) 424
 - nucleotide excision repair (NER) 263
 - N-use efficiency (NUE) 348, 370, 372–374, 377, 389, 390, 394, 395
 - nutrient imbalance 326
 - nutrient management 18
 - Nypa fruticans* 369
- O**
- oil palm-based biofuel agroecosystems 366
 - oligosaccharides 94
 - omics 728
 - OPR gene expression 336
 - Opuntia ficus-indica* 367
 - organic acids 190
 - organic carbon 251
 - organic carbon storage, sustainability 896–899
 - community forestry 898
 - forest types, managed by bureaucrats 898
 - government-initiated co-management approach 898
 - human-induced removal of woody biomass 898
 - hypothetical policy decisions 899
 - local communities, active participation 897, 898
 - local community socio-economic development 898
 - long-term support, communities 898
 - national forest policy 1994, Bangladesh 896, 897
 - plan/forest management plan 898
 - project-oriented annual development programs 898
 - REDD+ 899
 - sustainable forest management 896
 - carbon sequestration objectives 897
 - sustainable organic carbon storage, ways to achieve 897
 - top-down approach, forests management 897
 - unsustainable forests management 896
 - common causes 896
 - organic carbon storage trends, different forest types 889–892
 - biomass
 - closed forests 891
 - open forests 891
 - carbon density 891
 - closed forests 891
 - open forests 891
 - carbon pool, hill forests 890
 - larger-diameter trees 890
 - living biomass 889
 - carbon stock trend 891
 - living biomass removal 892
 - factor 892
 - most degraded forests, sal forests 889
 - deforestation reasons 890
 - non-living biomass 889
 - organic carbon sequestration 892
 - smaller-diameter trees, mangrove forest 890
 - standing volume of wood, different forest types 890
 - total organic carbon storage 891
 - village forests, highest organic carbon storage 891
 - organic matter 15, 18, 305
 - Oryza sativa* 299, 607
 - CBF1/DREB1B, introduced 575
 - miR398 743
 - *phs* mutants 737
 - plasma membrane fractions 692
 - up- and downregulation of miR896 513, 514
 - OsABA2* gene 609
 - OsZIP23* gene 593
 - OsDREB2B* expression 670
 - OsDREB1F* gene 610
 - Os03g0810800* gene 609
 - osmolytes 31, 42, 44, 94, 308
 - osmoprotectant molecules 423
 - osmoprotectants 40, 42, 277, 334, 439, 728
 - osmoregulation 277
 - osmotic adjustment 1047
 - acclimatization response 1047
 - osmolyte concentration 1047
 - pattern 1047
 - osmotic potential 1047
 - determination 1047
 - symptoms 1047
 - osmotic disequilibrium 326
 - osmotic potential 190

- osmotic stress responses 277, 1056–1065
 - alleviate osmotic stress, exogenously applied polyamines role 1058–1061
 - *Ctenanthe setosa* leaves 1061
 - exogenously supplied ABA 1061
 - exogenous polyamines 1058
 - photoadaptation 1061
 - polyamine biosynthesis 1061
 - polyamines inhibitors 1058
 - salinity stress, alga 1061
 - salt-tolerant Nonabokra 1059
 - scavenging system enzymes 1059
 - stress-related effects, mitigation 1058, 1059
 - waterlogging stress 1059
 - water stress tolerance 1059
- hypo/hyper-osmotic stress tolerance, plant response 1056–1058
- ADC activity 1056
- *Arabidopsis* 1058
- defense genes 1058
- drought-sensitive wheat “Yangmai 9” 1056
- drought stress (NaCl or sorbitol)-induced damage 1057
- drought-tolerant “Yumai 18” 1056
- ionic toxicity 1057
- moderate drought stress 1056
- more-tolerant Sahara cultivar 1057
- NaCl stress 1057
- osmotica 1056
- osmotic stress adaptation 1057
- osmotic stress-induced modulation, plant polyamines 1056
- polyamine biosynthesis 1057
- polyamine biosynthetic genes 1057
- polyamine levels alteration 1058
- polyamines different compositions 1056
- polyamines functions 1058
- polyamine signaling in plants, osmotic stress 1059
- Put aminopropyltransferase (PAPT) 1056
- reactive oxygen species (ROS)-scavenging activity 1057
- salt stress tolerance, polyamines importance 1058
- salt-tolerant hybrid pumpkin 1057
- salt-tolerant rice cultivars Nonabokra 1057
- Spd 1056, 1057
- Spm 1056, 1057
- transgenics expressing polyamines homo/hetero-logously 1060
- polyamine-mediated plant osmotic stress signal transduction 1062–1065
 - ABA-dependent transcription factor, OsBZ8 1063
 - ABA-dependent upregulation, ADC2 1063
 - ABA-mediated osmotic stress response 1064
 - ABA-mediated stress signaling 1063
 - abscisic acid (ABA) 1063
 - cross-talk 1062
 - ethylene signaling pathway 1062, 1063
 - functional cooperation between ABA, NO, and ROS 1064
 - iso-osmotic stress, NaCl 1065
 - NaCl-dependent regulation 1063
 - non-ABA-mediated osmotic stress response 1064
 - PAO 1062
 - polyamine biosynthetic pathways 1065
 - polyamine catabolism 1064
 - Put accumulation 1063
 - reactive oxygen species (ROS) induction 1062
 - reactive oxygen species (ROS) role 1062
 - salinity-stressed plant 1062
 - salinity stress, Spd/ABA, kinase interaction 1063
 - salt excretion process 1064
 - salt sensitivity 1062
 - steady-state ADC mRNA, transient upregulation 1064
 - transgenics, plant polyamines research 1061, 1062
 - *Agrobacterium*-mediated transformation 1061
 - *Arabidopsis* ADC2 gene 1061
 - enhanced antioxidant activity 1062
 - multiple abiotic stresses 1062
 - osmotic stress-induced ADC activity 1061
 - polyamine metabolism 1061
 - Put, tissue-specific accumulation 1061
 - Spd accumulation 1062
 - Spm deficiency 1062
 - tetracycline-inducible promoter 1061
 - transgenic tobacco plants 1061
- osmotic stress signaling
 - subclass III SnRK2s, positive regulators in 526–528
- osmotin 591
- ost2-1* and *ost2-2* mutant 85
- Ostreococcus lucimarinus* 277
- Ostreococcus tauri* 277
- oxidation number 349
- oxidative enzymes 234
- oxidative stress 152, 202, 209, 326, 515

- adaptive responses 257–259
 - lignin biosynthesis 913
 - *Potentilla atrosanguinea*
 - antioxidative mechanism 912
 - clones 911, 912
 - managing, nature's way 911–913
 - Mn-superoxide dismutase (SOD) 912
 - photosynthesis rate 911
 - reactive oxygen species (ROS) 911, 912
 - systematic analysis 912
 - temperature-responsive plant processes 911
 - winter dormancy 912
 - *PotSOD* gene 913
 - benefits 913
 - recombinant SOD 913
 - crystal structure 913
 - subzero temperature 912, 913
 - superoxide dismutase (SOD) 913
 - overexpression 913
 - superoxide radicals 913
 - transgenic adaptation to 260
 - 3-oxo-2(2'-pentenyl)-cyclopentane-1-octanoic acid 329
 - 12-oxophytodienoate reductase 3 (OPR) 329
 - 12-oxophytodienoic acid (OPDA) 328
 - oxygen (O) 349
 - ozone
 - downregulation of *miR398* under ozone and UV-B stress 515
 - levels in the preindustrial era 515
 - tropospheric 515
 - ozone depletion 252, 295, 301
 - land plant evolution 301
 - ozone tolerance 829, 830
 - biotechnological techniques 829
 - advancement 829
 - comparative studies 830
 - effects of ozone 829
 - elevated ozone 829
 - exposure results 829
 - multifactorial environmental conditions 829
 - ozone-resistance genes 829
 - regulated by 829
 - tropospheric ozone 829
 - concentration 829
 - yellow nutsedge, weed 829
 - control 829
 - tomato productivity 829
 - PAMP-triggered immunity (PTI)x 326
 - Panax ginseng* 147, 511
 - heat stress, to alter miRNA expression 511
 - Panicum virgatum* 365
 - parallel pathway to CBFs 570, 571
 - RAV1 (AP2) 571
 - ZAT12 (zinc finger) 571
 - P5CS2* gene 570
 - Penman–Monteith model 6
 - Pennisetum purpureum* Schumach. 365
 - pentose phosphate pathway 826
 - peptidyl–prolyl *cis–trans* isomerase (PPIase) activity 469
 - perk4* mutant 90
 - peroxidase 14
 - peroxidation, of lipids 202
 - peroxisomes 203, 329
 - peroxynitrite 138
 - pesticides 306
 - Phalaris arundinacea* 365, 367
 - Phaseolous trichocarpa* 513
 - differential expression miRNAs 513, 514
 - Phaseolous vulgaris* 513
 - upregulation of miRNS 513
 - phenolic compounds 709
 - Phleum pratense* 367
 - phloem proteins 141
 - phosphatases 269
 - phosphate acquisition, soil solution 992, 993
 - cytoplasmic Pi 992
 - diffusion 993
 - high/low-affinity Pi transporters 993
 - H⁺/Pi cotransport system 993
 - hydrostatic pressure, cumulative action 993
 - physiological pH 993
 - Pi concentration 992
 - Pi-deficient condition 993
 - Pi different routes, xylem upload 993
 - Pi distribution 993
 - pK values 993
 - plasma membrane, hypersensitization 993
 - phosphoenolpyruvate carboxylase (PEPCase) 911
 - phospholipases 438
 - phosphorus (P) 191, 991, 992
 - essential macronutrient 991
 - fertilizers 991
 - application 991
 - orthophosphate, inorganic form 992
 - orthophosphate Pi, inorganic form
 - H₂PO₄[–] and HPO₄^{2–} 992
 - Pi-acquisition efficiency 991
 - Pi-use efficiency 991
- p**
- Paenibacillus lentimorbus* 331
 - Palmae 366

- phosphorylation 267
- photoinhibition 429
- photomorphogenesis 38
- photon conversion efficiency 368
- photooxidation, of chlorophyll 82
- photorespiration 203
- photosynthesis 13, 42, 81, 82, 92, 368, 952–956
 - chloroplast structure 955, 956
 - chloroplast volume 955
 - salicylic acid, high concentration 955
 - downregulated 30
 - enhanced 390
 - factors effecting 29
 - inhibition of 29
 - leaf 955, 956
 - anatomy, alteration 955
 - ROS detoxification 956
 - light acclimation, redox homeostasis 953, 954
 - antioxidant enzymes inhibition 953
 - *Arabidopsis* mutants 953, 954
 - *Arabidopsis* plants cultivation 954
 - disease resistance 954
 - endogenous salicylic acid levels 953
 - foliar salicylic acid levels 954
 - genes expression regulation 953
 - reactive oxygen species (ROS) 953
 - regulation 954
 - ROS signaling pathway 953
 - salicylic acid deficiency 954
 - salicylic acid-dependent programmed cell death 954
 - salicylic acid signaling pathway 953
 - wild-type plants 954
 - mechanisms, of plants 686
 - non-selective contact herbicide, paraquat 956
 - deleterious effect 956
 - photosynthetic inhibition 728
 - plant-water relations 952–956
 - rate 4, 14
 - related traits 953
 - exogenous application, salicylic acid 953
 - foliar application, salicylic acid 953
 - presowing seed-soaking treatment 953
 - ribulose-1,5-biphosphate carboxylase/oxygenase (RuBisCO) 953
 - soybean study 953
 - RuBisCO activity 955, 956
 - declined activity 956
 - effect of salicylic acid 956
 - salicylic acid
 - chlorophyll content 953
 - foliar application 952, 953
 - low dose benefits 955
 - metabolic response, plants 952
 - pigments 952, 953
 - pigments synthesis 953
 - positive effect 956
 - seed-soaking treatment 952
 - stomatal closure 954, 955
 - ABA-deficient mutant 955
 - endogenous salicylic levels 955
 - phytohormonal control 955
 - salicylic acid-deficient genotypes 955
 - salicylic acid role 955
- photosynthetically active radiation (PAR) 910
- Photosystem II complexes 30
- Phragmites communis* 141
- Physcomitrella patens* 487, 691
- physiological mechanisms
 - abscisic acid (ABA) 846
 - exogenous applications 846
 - role 846
- cellular membrane systems 846
- critical temperature, heat tolerance in chickpea 846
 - heat stress reduces 846
 - heat tolerance 846, 847
 - heat-tolerant genotypes 846
 - high night temperature 847
 - damaging effect 847
- osmoprotectants 846
 - exogenous application 846
 - oxidative injury 846
- sensitive genotypes 846
- physiological processes
 - salicylic acid cross-talk, germination 944
 - abscisic acid (ABA) 944
 - biosynthesis, salicylic acid 944
 - gibberellin-induced α -amylase (*Amy32b*) expression 944
 - gibberellins 944
 - induced germination 944
 - salicylic acid-regulated 942–945
 - seed germination 943
 - under abiotic stress 943, 944
 - negative effect on 943
 - regulation 943
 - salicylic acid impact 943
- phytohormone 422, 486
- phytohormone abscisic acid (ABA) 522.
 - See also* abscisic acid (ABA)
- ABRE-dependent gene expression in signaling 530

- ABRE functions as a major *cis*-acting element 530, 532
- AREB/ABFs master transcription factors, role in regulation 529
- bZIP transcription factors 531
- mediated gene expression, controlled by 522
- potential application of ABA signaling components
 - to improve crop productivity 537, 538
- PYR/PYL/RCAR-PP2C-ABA receptor complex 524–526
- role in control of seed dormancy 532–534
 - quantitative trait locus analyses, providing novel factors for 535, 536
- SnRK2s as central regulators in 534, 535
- role in plant adaptation, to land and environmental changes 536, 537
- role in regulation of abiotic stress resistance 522
- signaling components, in response to cellular dehydration 522–524
- subclass III SnRK2s, positive regulators in signaling 526–528
- phytohormones 307, 712, 713
- phytohormones, Pi deficiency 1001–1003
 - auxin role, Pi deficiency 1001, 1002
 - auxin-dependent pathway 1002
 - auxin-independent pathway 1002
 - auxin response factors (ARFs) 1002
 - auxin-responsive mutants 1002
 - auxin treatment effect 1002
 - exogenous application, auxin 1001
 - Pi-deficient plants 1002
 - Pi-normal plants 1002
 - Pi-starved plants, auxin effect 1002
- cytokinin, Pi deficiency 1002
 - cytokinin receptor, *CRE1* 1002
 - PSI genes 1002
 - shoot growth promotion 1002
- ethylene, Pi deficiency 1002, 1003
 - *Arabidopsis* 1003
 - effects on roots 1003
 - ethylene biosynthesis 1002
 - ethylene biosynthesis inhibitor (AVG) 1003
 - ethylene precursor (ACC) 1003
 - ethylene responsiveness, roots 1002
 - Pi-sufficient conditions 1002
- gibberellic acid, Pi deficiency 1003
 - DELLA-deficient mutant plants 1003
 - DELLA protein degradation 1003
 - exogenous gibberellic acid application 1003
 - gibberellic acid biosynthetic genes 1003
 - gibberellic acid-dependent pathway 1003
 - gibberellic acid-independent pathway 1003
 - MYB62-overexpressed phenotype 1003
 - Pi starvation 1003
 - PSR response repression 1003
 - role 1003
 - transcription factor, MYB62 1003
 - transcriptome analysis 1001
- phytoremediation 786, 1028, 1029
- benzene, toluene, ethylbenzene and xylene (BTEX) 1028
- copper-resistant strains 1028
- ever-increasing role 1028
- genetically engineered strains 1028
- inoculated test plants 1028
- nitro-aromatic compound degradation 1028
- potential role, bacterial endophytes 1029
- quantitative gene expression 1029
- xenobiotic degradation 1028
- xenobiotic degradation pathway 1028, 1029
- Pi-acquisition/use efficiency, improvement strategies 1003–1007
 - *Arabidopsis* 1004, 1007
 - genes involved, Pi starvation adaptation, RSA changes 1005, 1006
 - genetic regulators, plant responses 1004
 - high-throughput genotyping platforms 1004
 - low-Pi tolerance mechanisms 1004
 - marker-assisted backcrossing 1004
 - marker-assisted selection (MAS) 1004
 - nutrient acquisition 1007
 - Pi deficiency, plant responses 1004
 - Pi deprivation signaling 1004
 - Pi, low availability 1003
 - Pi sensing/signaling mechanisms 1004
 - comprehensive analysis 1004
 - Pi-sufficient plants 1004
 - PTXD gene 1007
 - root-to-shoot Pi transport 1004
- Picea abies* 707
- lignin biosynthetic pathway 707
- Pi-deficiency 995–1001. *See also*
 - phytohormones, Pi deficiency
 - local Pi responses 995–999
 - developmental responses 995
 - lateral roots formation 999
 - primary root growth, effect on 995–997
 - root hair proliferation 997–999
 - root system architecture (RSA) 995
 - responses 1001
 - hexokinase-dependent signaling pathway 1001

- hexokinase-independent signaling pathway 1001
- high-affinity phosphate transporters 1001
- Pi deficiency increases 1001
- Pi homeostasis 1001
- Pi-starved conditions 1001
- systemic Pi response 999–1001
- genetic network regulating systemic response, Pi starvation 1000 (*See also* genetic network regulating systemic response)
- phosphate starvation response (PSR) 999
- Pi-containing organic compounds 999
- Pi homeostasis 999
- Pinus densata* 231
- Pinus sylvestris*
 - polyphenolic antioxidant catechin 707
- Pinus tabulaeformis* 231
- Pinus yunnanensis* 231
- Piriformospora indica* 331, 332, 470
- Pi status sensing, plants 993, 994
 - external Pi level, perception 994
 - phosphite 994
 - Pi-deficiency responses 994
 - Pi sensing 993
 - Pi starvation 994
 - induced genes, PSI 994
 - local signaling 994
 - suppression 994
 - systematic signaling 994
 - Pi-utilizing proteins 994
 - putative Pi sensors 993
 - Pho81, cyclin-dependent kinase (CDK) inhibitor 993
 - PhoR, membrane-associated sensor kinase 993
 - putative signaling molecules 994
 - “split-root” design 994
 - transcriptomic studies 994
 - two-component signaling system 993, 994
- Pisum sativum*
 - Cd-, Cu-, or Zn-exposed roots 147
 - HSP18.1 471
- plain land sal forests 887
 - current situation 887
 - located in 887
 - map 885
 - main tree species 887
 - total area 883, 887
 - types 887
- plant acclimation 28
- plant adaptation, and mitigation strategies
 - heat shock proteins 429–431
 - for heat stress tolerance 427–429
 - thermal stability of cell membranes 429
- plant adaptations 28, 934
 - omic architecture 934
 - to salinity stress 686
 - stress, in response of 934
 - to thermotolerance 445, 446
- Plantago maritima* 190
- plant biomass 81, 82
 - productivity 81
- plant composition 18
- plant-derived organics 71
- plant diversity
 - conservation 934
- plant domestication/cultivation 508
- plant growth factors 326–328
- plant growth-promoting (PGP) fungi 332
- plant-growth promoting rhizobacteria (PGPR) 426
- plant growth regulation 949, 950
 - *Arabidopsis*
 - growth 949
 - MPK6 949
 - map kinase phosphatase 1 (*mkip1*) 949
 - growth defects 949
 - mitogen-activated protein kinase (MAPK) pathway 949, 950
 - cascades 950
 - plant growth homeostasis 950
 - PR gene expression 949
 - protein tyrosine phosphatase 1 (*ptp 1*) 949
 - growth defects 949
 - reactive oxygen species (ROS) 949, 950
 - signaling 950
 - salicylic acid 949, 950
 - accumulation 949
 - biosynthesis 949, 950
 - induced protein kinase (SIPK) 949
 - signaling pathway 949
- plant metabolome alterations, adverse conditions 757–770
 - abiotic stress 757
 - cellular level 757
 - environmental conditions 757
 - response 757
 - adaptation 757
 - biochemical, plants 758
 - *Arabidopsis* 757
 - complete genome sequence 757
 - drought 763–765
 - ABA biosynthesis 765
 - ABA-dependent regulatory systems 765
 - ABA-independent regulatory systems 765

- antioxidant systems 764
- carbohydrate metabolism 764
- carbohydrate-related genes 765
- 9-*cis*-epoxycarotenoid dioxygenase (NCED) 765
- dehydration-responsive element (DRE) 765
- drought-modulated different secondary metabolites 765
- drought-responsive metabolites 765
- drought stress perception 764
- fructans 764
- mannitol 764
- mechanical barrier, pathogens 764
- metabolic profiling analysis 765
- morphological structures 763
- phytohormone ABA 765
- proline accumulation 764
- proline biosynthesis 764
- stress-related genes, expression 765
- stress-tolerant organisms 764
- water-limiting environment 763
- heavy metals 768–770
- adverse affects 768
- antioxidant systems 769
- Cr-stressed rice roots 769
- cysteine-rich polypeptides (metallothioneins) 769
- metabolite content analyses 770
- metabolite profiling 769
- metal ion chelation, ubiquitous mechanism 769
- nitrogen metabolism 769
- NMR-based metabolic fingerprinting 769
- proline accumulation 769
- reactive oxygen species (ROS) generation 768
- ROS-detoxifying antioxidant defense machinery 769
- toxic effects, metabolic pathways 768
- hypoxia 767, 768
- genes downregulation 768
- genes upregulation 768
- hazardous effects 767
- hypoxia treatments 767
- metabolite accumulation changes 768
- metabolite profiling 767
- metabolome analyses 768
- oxygen availability, tolerate limitations 767
- proteins accumulation 768
- proteome analyses 768
- response analyses 767
- tolerance 767
- intergovernmental panel, of climate change 757
- light 758, 759
- alternative protective mechanisms 759
- *Arabidopsis* 758
- chlorophyll fluorescence imaging 759
- dark reaction 758
- flavonoids 759
- GC/time-of-flight-MS analysis 759
- glycolysis 759
- high-light stress 758
- isoprene-expressing wild-type plants 759
- light-harvesting antenna complexes (LHCs) 758
- light intensities, increment 758
- long-term exposure, high light intensity 758
- metabolic adjustments 758
- metabolite pools 759
- photosynthetic light reaction 758
- photosynthetic organisms 758
- photosystem II (PSII) 758
- RNA interference-mediated non-isoprene-expressing poplars 759
- time-course experiments 758
- tricarboxylic acid cycle (TCA) 759
- UV radiation 759
- metabolic pathways 758
- omics approaches 757
- salinity 766, 767
- ATP biosynthesis 766
- choline-oxidizing enzymes 767
- constraints, salt stress 766
- deleterious effect 766
- differential salt tolerance 766
- metabolite profiles 767
- metabolites 766
- metabolome analysis 766
- physiological mechanisms 766
- plant responses 766
- salt stress signaling 766
- Soil salinity 766
- transgenic plants 766
- UV detection 767
- water-deficit-treated plants 767
- temperature 760–763 (*See also* temperature)
- cold stress 761–763
- global climate change 760
- high temperature 760, 761
- transposon-tagged mutant lines 757
- plant–microbe interactions 326
- plant-microbe interactions under abiotic stress 428

- plant molecular adaptation 422
- plant productivity 15
 - stomatal responses in control of 82, 83
- plant resources, modifying 15
- plant response, climate change parameters
 - biodiversity-rich zone, Himalayan 910
 - relevant questions 910
 - Himalayan flora 908–919
 - increased temperature affects 908, 910
- plant responses, to abiotic stress 252–256
- plant response, to drought stress 211
 - gas exchange 217–219
 - germination 211–213
 - photosynthesis 219–223
 - plant growth 214–216
 - plant-water relations 216–217
 - reproductive development 223–225
 - seed formation 223–225
 - stomatal conductance 217–219
 - yield attributes, and yield 226, 228, 229
 - reduction in yield, crop species 227
- plant secondary metabolites 705, 706
- plant signaling machinery 460–465
- Plumbaginaeae 686
- Poaceae 686
- Poa pratensis* 231
- pollination 29
- pollution 398
- poly(ADP-ribose) (PAR) 271, 272
- poly(ADP-ribose) polymerase (PARP) 265, 272
- poly(ADP-ribosylation) (PARylation) 271
- polyamine-induced NO generation 142
- polyamine levels 142
- polyamine oxidase 142
- polyamines 31, 42, 44, 314, 708, 1053
 - in abiotic stress tolerance in plants 425, 426
 - biosynthetic genes 1053
 - in chilling stress 715
 - for chilling tolerance 315
 - definition 1053
 - implication in cellular events 1054
 - mode of action 1054
 - physiological function, research focus 1054
 - in plants 1054
 - spermine 1054
 - in stress-induced modulation of RSA 314
- polyamines metabolism, plants 1055, 1056
 - abscisic acid (ABA) 1055
 - desiccation tolerance 1055
 - homeostasis 1055
 - intracellular concentrations, free
 - polyamines 1055
 - conjugation 1055
 - oxidative deamination 1056
 - osmotic stress effect 1055
 - polyamine biosynthetic pathway 1055
 - transgenic plants analyses 1055
- poly(ethylene glycol) (PEG) 608
- polyunsaturated fatty acids (PUFAs) 202, 328, 336
- Pongamia pinnata* 366
- Populus euphratica* 91
- Populus przewalskii* 231
- Populus tomentosa* 511
 - heat stress, to alter miRNA expression 511
- Populus tremula*
 - miRNA families to be upregulated by UV-B radiation 490
- Populus tremuloides* 707
- Populus trichocarpa* 277, 371
- Porteresia coarctata* 689
- postgenomic era 728, 925
- post-transcriptional gene silencing (PTGS) 482
- post-translational protein modification 271
- potassium 191
- poverty 59
- PP2C phosphatases 659
- precipitation 5, 15, 252, 305
 - changes in 5
 - erratic intense 508
 - rates 507
 - variability 6
- preharvest sprouting 522
- primary root growth 995–997
 - *Arabidopsis*
 - experiments 995
 - primary root 995
 - ethyl methanesulfonate-induced *Arabidopsis pdr2* mutant 995
 - external Pi concentration sensing 995
 - Fe-mediated reactive oxygen species (ROS) production 996
 - inhibition 996
 - iron uptake 996
 - low-Pi patch 995
 - multi-copper (Cu) oxidase-encoding gene 996
 - MYB transcription factors 996
 - phosphate deficiency response 2 (PDR2) 995
 - Pi-dependent root architecture modifications 997
 - Pi, local availability 995
 - Pi-responsive genes 995
 - Pi-sensitive checkpoint 996

- quantitative trait loci (QTLs) 995
 - RNA interference (RNAi) 997
 - root apical meristem (RAM) 995
 - fine-tunes RAM activity 996
 - quiescent center (QC) identity 995
 - restoration 996
 - SCR (scarecrow) protein 996
 - split-root growth assay 995
 - Pringlea antiscorbutica* 715
 - probiotics 58, 72
 - productivity, salicylic acid 945, 946
 - “progressive nitrogen limitation”
 - hypothesis 18
 - proline 315, 569
 - 14-3-3 protein 591
 - proteinase inhibitors 591
 - protein carbonylation 202
 - protein degradation 522
 - protein disulfide isomerase (PDI) 469
 - protein kinase inhibitors 268
 - protein kinases 39, 40, 591, 740, 741
 - protein oxidation 202
 - proteins involved in metabolic regulation 465–469
 - proteins, stabilization of 469–471
 - proteins three-dimensional structures,
 - involved in abiotic stress tolerance 460
 - proteome 439
 - proteomics 729
 - Pseudomonas fluorescens* 332
 - Pseudomonas syringae* 330
 - Puccinellia tenuiflora* 691
 - putrescine 315, 735
 - pyrabactin-1-like proteins (PYLs) 423
 - pyrabactin resistance 1 (PYR1) 423
 - pyruvate phosphate dikinase (PPDK) 822
 - expression profile, gene coding for 822
- q**
- quadruple squeeze 858, 859
 - first squeeze 858
 - fourth squeeze 859
 - second squeeze 858, 859
 - third squeeze 859
 - quantitative trait locus (QTL) analysis 307, 535, 536
- r**
- radiation-use efficiency (RUE) 389, 395
 - Rad50 protein 265
 - raffinose 94
 - rainfall 14, 18, 63, 73
 - amount and pattern
 - change 817, 818
 - current farming system 817
 - intergovernmental panel on climate change (IPCC), prediction 817
 - distribution 15
 - impact rice production 29
 - pattern 74
 - random amplified polymorphic DNA (RAPD) 867
 - rd29A* gene 575
 - RD29A* promoter 662
 - RDO4* gene 535
 - reactive nitrogen species (RNS) 137, 138, 141, 147, 152, 190, 739, 741, 1048
 - reactive oxygen species (ROS) 84, 85, 88, 137, 138, 210, 296, 327, 422, 431, 432, 439, 591, 606, 686, 728, 735, 739, 942, 1048
 - DNA lesions induced by 265
 - generation after UV-B irradiation in plant cells 297
 - harmful effects 1048
 - induce genotoxic stress 256, 257
 - mutualistic symbiotic interactions 1048
 - key requirements 1048
 - pathogenesis 1048
 - plant–microbe interactions 1048
 - reactive nitrogen species (RNS) 1048
 - scavenging, and detoxifying enzymes 255
 - as signaling molecules 1048
- reactive oxylipins (RES) 328
- recalcitrance, of soil organic matter 402
- recalcitrant carbon, in bioenergy crops 354, 355
- receptor-like kinases (RLKs) 88
- receptor protein kinase 591
- REDD (Reduced Emissions from Deforestation and Forest Degradation) 376
- Reducing Emissions from Livestock Research Program 71
- regulatory component of ABA receptor 1 (RCAR1) 423
- respiration 956, 957
 - alternative oxidase pathway (AOX)
 - *de novo* synthesis 957
 - limiting reactive oxygen species (ROS) production 956
 - nonproton-driven carrier 956
 - overexpression 956
 - salicylic acid regulation 956, 957
 - ubiquinol oxidation 956
 - alternative respiration pathway 956
 - induction 956
 - *de novo* protein synthesis 957

- isolated mitochondria 957
- *NtAOX1* gene 956
- expression 956
- transcript 956
- salicylic acid
 - concentration 957
 - impact on mitochondrial function 957
 - induced inhibition 957
 - treated tobacco cells 957
- stimulation 957
- tobacco cell suspension culture 956
- rhizobacteria 332
- Rhizobium* 486
- Rhizophoraceae 686
- Rhizophora mangle* 689
- Rhizophora mucronata* 201
- rhizosphere fungi 332
- rhizosphere/rhizodeposition 1015, 1016
 - autotrophic organisms, plants 1015
 - bacteria 1015
 - plant associated 1015
 - plant interactions 1015
 - definition 1015
 - ecological importance, rhizodeposition 1015
 - neutrophil extracellular traps 1015
 - photo-assimilated compounds, 1015
 - plant root secretion 1015
 - root colonization 1015
 - root exudates, role 1015
- ribulose biphosphate carboxylase (RuBisCO) 299
- ribulose biphosphate (RuBP) 30
 - nitrogen for regeneration 30
- rice
 - chimeric bifunctional *TPSP* gene 42
 - cultivation as main source of GHG emissions 69
 - DREB transcription factor 610
 - expression of *OsABI5* 593
 - *OsTPP1* expression in 42
 - pollen germination in 29
 - protoplast, ABA require activation of 529
 - submergence tolerance 66
 - Swarna-Sub1 66, 67
- rice domestication
 - amino acid substitution (K79N) 656
 - BEL1-type homeobox gene 656
 - genomic fragment 656
 - MYB3 transcriptional regulator 655
 - *O. rufipogon* 656
 - *O. sativa* ssp. *indica* 656
 - plant propagation 655
 - progenies 655
 - reducing grain shattering 655, 656
 - seed 655
 - seed shattering in chromosome 1 (*qSH1*), QTL 656
 - varieties, seed shattering 656
- Rice-Wheat System
 - adaptations 65
- risk management 74, 75
- RNA interference (RNAi) 482
 - silencing 449
- RNA-mediated gene suppression
 - methods 497
- RNA polymerases 256, 606, 620
- RNA, stabilization of 469–471
- RNS. *see* reactive nitrogen species (RNS)
- Rof1* gene 446
- role of transcription factors
 - *escherichia coli* 641
 - gene expression 641
 - activation domains (ADs) 643
 - diagram, eukaryotic promoters 643
 - eukaryotes 641, 642
 - heterodimerization 643
 - proteins 642, 643
 - sigma 642
 - historical perspective 641–644
- root hair proliferation 997–999
 - *Arabidopsis* 998
 - mutants screening 998
 - Pi-deficient roots 998
 - basic helix–loop–helix (bHLH) transcription factor, RSL4 997
 - DELLA protein accumulation 999
 - ethylene signaling 998
 - F-box protein, FBX2 997
 - gibberellic acid–DELLA system 999
 - gibberellins role 998
 - glycerophosphodiester phosphodiesterase genes 998
 - key regulators 997
 - novel transcription factor BHLH32 997
 - phenes 997
 - Pi-deficiency responses 998
 - Pi starvation response 997
 - Pi-sufficient conditions 997
 - plant hormones role 998
 - proteasome-mediated protein degradation inhibition 998
 - root hair density 997
 - secondary messengers 997
 - soil P availability 997
 - ubiquitin proteases 998
 - role 998

- ubiquitin-specific protease, UBP14 998
- roots, as primary organ sensing soil environment 330, 331
- root-shoot communication 333
- signaling compounds 333
- root “stem cell niche” 307
- root system architecture (RSA) 306
- application of phenomics in stress-associated 319, 320
- breeding approaches to improve 308
- conventional breeding approach 308, 309
- genomic approaches
- to identify regulators, associated with abiotic stress tolerance 311, 312
- identification of QTLs
- associated with specific traits, and stress tolerance 309–311
- marker-assisted selection (MAS) 308
- minirhizotrons tool, use of 320
- polyamines and osmotic regulators in stress-induced modulation 314, 315
- RooTrak software 320
- small RNA-mediated regulation of 317–319
- traits 308
- transgenic approaches to improve, abiotic stress tolerance 313, 314
- use of “RootReader3D” 320
- ROS. *see* reactive oxygen species (ROS)
- RSA. *see* root system architecture (RSA)
- RuBisCO degradation 30
- RuBisCO enzyme 818
- activity 818
- RUS2 protein 298
- S**
- SAAT sequences 335
- Saccharomyces cerevisiae* 263, 463
- Saccharum officinalis* 1022
- SAC51 gene 43
- Salicornia bigelovii* 695
- Salicornia brachiata* 686
- salicylic acid 326, 327, 608, 712
- salicylic acid (*ortho*-hydroxy benzoic acid) 940
- derivatives 941
- structure 941
- varying basal levels 940
- salinity 163, 179, 326, 368
- chlorophyll content in plant tissues 196
- genes conferring tolerance 67
- irrigation-induced 163
- soil 164
- tolerance by plants 189
- salinity stress, abiotic stress 966, 967
- antioxidant enzymes 967
- activities 967
- damaging effects salinity 967
- enhanced tolerance 966
- metabolic perturbations, plants 966
- osmolytes (proline) 966
- accumulation 966
- preadaptation, seedlings 967
- salicylic acid 966
- capability 967
- effects on plants 966
- exogenous application 966
- pretreatment 967
- salinity-stressed plants 967
- tolerance 967
- salinity stress signaling 82
- salinity tolerance 444, 445, 686, 824–826, 825
- comparative study 824
- controlled by 824
- effects on plant performance 824
- leaf hydrogen peroxide 826
- non-stressed rice plants leaves 826
- oxidative injury 826
- plant adaptations 825
- pokkali rice cultivation 824, 825
- feature 825
- rice cultivation areas survey 825
- salicylic acid 826
- foliar application 826
- salt-induced ROS 826
- salt tolerance engineering 826
- weeds 824
- *Diplachne fusca* 824
- studies 824
- salinization 334
- Salsola baryosma* 686
- Salsola kali* 686
- salt concentrations (NaCl), in soil 326, 334
- salt overly sensitive (SOS) kinases 438
- salt stress 179, 713, 714
- defined 164–167
- perception 167, 168
- primary/secondary effects 172
- osmotic and ionic phases 172–174
- oxidative stress 176, 177
- role of SOS pathway in ion homeostasis 174, 175
- signaling 168
- abscisic acid signaling 169, 170
- Ca²⁺ signaling 168, 169
- pH in signaling 169
- phospholipid signaling 171, 172
- salt stress sensitivity 140

- salt tolerance 38
- salt-tolerant genes 687
- Salvia officinalis* 970
- SAMDC gene 43
- Sapindus mukorossi* 366
- Sapindus trifoliatius* 366
- saturated hydraulic conductivity 16
- saturation vapor pressure 6
- Schizosaccharomyces pombe* 263
- seasonality 5
- seasonal water patterns in maize 15
- sea waters 63
- secondary metabolism impact
 - catechin biosynthesis 923
 - climate change scenario 919–924
 - CSIR-institute of Himalayan bioresource technology, India 922, 923
 - *de novo* transcriptome sequencing 920
 - endangered species 920
 - flavonoid biosynthetic pathways 923
 - drought stress effect 924
 - free air CO₂ enrichment (FACE) 922
 - free air temperature increase (FATI) 922
 - high-altitude region confined species 919
 - iridoid moiety 920
 - monooxygenase activity 922
 - phenylpropanoid biosynthetic pathways 923
 - drought stress effect 924
 - phenylpropanoid pathways 923
 - photosynthesis 919
 - primary products 919
 - secondary products 919
 - *Picrorhiza kurroa* Royle ex. Benth. perennial herb 920
 - genes overexpression functional categories, compared to 922
 - hepatoprotectivity 920
 - picrosides biosynthetic pathway, temperature change 921
 - relationship between 920
 - primary and secondary metabolism 920
 - secondary metabolites 919
 - stevia 923
 - diterpenoid steviol glycosides 923
 - tea 923
 - catechins 923
 - climate change, response 923
 - leaves major source of 923
- secondary metabolites roles
 - under changing climatic conditions 706, 707
 - in drought, salt, temperature, cold, and chilling stress 713
- second-generation biofuel crops 366, 367
- second-generation Hadley centre regional model (HadRM2) 907
- semi-arid tropic (SAT) region 58
- semideciduous forests 16
- senescence 960–963, 961
 - associated genes (SAGs), salicylic acid regulation 960
 - *Arabidopsis* senescent leaves 960
 - genes enhancement 960
 - salicylic acid signaling pathway 960
 - jasmonic acid signaling, senescence regulation 960, 961 (*See also* jasmonic acid signaling)
 - occurs due to 960
 - salicylic acid accumulation 960
 - probably consequences 960
 - salicylic acid biosynthesis 960
 - salicylic acid signaling pathway 961
 - different tissues senescence process, conservation of 961
 - phytohormones 961
- serotonin 713
- Ser/Thr kinases 88, 267, 741
- Sesuvium* 686
- severe erosion 16
- sfr6* mutant 569
- SHN1/2/3* genes 608
- shoot culture, *Aloe vera* 872, 873
 - shoot multiplication, enhanced rate 872
- shoot multiplication 936, 937
 - BAP 936
 - *in/ex vitro* rooting 936, 937
 - kinetin 936
 - large scale 937
 - large-scale plantation 937
 - micropropagation 937
 - *in vitro* induced shoots 937
 - *Blepharis indica* 937
 - *Haloxylon recurvum* 937
 - *Haloxylon salicornicum* 937
 - *Leptadenia pyrotechnica* 937
- 13(*S*)-hydroperoxy fatty acids 328
- signaling molecules 31
- signaling molecules, role during abiotic stress 711
- signaling, salicylic acid 941, 942, 947, 948
 - *Arabidopsis* 947
 - *Arabidopsis NahG* transgenic plants 947
 - expression 947, 948
 - growth rate 947
 - benzothiadiazole S-methylester (BTH) 948
 - brassinosteroid signaling pathways 948
 - cell suspension culture 942

- complex role 948
- *de novo* induction 942
- *de novo* synthesis, proteins 942
- endoreduplication 947
- enzyme control 942
- key role 941
- methyl-salicylate (MeSA) 942
- esterase activity 942
- production 942
- *NahG* mutants 942
- radiolabeled salicylic acid 941
- salicylic acid depletion effect 947
- as signaling molecule 942
- signaling pathway 947
- transgenic plants, characterization 947
- systemic acquired resistance (SAR) 941
- transcription factor *OBP3*
- salicylic acid-inducible DOF (DNA binding with one finger) 947
- signal transduction 267
- singlet oxygen 210
- Sinorhizobium meliloti* 333
- SIPK (salicylic acid-induced protein kinase) 268
- siRNAs 482
- slow anion channel-associated 1 85
- small interfering RNAs (siRNAs) 277, 482
- biogenesis of 492
- role in plant stress responses for adapting to climate change 492–494
- small RNA-mediated suppression
 - approaches, application of 497–499
- small RNA (sRNA)-mediated suppression constructs 498
- small non-coding RNAs in plants 508, 509
- small RNAs 267, 425, 483
- classes of 483
- S-methylmethionine 315
- smokestack emissions 352
- SNAC1* and *SNAC2* gene 669
- SNF1-related protein kinases 2 (*SnRK2*) 423, 659
- S-nitrosogluthathione 138
- S-nitrosothiols 138, 151
- S-nitrosylation 138, 141, 151
- SNP-associated alternative tags (SAATs) 334
- sodium 191
- soft inheritance 274
- soil 63
- biodiversity 71
- C : N ratio 63
- C sequestration in soil 71
- degradation (*see* soil degradation)
- erosion 15, 16, 63
- evaporation 16
- fertility 17
- management 16, 17
- moisture 335
- nitrogen (N) mineralization 63
- nutrients 16
- organic C loss 71
- organic matter 15, 63
- salinity 28, 164, 686
- structure 16
- temperature 63
- water 15, 17
- availability 15, 17
- evaporation 15
- holding capacity 5, 15
- soil-bioenergy crops
- temporal changes of carbon 360–361
- soil C, and P loss 353
- soil conditions 28
- soil C sink potential 366
- soil degradation 15, 16
- continuous cropping practice and 16
- conventional tillage 16
- double-cropping systems 16
- eastern Paraguay 16
- Ethiopia 16
- Fraser Valley in British Columbia 16
- mechanical tillage resulted in 16
- Nepal 16
- Paudel 16
- southern Brazil 16
- Taddese 16
- Thapu 16
- soil surface, exposed to direct sunlight 15
- Solanum lycopersicum* 114, 117, 118, 121, 122, 124, 623, 646, 773, 1057, 1062
- Solanum tuberosum* 514
- solar radiation 4, 5, 251
- variability 6
- Sonneratia alba* 201
- SOS1* gene 691
- SOS genes 644
- SOS3* mutant 445
- SOS pathway 444
- Soybean
- *GmWRKY21* 38
- *GmWRKY54* 38
- miRNA expression 513
- WRKY genes 38
- soybean
- free-air carbon-enrichment (FACE) study 17
- Spartina alterniflora* 687

- spermidine 735
- spermine 142, 735
- Sporobolus spicatus* 688
- srk2d/e/i* mutant 534
- starch 348
- StMYB1R-1* gene 594
- StMYB1R-1* transgenic plants 620
- stomata
 - aperture, signal mediation of 84–86
 - guard cell development 86, 87
 - responses in control of plant productivity 82, 83
- stomatal aperture, signal mediation of 84–86
- stomatal conductance 12, 13, 14
 - C3 vs. C4 species 13
 - effects of elevated CO₂ on 13
- storms 252
- stress-activated protein kinase 268
- stress hormones 736, 737
 - ABA 737, 738
 - ethylene 738, 739
 - jasmonic acid 738, 739
 - salicylic acid 738
- stress mitigation
 - salicylic acid 963–971
 - effects on abiotic stress, model 964
 - effects on biotic stress, model 964
- stress proteins 710, 740, 741
- stress resistance
 - in plants 31
- stress-resistant genes 327
- stress-responsive gene expression 741, 742
- stress-responsive genes 556
- stress-responsive transcription 90, 91
- stress tolerance 42, 43, 1025
 - abiotic stresses 1025
 - bacterial endophytes 1025
 - cold-tolerant 1025
 - host plant physiology 1025
 - photosynthesis-related proteins 1025
 - production 1025
- stress tolerance, endophytic fungi 1042–1046
 - abiotic stress tolerance types
 - endophytes to plant species 1044, 1045
 - class 1 endophyte tolerance 1042
 - drought stress 1043
 - definition 1043
 - endophytic fungi infection 1043
 - endophytic symbiotic association 1043
 - photosynthetic pigments, changes in 1043
 - water stress 1043
 - heavy metal stress 1046
 - detoxification, heavy metals 1046
 - endophytic bacteria 1046
 - mitigating Cd stress 1046
 - species showing abiotic stress tolerance 1046
 - salt stress 1045, 1046
 - area under salt stress 1046
 - complex genetic traits 1046
 - plant response, salinity 1046
 - soil salinity 1045
 - symbiotic relationships 1046
 - UN environment program 1045, 1046
 - yield stability 1046
 - temperature stress 1043–1045
 - antioxidant enzymes 1045
 - crop species 1043
 - endophytic bacterium, effect 1045
 - heat shock proteins (HSPs) 1043
 - response mechanisms, stress tolerance 1043
 - stress tolerance mechanisms 1046–1049
 - abiotic stresses, plant response 1046
 - symbiotically conferred stress tolerance 1046, 1047
- stress-tolerant crops 31
- stress-tolerant varieties 66, 67
- strigolactones 332
- Suaeda altissima* 691
- Suaeda corniculata* 696
- Suaeda fruticosa* 686
- Suaeda japonica* 686
- Suaeda maritima* 686
- Suaeda monoica* 686
- Suaeda nudiflora* 686
- Suaeda salsa* 140, 687
- submergence-tolerance genes 67
- sucrose 42
- sucrose phosphate synthase (SPS) 570
- sucrose synthase (SuSy) 570
- sugar alcohols 94
- sulfate 251
 - assimilation 516
- sulfotransferases activity 466
- summer crop 5
- sunshine duration 5
- superoxide 93, 190, 210, 686
- superoxide dismutase (SOD) 14, 297, 694, 709
- superoxide radical-scavenging mechanisms 728
- SuperSAGE data, on transcript induction 335
- Swarna-Sub1 67
 - adoption rate 67
 - National Food Security Mission 67
 - yield advantage 67

- Swarna tolerant to multiple stresses 67
- switchgrass 365
- symbiosis 332
- symbiotic microorganisms 331, 332
- alleviate/improve abiotic stress tolerance of host plants 332, 333
- t**
- TaE3* gene 442
- TAL-effector nucleases (TALEN) 96
- Tamarix hispida* 687
- TaOPR1* (12-oxophytodienoic acid reductases) gene 444
- TaqMan probes 334
- temperature 4, 6, 760–763, 917–919
 - cell cycle/cell division 918
 - cold stress 761–763
 - acclimation 762
 - *Arabidopsis* 763
 - chilling injury 761
 - chilling/non-freezing temperatures 761
 - chilling tolerant, temperate regions 761
 - cold acclimation 762
 - cold-induced adaptation 762
 - cold-induced gene expression 762
 - cold-responsive genes 762
 - cold-responsive transcription factor 762
 - C-repeat binding factors (CBFs) cold-responsive pathway 763
 - cryo-damage 763
 - freezing injury 761
 - gene expression, reprogramming 762
 - low-temperature stress 762
 - maltose 763
 - perception 762
 - response 762
 - signaling events, low temperature plants response 762
 - stress-tolerant crop plants 762
 - transcription factors CBF/DREB1, overexpression 763
 - transgenic *Arabidopsis* 763
 - transgenic tobacco plants 762
 - cryoprotectants, generating 919
 - effect on plant performance 917
 - expression of genes 919
 - genes downregulation 918
 - global climate change 760
 - growth temperature effect 919
 - high temperature 760, 761
 - abscisic acid (ABA) 761
 - alkaloids 761
 - genetic engineering approach 761
 - glycine betaine-deficient species 761
 - hormonal balance alteration 760
 - hormonal homeostasis alteration, heat stress 761
 - low-molecular-mass organic compounds, osmolytes 760
 - membrane fluidity changes 760
 - phenolics biosynthesis 761
 - phytohormones 761
 - secondary metabolites 761
 - secondary plant products 761
 - thermotolerance 760
 - transgenic plants 761
 - histone proteins 918
 - increase, effects of 841, 842
 - low-temperature-responsive mechanisms 917
 - molecular response 917–919
 - optimum for vegetative growth 29
 - responses 7
 - annual crops 7–11
 - major challenges 11
 - stress 714, 715
 - stress-inducible genes, upregulation 918
 - stress-responsive genes 918
 - expression 919
 - tea 917
 - transcriptome 919
 - winter dormancy 919
- temperature stress, abiotic stress 967–969
 - cold stress 968, 969
 - CAT activity 968
 - chilling injury 968
 - electrolyte leakage 968
 - enhanced germination percentage 969
 - exogenous salicylic acid 968
 - hydroponic solution 969
 - treatment, salicylic acid 968
 - gaseous emissions 967
 - global mean temperature 967
 - heat stress 967, 968
 - applications, salicylic acid 968
 - heat-stressed plants 968
 - negative effects, plants 967
 - photorespiration rate, increases 968
 - photosynthetic functions 968
 - pretreatment, salicylic acid 968
 - salicylic acid, foliar spray 968
 - treatment, salicylic acid 968
 - optimum temperature, deviation 967
- Teten's equation 12
- Thellungiella halophila* 687
- thermotolerance 821–823

- acquisition of 822
- antioxidant defense mechanism 822
- antioxidant enzymes 823
 - ascorbate peroxidase (APX) 823
 - heat-stable isozymes of 823
 - superoxide dismutase (SOD) 823
- cool-season turfgras
- heat-acclimated vs. non-acclimated 822
- cytosolic Cu/Zn-superoxide dismutase (SOD) 822
- defense mechanisms 822
 - *Eupatorium adenophorum* Spreng 821
 - vs. *Eupatorium odoratum* 821
 - *Eupatorium odoratum* L. 821
 - heat shock proteins (HSPs) 822
 - accumulation 822
 - synthesis 822
- heat stress 822
- maize plants 821
 - intolerance to low temperature 821
- *Miscanthus × giganteus*, C4 grass 821
- Mn-SOD 822
- oxidative stress 821
 - protection 822
- plant adaptation 822
 - pyruvate phosphate dikinase (PPDK) 822
 - expression profile, gene coding for 822
- reactive oxygen species (ROS) 821
 - autocatalytic peroxidation 822
 - generation 822
 - reaction 822
- stress-related proteins 823
 - synthesis 823
- thermotolerance factors 431, 432
- thylakoid membranes 29, 297
- tillage
 - conventional 71
 - reduced tillage on soil C content 71
- TIR1 (transport-inhibitor response 1) 516
- tissue culture-raised plants 867
- tocopherols 255, 708
- tpk1* mutant 86
- TPP* gene 42
- TPS* gene 42
- TPSP* gene 42
- trans-acting siRNAs (ta-siRNAs) 277
- transcriptional gene silencing (TGS) 481
- transcription factors 31, 91, 424, 438, 515, 556
 - bZIP 35
 - combinatorial activity maintain root “stem cell niche” 307
 - DREB/ERF 36
 - gene discovery in stress responses 663–665
 - historical perspective 641–644
 - modulating plant adaption, classification of 592
 - AREB/ABF proteins 593
 - CBF/DREB transcription factors 594, 595
 - MYC/MYB transcription factors 593, 594
 - NAC and ZF-HD proteins 595, 596
 - WRKY transcription factors 596
 - ZF proteins 596, 597
 - NAC 37, 38
 - plant families, implicated in drought and salinity 644, 645
 - AP2/ERF family 651, 652
 - bHLH family 649
 - bZIP family 649, 650
 - HD family 653, 654
 - MYB family 37, 645
 - NAC family 650, 651
 - WRKY family 652, 653
 - role of 336
 - simplified model, representing phylogenetic distance 668
 - WRKY 38
 - ZF 38, 39
- transcriptome 94
- transcriptome analysis 914
 - chaperones expression 919
 - climate change scenario 914, 919
 - challenge 919
 - objective 914
- environmental cues tolerance genes 914
- environmental cues tolerance, gene suits 914
- transcriptomics 728, 729
- trans*-esterification 364
- transgenerational inheritance 274–278
- transgenic expression of *HaHb4* in soybean 671
- transgenic maize plants overexpressing ZmNF-YB2 672
- transgenic plants 94
 - drought-resistant, engineered by using bZIP transcription factors as transgenes 539, 540
 - functional analysis of DREB genes in 574, 575
- transgenic rice plants 38
- transgenic wheat plants 42
- transpiration 12, 13, 328
 - rate 14, 30
- transport, salicylic acid 941, 942
 - infection tolerance 942
 - reactive oxygen species (ROS) 942
 - surface-applied salicylic acid 942

- synthesis in cells 941
 - trees abiotic stress tolerance, different forest types 892–894
 - biomass production 894
 - climate change-induced abiotic stresses 892
 - degree of tolerance 892
 - dominant trees 894
 - regeneration 894
 - drought sensitive trees 894
 - extreme temperature trees 894
 - genetic variation 892
 - likely abiotic stress tolerance, organic carbon storage trends 893
 - sal coppices 894
 - Sundarban trees 892
 - *H. fomes* (Sundri), most dominant tree 892, 893
 - top dying of Sundri 893
 - trehalose 42
 - biosynthesis pathway, heterologous expression of genes 42
 - Trifolium alexandrinum* 608
 - tripeptide glutathione (GSH) 709
 - Triticum aestivum* 11, 41, 112, 113, 116, 117, 222, 231, 259, 262, 443, 460, 493, 511, 512, 575, 590, 610, 619, 1043–1045
 - heat stress, to alter miRNA expression 511
 - *TaDREB1* gene 609
 - Triticum dicoccoides* 317, 512–514
 - drought-responsive miRNAs 513
 - miRNA expression patterns 488
 - upregulation *miR398* 514
 - tropical cyclones 507
 - tropospheric ozone 818
 - agricultural yield reduction 818
 - mechanisms 818
 - meta-analytic techniques, results 818
 - tryptophan
 - in UV-B perception 299
 - tyrosine nitration 141
- u**
- ubiquitin-proteasome system (UPS) 944, 945
 - DELLA proteins 945
 - gibberellin signaling repressors 945
 - reactive oxygen species (ROS) modulation 945
 - salicylic acid/jasmonic acid signaling modulation 945
 - hormone signaling pathways 944, 945
 - NPR1 945
 - activity regulation 945
 - pathogen-infected cells 945
 - salicylic acid-induced phosphorylation 945
 - salicylic acid signaling transducer 945
 - NPR1 945
 - ultraviolet (UV)-B 295
 - effect on plants 296
 - response of plant roots to 297, 298
 - root evolution under a UV-B environment 299, 300
 - tryptophan in UV-B perception 299
 - UV-B receptors found in roots 298, 299
 - ultraviolet (UV) wavelength regions 295
 - underground water 508
 - UN food and agriculture organization (FAO) 883
 - UniTags 334
 - urban growth 73
 - UV-B radiation 252, 514, 819, 820
 - Asia-Pacific region, effects on 819, 820
 - enhanced UV-B 819
 - effects 819
 - gene pool 820
 - genetically engineered genomes 820
 - aim 820
 - high levels 819
 - effects 819
 - natural evolution, crop plant species 820
 - spring wheat, effects on 819
 - stratospheric ozone layer, depletion 819
 - effects of 819
 - tolerance (*see* UV-B radiation tolerance)
 - UV-B photoreceptors 819
 - UV-B radiation tolerance 828
 - enhanced peroxidase activity 828
 - increased UV-B radiation 828
 - plant responses 828
 - ozone layer 828
 - plant sensitivity 828
 - stratospheric ozone depletion 828
 - UV-B-absorbing compounds 828
 - UV-B exposure 828
 - biological responses 828
 - UV-B protection, protection mechanism 828
 - weeds 828
 - exposure study 828
 - UV-mediated ROS generation 296, 297
 - UV radiation, ozone stress 969, 970
 - ozone, deteriorating effects 969
 - ozone exposure 969
 - photochemical reactions 969
 - phytohormone signaling role
 - ozone-induced cell death 970

- protection mechanisms 969
- salicylic acid
- accumulation 969
- exogenous application 969
- photochemical efficiency 969
- UV-B-stressed plants 969
- UVR8 gene 298

V

- vacuolar H⁺-PPase (V-PPase) 695, 696
 - valley exit jet 904
 - Vallisneria spiralis* 692
 - vaporization 6
 - V-ATPase 695
 - vegetative growth 946, 947
 - cost-beneficial phytohormones 946
 - exogenous salicylic acid 946, 947
 - salicylates treatment 946
 - salicylic acid 946
 - antiproliferative properties 947
 - exogenous application 946
 - foliar application 946
 - growth-promoting effects 947
 - growth-stimulating effect 947
 - rooting enhancement 946
 - soil nutrient solution 946
 - Verticillium dahliae* 43
 - Vicia faba* 263
 - village forests 886, 887
 - biological diversity 886
 - dominant trees 887
 - homestead forests, Bangladesh 886
 - products 886
 - *in situ* conservation, agro-biodiversity 886
 - surplus products 886
 - total area 883, 886
 - vitamin A-improved rice 787
 - vitamin A, malnutrition 787
 - vitamin C 255
 - vitamin E 255
 - VP1/ABI3 gene 532
- ## W
- warming temperatures 4
 - water 62
 - balance 63
 - cycles 346
 - demands 5
 - erosion 16
 - holding capacity, of plant tissue 306
 - productivity 81
 - responses 11–17
 - salinity 63
 - protection mechanisms of photosynthesis during 92–94
 - salicylic acid 970
 - ameliorating role 971
 - damaging effects alleviation 970
 - efficiency depends on 971
 - exogenous application 970
 - pivotal role 970
 - treatment 970
 - tolerance 82
 - key transcription factors 91, 92
 - signaling mediation by membrane-localized proteins 87–90
 - signaling/transcriptional control 87
 - stress-responsive transcription 90, 91
 - water-use efficiency (WUE) 12, 81, 348, 1048
 - basic equation for 12
 - critical parameter in 15
 - definition 1048
 - endophyte-positive plants
 - vs. endophyte-negative plants 1048
 - enhancement through genetic selection/crop management 15
 - evaluation 13
 - growth and development processes and 17, 18
 - measure of 1048
 - *Neothyphodium coenophialum* infection 1048
 - optimizes WUE of C assimilation, in arid habitats 367
 - physical variables 14
 - physiological variables 14
 - water vapor 6, 507
 - wax content 14
 - WCS120 gene 564
 - weather patterns 28
 - wheat
 - *Cor/Lea* genes 448
 - genes, used for raising abiotic stress-tolerant transgenics 440
 - genotypes 30
 - improvement

- functional and comparative genomics approaches for 449, 450
 - overexpression of *mtlD* gene 42
 - physiological and molecular adaptive strategies in 438–440
 - systems 17
 - *TaWRKY2* and *TaWRKY19* genes 38
 - wild underutilized species 934
 - importance 934
 - wilting coefficient 191
 - wind velocity 63
 - WIPK (wound-induced protein kinase) 268
 - woody plants, grow in intertidal zones with high salinity 190
 - World Water Day 859
 - WRKY family
 - five subgroup 652, 653
 - ABA signaling 653
 - electrophoretic mobility shift assays (EMSAs) 653
 - homo- and heterodimerization, proteins 653
 - immunoprecipitation 653
 - N-terminal LZ 653
 - higher plants 652, 653
 - W-box 652
 - zinc finger-like motif, three groups 652
 - WRKY transcription factors 596, 624–626
 - WUE. *see* water-use efficiency (WUE)
- x**
- x-ray repair cross-complementing factor 1 (XRCC1) 271
 - xylem 333
 - xylem biopolymers 366
- y**
- yeast one-hybrid assays 659
 - yield gap 210
- z**
- Zea mays* 34, 115, 119, 261, 353, 560, 575, 590
 - zeaxanthin 707
 - ZF-HD regulon 614, 615
 - ZF transcription factors 596, 597
 - zinc 608
 - zinc finger homeodomain 1 (ZFHD1) 662
 - zinc finger nucleases (ZFN) 96
 - Zinc finger transcription factors 93
 - ZmbZIP17* gene 593
 - Zostera* 686
 - Zosteraceae 686
 - Zostera marina* 692