

Index

a

- angular overlap model (AOM)
 - angular distortion 42
 - f orbital energy 41
 - parameters 41
 - properties 41
 - superposition model 41
- aquo* ions
 - angular distortion 40
 - best fit factors 39
 - coordination geometry 38
 - diagonal matrix 39
 - ground multiplet 40
 - non-diagonal matrix 39
 - symmetry operations 39
- atomic force microscopy (AFM) 298
- atomic mean field integrals (AMFI) 86

b

- ball and stick (BS) approach 173, 175
- bidentate ligands 51
- Brillouin equation 70
- Brillouin function 81
- broken symmetry approach (BS) 130

c

- carboxylates 17, 62
- chemical exchange saturation transfer (CEST)
 - technique
 - advantage 417
 - diaCEST agents 418
 - magnetization transfer asymmetry 416
 - PARACEST agents 418
 - proton exchange 417, 418
 - saturation transfer technique 415
 - spin distribution 415, 416
 - Z-spectrum 417

chemisorption 306

colossal magnetoresistance (CMR) 297

configuration interaction (CI) method 85

copper acetate

- *ab initio* techniques 132
- anisotropic components 133
- covalency effects 133
- DDCI3 133
- dinuclear structure 132
- effective Hamiltonian theory 135, 136
- perturbation theory 133
- post-CASSCF method 134
- relative energy 135
- singlet triplet separation 132
- spin orbit contribution 134
- T dependence 132
- temperature dependent magnetic susceptibility 132

crystal field (CF)

- Dq parameter 38
- group theory 37
- non-vanishing parameters 37
- operator equivalent 35
- parameters 35
- qualitative agreement 35
- radial integrals 35
- Stevens/Wybourne approach 36
- symmetry requirements 36
- tetragonal environment 36
- transition metals 38
- crystal field strength 80
- Curie law 70
 - Curie–Weiss law 71
 - effective magnetic moment 70
 - Ln(III) ions magnetic information 71
 - transition metal ions, wave functions 72
- Curie spin relaxation 402
- Curie–Weiss law 71

- cyclooctatetraene 17
- cyclooctatetraene dianion (COT)
 - chloride salt 55
 - coordination sphere 55
 - crystal structure 55
 - cyclopentadienyl derivatives 55
 - sodium salt 55
- cyclopentadiene 17

- d**
- d* magnetic orbitals
 - HS-LS transition 11
 - octahedral environments 12
 - SCO 10, 12
 - SMM 13
- d* orbitals
 - aufbau approach 163
 - chirality
 - crystal structure 170
 - ECM 169
 - magnetic chirality 169
 - MChD 169, 170
 - non-enantiopure materials 169
 - non-zero contribution 170
 - refraction and absorption 170
 - space groups 170
 - properties 169
 - spin chirality 169
 - 4*d* and 5*d* ions
 - exchange coupling constants 168
 - J coupling constants 165
 - Jahn Teller distortion 167
 - magnetic properties 165
 - NiM₂ adduct 168, 169
 - octahedral coordination 165
 - photomagnetism 165
 - pseudo-spin operator 167
 - six carbon donors 165
 - trigonal bipyramidal pentanuclear cluster 165
 - 3*d* metal ions 164, 165
 - DFT
 - Bn parameters 172
 - BS approach 173, 175
 - epochal paper 172
 - molecular structure 172
 - dynamic Jahn-Teller 163
 - f-d interactions 171
 - Gd transition metals 176
 - Ham effect 163
 - high spin state 163
 - low spin state 163
- density functional theory (DFT)
 - Coulomb repulsion 87
 - *D* values 91, 92
 - electron density 87
 - electron kinetic energy 87
 - electron nucleus interaction 87
 - exchange correlation energy 87
 - *in silico* approach 91
 - Kohn Sham equations 87
 - Koopmans theorem 87, 88
 - octahedral environment 92
 - polynuclear system 90
 - quasi-routine technique 87
 - wavefunction approach 87
 - Xα method 87
- diamagnetic (diaCEST) agents 418
- diamagnetic susceptibility 69
- dI/dV mapping 315
- difference dedicated configuration interaction (DDCI) 135
- diketonates 17, 58, 59
- discrete Fourier transform (DFT)
 - Bn parameters 172
 - BS approach 173, 175
 - epochal paper 172
 - molecular structure 172
- double decker Tb(III) derivative 324
- double electron-electron resonance (DEER) sequence 391, 392
- double quantum coherence (DQC) 159
- Douglas–Krell–Hess (DKH) Hamiltonian 86
- dressing ions
 - ligand modes 33
 - metal-ligand interaction 33
 - paddlewheel motif 33
 - transition metal ions 33
- Dy-Pc based devices 316
- dysprosium (Dy) 18
 - AFM interaction 283
 - angular dependence 279, 280
 - archetypal Dy₃ 283
 - axial symmetry 279
 - binary symmetry 279
 - bottom up approach 287
 - chirality 282
 - components 277
 - dynamic properties 286
 - Eu(III) ion 291
 - exchange coupling constants 291
 - 4*f* orbital 279
 - features 277
 - field dependence 280, 281
 - Heisenberg cluster 283
 - Ising spins 285
 - Kramers doublets 280, 281
 - local anisotropy axes 285

- magnetic properties 277, 278
- magnetic susceptibility 288
- magnetic tensor 279
- magnetization 283
- microstates 284
- monoclinic symmetry 278
- normalized magnetization 289
- Pauli matrices 284
- performing measurement 278
- perturbation 292, 293
- relaxation mechanism 289
- single-ion easy anisotropy axes 288
- spin chirality 286
- toroidal dipoles 282
- triangular cluster 284
- triangular moieties 288, 289
- triclinic crystals 289
- Zeeman splitting 285

- e**
- effective magnetic moment 70
- electron paramagnetic resonance (EPR)
 - DEER sequence 391, 392
 - dipolar fields 378
 - fine structure 377
 - GdDTBSQ
 - axial symmetry 379
 - fourfold symmetry 381
 - g values 381, 382
 - GdTrop 380
 - HF-EPR spectra 379
 - magnetic evidence 382
 - rhombic symmetry 380
 - X band spectrum 379
 - zero field splitting tensor 380, 381
 - ground states 375
 - hyperfine term 375
 - intense lines 377
 - intrinsic intensity 376
 - Ln nicotinates
 - angular dependence 388
 - components 389
 - dinuclear structure 388
 - dipolar and exchange interaction 388, 389
 - hyperfine splitting 391
 - Ising symmetry 388
 - LaND 389
 - pseudo-XY symmetry 388
 - orbit 382, 383
 - tetragonal symmetry 376, 377
 - transition metal
 - coordination polyhedron 384
- -- experimental curves 386
- -- polycrystalline powder 384, 385
- -- principle 384
- -- two not-equivalent molecules 385, 386
- -- x, y and z axes 386, 387
- -- Zeeman energy 375
- electron spin resonance (EPR), *see* electron paramagnetic resonance (EPR)
- electron spin resonance (ESR) spectroscopy 153
- electronic structures 16
- electronic structures, free ions
 - Aufbau configuration 25
 - cation and electric dipoles 25
 - 4f orbitals 26
 - Fermi-Dirac statistics 27
 - ground multiplets 27, 28
 - magnetic field 31, 32
 - Russell-Saunders coupling scheme 27
 - spin orbit coupling
 - f orbitals 30
 - features 29
 - Hamiltonian energy 29
 - interelectronic repulsions 28
 - j-j coupling scheme 31
 - lanthanides 30
 - quantum number 29
 - Slater determinant 30
 - transition metal ions 28
 - tripositive lanthanide ions 29
 - enantiopure chiral magnets (ECM) 169
 - exchange and superexchange interaction 17
 - experimental susceptibility 69

- f**
- f magnetic orbitals 13, 14
- f orbital system 176, 177
- ferromagnets 17
 - anisotropy 117
 - bricks, infinite array
 - icosidodecahedral structure 103
 - types 103
 - unpaired electrons 102
 - zero dimensional 102
 - building blocks 99, 100
 - dipolar interactions
 - coordinate system 112
 - D12 tensor 111
 - EPR spectroscopy 113
 - equivalent Hamiltonian 111
 - exchange interaction 113
 - g tensors 111
 - Gd(III) ions 110, 111, 112
 - slow decay 110

- f**
- ferromagnets (*contd.*)
 - – Y ions 111
 - giant spin 114
 - isotropic and anisotropic interaction 115
 - magnetic interactions
 - – Cr(III) free ion 104
 - – Curie constant 106
 - – Curie–Weiss law 106
 - – dipolar 105
 - – exchange interaction 105
 - – group theory 108
 - – HDVV Hamiltonian 104
 - – Ising approach 108
 - – Landé interval rule 105
 - – magnetic ordering 108, 109
 - – spin alignment 109
 - – spin angular momentum 104
 - – temperature dependence 106
 - – Van Vleck equation 105
 - Mn12 molecule 116, 117
 - non collinearity 117
 - one dimensional array 100
 - orbital degeneracy
 - – diamagnetic dilution 123
 - – electronic ground states 119
 - – exact diagonalization 120
 - – isotropic and anisotropic terms 124
 - – Kramers doublets 122
 - – lanthanide derivatives 123
 - – Lines model 120
 - – Ln isolation 121
 - – matrix elements 121
 - – Pauli matrices 120
 - – qualitative diagram 123
 - – SH interaction 121
 - – SOC 120
 - – Zeeman term 122
 - paramagnetic system 100
 - single building block 115
 - spin Hamiltonian approach 113, 114
 - spin Hamiltonian parameters 116
 - spin orientation 100
 - three dimensional representation 101
- g**
- generalized gradient approximations (GGAs)
 - functionals 92
 - giant magnetoresistance (GMR) 296
 - Glauber model
 - configurations 254
 - dinuclear copper compound 255
 - domain wall 256
 - infinite array 255
 - Orbach mechanism 254
- h**
- Hamiltonian approach 16
 - highly oriented pyrolytic graphite (HOPG) 306
 - hybrid organic magnetic molecules 318
- i**
- intermediate coupling 299
 - intrinsic optical bistability (IOB) 113
 - ions magnetism
 - Curie law, *see* Curie law
 - diamagnetic susceptibility 69
 - experimental susceptibility 69
 - magnetic induction 69
 - paramagnetic magnetization 70
 - Van Vleck equation, *see* Van Vleck equation
 - irreducible tensor operators (ITO) 367
- j**
- Jahn-Teller theorem 77, 136
- k**
- Kondo effect 300, 301, 316
- l**
- Landau–Zener–Stückelberg (LZS) model 208
 - lanthanide induced relaxation (LIR) 397
 - lanthanide induced shift (LIS) 397
 - lanthanide shift reagents (LSR)
 - diamagnetic molecules 406
 - g tensor 405
 - linear expansion 404
 - secondary internal field 404
 - shift reagent 405
 - light-induced spin-state trapping effect (LIESST) 12
 - Lines model 120
 - luminescence
 - absorption/emission processes 425, 426
 - antenna effect 429
 - biological assays 432
 - charge transfer transitions 425
 - definition 424
 - 4f–5d orbitals 424, 425
 - forbidden transitions 426
 - hypersensitive/pseudo quadrupolar transitions 426, 428
 - light emission 426

- medical imaging 432
- non-radiative pathways 429
- OLED 429–432
- properties 426, 427

- m**
- magnetic anisotropy
 - *4f* charge density 49
 - *f* orbitals 48
 - properties 47
 - quadrupolar moment 48
 - spin orbit coupling 47
 - uniaxial anisotropy 47
- magnetic circular dichroism (MCD) 169
- magnetic coercitivity 296
- magnetic resonance imaging (MRI)
 - CEST
 - advantage 417
 - diaCEST agents 418
 - magnetization transfer asymmetry 416
 - PARACEST agents 418
 - proton exchange 417, 418
 - saturation transfer technique 415
 - spin distribution 415, 416
 - Z-spectrum 417
 - chemical shift 414
 - clinical applications 412
 - contrast agents 413
 - contrast reagents 409, 410
 - coordinated water molecules 413
 - coordination number 412
 - Curie spin relaxation mechanism 411
 - electronic relaxation time 412
 - inner-sphere relaxation mechanism 410
 - molecular structures 413
 - nuclear longitudinal 410
 - principles 409
 - relaxation time 409
 - residence time 414
 - transverse relaxivity 410
 - ZFS 411
- magneto chiral effect (MChE) 169, 170
- magnetocaloric effect (MCE)
 - commercial application 423, 424
 - constant pressure 421
 - magnetic degrees of freedom 423
 - magnetic entropy 421
 - n-non-interacting spin 423
 - paramagnetic demagnetization 422
 - S-T diagram 421
 - spin frustration 423
 - thermal hysteresis 422
 - vapor-compression cycle 424
- magnetocrystalline anisotropy, *see* magnetic anisotropy
- magnetogiric ratio 398
- metal organic frameworks (MOFs) 18
 - analysis 361
 - anisotropy
 - applications 363
 - ball-and-stick view 364
 - magnetization value 364
 - symmetric core 363
 - temperature dependence 365
 - cationic cluster 361
 - cobalt(II) 355
 - crystallographic analysis 360
 - Cu-dimers 366
 - 2D structure 369, 370
 - 3D system 354
 - definition 353
 - depopulation mechanism 366
 - development 352
 - 3d–4f heterometallic cluster 365
 - exchange mechanism 366
 - features 363
 - gadolinium 353
 - giant clusters 358
 - growth mechanism 351
 - inner shell 360
 - ITO 367
 - magnetic susceptibility 362
 - metal ions 362, 363
 - metamagnet 355, 356
 - M–L–M connectivity 353
 - molybdates
 - antiferromagnetic coupling 359
 - Hilbert space 358, 359
 - isomorphous 360
 - structures 358
 - temperature dependence 359, 360
 - monoclinic 355
 - M–X–M bonding 354
 - one-dimensional coordination polymers 354, 355
 - orthorhombic modification 355
 - photoluminescence properties 357
 - point group symmetry 367
 - quantitative indication 367
 - red fluorescence 357
 - spin states 367
 - structures 352
 - Yaghi's secondary-building-units 354
 - ZFC-FC measurement 356
- molecular magnetism (MM) 297
 - complexity 7
 - *d* magnetic orbitals

- molecular magnetism (MM) (*contd.*)
 - HS-LS transition 10
 - octahedral environments 12
 - SCO 10, 12
 - SMM 13
 - diamagnets 2
 - dipolar magnets 3
 - exponential growth 15
 - *f* magnetic orbitals 13, 14
 - *f* orbital approach 1
 - goals 14
 - graphic outline 16
 - lanthanides
 - heterogeneous catalysis 19
 - magnetocrystalline anisotropy 20
 - mischmetal/Auer metal 18
 - Nd, Dy, Eu, Tb, and Y 20
 - optical spectra 19
 - peculiar property 19
 - permanent magnet industry 20
 - SmCo₅ and Nd₂Fe₁₄B 19, 20
 - material description 2
 - metallic and ionic lattices 4
 - organic molecules 4
 - origin 5
 - *p* magnetic orbitals
 - dithiadiazolyl radical 9
 - EPR spectroscopy 8
 - inorganic radicals 9
 - intermolecular reactivity 9
 - nitroxide and tetracyanoethylene 8
 - quantum mechanics 3, 4
 - SI vs. emu 21
 - tantalizing expression 1
 - time and space 2
 - molecular orbital (MO) 17
 - charge transfer processes 127, 128
 - complexity
 - D tensors 88, 90
 - EPR spectra 89
 - g tensors 88
 - ground state 90
 - INDO methods 90
 - LF formalism 88
 - spin orbit coupling 89
 - ZFS parameters 89
 - copper acetate
 - *ab initio* techniques 132
 - anisotropic components 133
 - covalency effects 133
 - DDCI3 133
 - dinuclear structure 132
 - effective Hamiltonian theory 135, 136
 - perturbation theory 133
 - post-CASSCF method 134
 - relative energy 135
 - singlet triplet separation 132
 - spin orbit contribution 134
 - T dependence 132
 - temperature dependent magnetic susceptibility 132
 - correlation effects
 - 4f orbitals 86
 - *ab initio* techniques 84
 - AMFI 86
 - CASSCF/CASPT2 states 85
 - configuration interaction 84
 - Coulomb correlation 84
 - coupled cluster 84
 - Dirac approach 85
 - DKH Hamiltonian 86
 - electron correlation 84
 - g tensor 86
 - MBPT 84
 - SCF 84
 - d orbitals, structural requirements 129
 - degenerated states 136, 137
 - DFT
 - Coulomb repulsion 87
 - *D* values 91, 92
 - electron density 87
 - electron kinetic energy 87
 - electron nucleus interaction 87
 - exchange correlation energy 87
 - *in silico* approach 91
 - Kohn Sham equations 87
 - Koopmans theorem 87, 88
 - octahedral environment 92
 - polynuclear system 90
 - quasi-routine technique 87
 - wavefunction approach 87
 - X α method 87
 - DOTA complexes
 - CASSCF-CASPT2/RASSI-SO method 94
 - Dy-Na interaction 94
 - experimental and calculated luminescence spectra 94
 - LnDOTAa 95
 - magnetic anisotropies and luminescence 93
 - Na[Dy(DOTA)(H₂O)]·4H₂O compound 93
 - pseudo four-fold symmetry axis 95
 - excited state 128
 - *f* orbitals and orbital degeneracy
 - Cartesian components 139
 - HDVV 139

- - octahedral symmetry, splitting 138
- - techniques 138
- - wavefunctions 139
- - Yb ions 138
- ferromagnetic contribution 128
- ligand field model 83
- physical properties 127
- quantitative calculations 83
- schematic diagram 128
- SH parameters
- - broken symmetry approach 130
- - covalency effects 130, 131
- - eigenfunctions 130
- - error cancellation effects 131
- - exchange parameter 132
- - Heisenberg constant 130
- - maximum delocalization 130, 131
- valence bond theory 129
- molecular spintronics
 - CMR 297
 - detection process 308
 - detection schemes 303, 304
 - GMR 296
 - hybrid organic magnetic molecules 318
 - magnetic field sensors 296
 - magnetic multilayers 296
 - magnetic substrate 319
 - magnetism measurement 310
 - Mn₁₂ 317
 - molecular organization on surfaces 303
 - molecules and mobile electrons
 - - Coulomb blockade 300
 - - Coulomb oscillations 302
 - - electron beam lithography fabrication 301
 - - HOMO/LUMO properties 299
 - - hybridization effects 299
 - - Kondo effect 300, 301
 - - metal thiolates 301
 - - MM 297
 - - molecular electronics 298
 - - molecule-electrode interaction 298
 - - QD 297
 - - Schottky barrier 299, 300
 - - SMM 297
 - - STM 298
 - molecules and surfaces selection
 - - bottom up approach 307
 - - chemisorption 306
 - - immobile molecules 307
 - - MPC monolayers 305
 - - multilayer techniques 305
 - - physisorption 305
 - - self-assembly/self-organized growth 307
 - - sulfur strategy 306
 - - molecules purity 307
 - - nanostructured magnetic devices 295
 - - nuclei
 - - electronic magnetic moment 321
 - - ground states energy 322
 - - non coherent tunneling 322
 - - nuclear spins 321
 - - pc ligands 322
 - - spin flip dynamics 324
 - - STS 322
 - - ¹⁶⁵Tb nucleus 322
 - - TbPc₂ properties 322
 - - transition matrix, QTM 323
 - - waiting time 323
 - - organizing and addressing molecules 302
 - - Pc family
 - - dI/dV mapping 315
 - - Dy-Pc based devices 316
 - - four-lobe molecule analysis 315
 - - Kondo effect 316
 - - Kondo temperature 316
 - - Ln molecules 314
 - - Ln(III)-Pc molecules 316
 - - OFETs 314, 316
 - - Tb(Pc)₂, SIM nature 316
 - - TbNPcPc double-decker molecules 317
 - - TbPc₂ molecule 314
 - - single radical transport
 - - archetypal radical 312
 - - Co-Salen molecule 312
 - - electrode with radical bridge 312
 - - high-resolution topography 314, 315
 - - 2'-nitronylnitroxide-5'-methyl-[1,1'; 4',1'']terphenyl-4,4''-dicarbonitrile radical 315
 - - nitronyl nitroxides 313
 - - SAM 312
 - - TOV 312
 - - verdazyl derivatives 312
 - - spin valve device 324
 - - TMR 297
 - - VASP 319
 - - X-rays
 - - absorption 308, 309
 - - dipole selection rule 309
 - - L_{2,3} edges 310
 - - one-electron model 310
 - - XMCD 308
 - - XASs 308
 - molybdates
 - - antiferromagnetic coupling 359
 - - Hilbert space 358, 359
 - - isomorphous 360

- molybdates (*contd.*)
 - structures 358
 - temperature dependence 359, 360
- monodentate ligands 51

- n**
- nanostructured magnetic devices 295
- n-electron valence second-order perturbation theory (NEPTV2) method 135
- nitronyl nitroxides (NITs) 17
 - automatized method 146
 - bidentate ligand 60
 - computational analysis 159
 - diamagnetic nitrone 156, 157
 - DQC 158
 - energy levels 156
 - experimental evidence 62
 - first principle bottom up 146, 147
 - initial reagents 60
 - J_{AB} values 147
 - Kramers doublets 156
 - lead dioxide 60
 - metal ion/radical ratio 61
 - molecular structure 158
 - N–N distances 160
 - parameters 156
 - PELDOR 158
 - β -phase crystals 146
 - polymorphs 146
 - sodium periodate 60
 - spin–spin interaction 160
 - structural instability 146
 - χT product 155, 156
 - temperature dependence 148
 - Van Vleck equation 147
 - X band spectrum 158
- nuclear magnetic resonance (NMR)
 - *f*-block element 395
 - lanthanide ion
 - - crystal field effects 400
 - - Curie relaxations rates 403
 - - Fermi contact 397, 398
 - - induced magnetic moment 398
 - - isostructural 396
 - - isostructural series 399
 - - isotropic hyperfine coupling constant 398
 - - LIR 396, 397
 - - LIS 396, 397
 - - LnDOTAM series 401
 - - non-zero spin 401
 - - paramagnetic shift 400
 - - paramagnetism 395
 - - pseudocontact shift 399
 - - reference compound 400
 - - relaxation mechanism 402, 403
 - - resonant nuclei 397
 - - rotational correlation time 404
 - - second-order magnetic axial anisotropy 400
 - - signal linewidth 404
 - - LSR
 - - diamagnetic molecules 406
 - - *g* tensor 405
 - - linear expansion 404
 - - secondary internal field 404
 - - shift reagent 405
 - - properties 396
 - - uses 395

- o**
- orbital degeneracy 176, 177
- organic field-effect transistors (OFETs) 314, 316
- organic light-emitting diode (OLED)
 - advantages 430
 - cell structure 430
 - Dexter transfer mechanism 431
 - dipole–dipole coupling 431
 - electrons transport mechanism 430
 - guest phosphorescent molecules 431
 - parameters 431, 432

- p**
- p* magnetic orbitals 17
- dithiadiazolyl radical 9
- EPR spectroscopy 8
- inorganic radicals 9
- intermolecular reactivity 9
- nitroxide and tetracyanoethylene 8
- organic ferromagnet 8
- p* magnetic orbitals system
 - allyl radicals 145
 - antiferro- and ferromagnetic interactions 144
 - charge transfer mechanism 145
 - ferrimagnetic approach 143
 - metallorganics magnets
 - building blocks 149
 - - *d* orbitals 149
 - - *f* orbitals 149
 - - TCNE 150
 - - TCNQ 151
 - nitronyl nitroxides
 - - automatized method 146
 - - computational analysis 159
 - - diamagnetic nitrone 156, 157
 - - DQC 158

- - energy levels 156
 - - first principle bottom up 146, 147
 - - J_{AB} values 147
 - - Kramers doublets 156
 - - molecular structure 158
 - - N–N distances 160
 - - parameters 156
 - - PELDOR 158
 - - β -phase crystals 146
 - - polymorphs 146
 - - spin–spin interaction 160
 - - structural instability 146
 - - χT product 155, 156
 - - temperature dependence 148
 - - Van Vleck equation 147
 - - X band spectrum 158
 - polycarbene 144
 - semiquinone radicals
 - - A–B pair 153
 - - A–D pair 153
 - - AFM coupling 154
 - - antiferromagnetic coupling 155
 - - asymmetric unit 152
 - - Brillouin function 152, 153
 - - DTBSQ 153
 - - EPR spectroscopy 153
 - - exchange mechanism 152
 - - magnetic measurement 153
 - - static deformation density 154
 - - structures 152
 - - π -symmetry 155
 - - valence tautomerism 152
 - thioradicals 147
 - paramagnetic chemical exchange saturation transfer (PARACEST) agents 418
 - paramagnetic Weiss temperature 71
 - Pascal's approach 69
 - Pc family
 - dI/dV mapping 315
 - Dy-Pc based devices 316
 - four-lobe molecule analysis 315
 - Kondo effect 316
 - Kondo temperature 316
 - Ln molecules 314
 - Ln(III)-Pc molecules 316
 - OFETs 314, 316
 - Tb(Pc)₂, SIM nature 316
 - TbPc₂ molecule 314
 - photon scanning microscopy (PSTM) 298
 - phthalocyanines (Pc) 17, 223, 224
 - aromatic rings 54
 - chromatography 53, 54
 - Ln(III)
 - - cube dodecahedron and squared antiprism 43
 - - energy splittings 44, 45
 - - Jz operator 43
 - - pyrrole nitrogen atom 42
 - - SAP 43
 - - structural parameters 43
 - porphyrins (RP) 53
 - sandwich compounds 54
 - physisorption 305
 - polyoxometalates (POMs) 17, 231, 234
 - angle dependent energy 45
 - characteristics 58
 - $e\lambda$ parameters 44
 - gadolinium 46
 - geometrical perturbation 45
 - Ising features 44
 - ligand field parameters 46
 - magnetic ions 56, 57
 - magnetic properties 44
 - sodium molybdate 56
 - spin localized mixed valence derivatives 57
 - spin-delocalized mixed valence derivatives 57
 - structures 57
 - Wybourne notation 47
 - pulsed electron-electron double resonance (PELDOR) 159
 - pyrazolylborates 17
 - trifluoromethanesulfonate 52
 - trigonal prismatic geometry 52, 53
 - trispyrazolylborate anion 52
 - pyrene moiety 324
- q**
- quantitative model 334
 - quantum computing (QC)
 - definition 329
 - detour 334
 - endohedral fullerenes 335
 - optical approach 331
 - philosophical aspects 347
 - QIP, *see* quantum information processing (QIP)
 - SQUID devices 330
 - transmon 330
 - 15 vanadium(IV) 346
 - quantum dots (QD) 297
 - quantum information processing (QIP)
 - bit 331
 - Bloch sphere 332
 - coherence and decoherence 333
 - definition 330
 - dephasing 338

- quantum information processing (QIP)
(contd.)
- electron and nuclear spins 332
 - entangled state 334
 - entanglement 333
 - inorganic compound 340
 - molecular rings
 - - anisotropy analysis 344
 - - 7 chromium(III) and one nickel(II) 342
 - - copper compounds 345
 - - Cr₇Ni moieties 345
 - - crystal K band spectrum 342
 - - 1D materials 341
 - - EPR and INS 343
 - - ESR and INS 342
 - - frozen solution 344
 - - giant spin model 343
 - - isotropic exchange interaction 342
 - - spin-Hamiltonian parameters 344
 - qubit 331, 333, 347
- r**
- radical pair model 334
 - remnant magnetization 200
- s**
- scanning tunneling microscopy (STM) 298
 - scanning tunneling potentiometry (STP) 298
 - Schiff bases 17
 - 3d-4f exchange interactions 62
 - homo and heterobinuclear compounds 64
 - primary imine 62
 - Schottky barrier 299
 - self-assembled monolayers (SAM) 312
 - self-consistent field (SCF) 84
 - sequence for phase error amplification (SPAM) 337
 - single chain magnets (SCM)
 - antiferromagnetic chain 251
 - chain-to-chain interaction 251
 - d and p orbital
 - - anisotropic Heisenberg Hamiltonian 259
 - - CoPhOMe 257
 - - crystal structure 258
 - - flipping energy 259
 - - isolated chain 258
 - - magnetic relaxation 258
 - - nitronyl nitroxides 257
 - - temperature dependence 258
 - diamagnetic defect 252
 - f orbitals, Dy
 - - ac measurements 269
 - - coordination environment 268
 - - CoPhOMe 268
 - - 1D system 271
 - - g tensor 269
 - - interchain dipolar interactions 267
 - - metamagnetic transition 271
 - - monomer and polynuclear systems 268
 - - puzzling result 269, 270
 - - reference frame 269
 - - supramolecular interactions 267
 - - versatility 269
 - f orbitals, Gd
 - - cell representation 262
 - - design 262
 - - Ising approach 264, 265
 - - magnetic properties 262
 - - NNN magnetic interactions 265
 - - paramagnetic phase 266
 - - pitch Q 265, 266
 - - $\chi;T$ values 263, 265
 - - temperature 263
 - Glauber model
 - - configurations 253
 - - dinuclear copper compound 255
 - - domain wall 256
 - - infinite array 255
 - - Orbach mechanism 254
 - - scaling function 256
 - - stochastic techniques 254
 - - temperature independent behavior 254
 - - time evolution 254
 - - magnetic properties 251
 - - non-collinear system 260–262
 - - spin correlation function 252
 - - spin dynamics 252
 - - spin glass 259, 260
 - - static and dynamic properties
 - - - anisotropic systems 272
 - - - 1D system 271
 - - - exponential divergence 272
 - - - f-p chains 273
 - - - magnetization 272
 - - - mixed metal chain 273
 - - - parameters 272
 - single ion magnet (SIM) 180
 - anionic double decker 224, 225
 - CF parameters 225, 226
 - magnetic site-dilution measurement 233
 - phthalocyanine 223, 224
 - POM 231, 232
 - resonant tunneling
 - - dipolar interaction 221
 - - features 221
 - - hyperfine constant values 221
 - - spectral resolution 219
 - ring-shaped aromatic ligands 233

- slow magnetic relaxation, Ho(III)
 - effective spin 218
 - hysteresis signals 219
 - irreducible representations 218
 - magnetic properties 218
 - micro SQUID 219
 - zero field 218
- Tb and Dy molecules
- didactic approach 227
- distortion angle 228
- electron-phonon interaction 228
- matrix element 227
- Orbach and Raman contributions 227
- properties 227
- relaxation times 228
- transverse components 222
- triple decker compounds 229, 230
- single-molecule magnets (SMMs) 18
 - ac susceptibility 243
 - Arrhenius plot 242
 - Arrhenius plot τ_0 207
 - Berry phase effects 197
 - blocking temperature 240
 - dinuclear compounds 245
 - energy quanta 196
 - $4f$ orbital 245
 - Fe4 structure
 - Arrhenius relaxation 205
 - H_3L ligands 202
 - helical pitch γ 203
 - Hilbert space 201
 - iron(III) clusters 201
 - isotropic and Zeeman components 204
 - local binary axes 204
 - low lying level 202
 - magnetic properties 203
 - Orbach model 203
 - quantum tunneling 205
 - robustness 205
 - trigonal symmetry 202
 - Jahn-Teller elongation axis 242
 - Kramers doublets 246
 - Kramers ion 243
 - magnetic dilution effects 210
 - magnetic hysteresis 196
 - metallacrown ligands 241
 - Mn12Ac analysis 196, 197
 - Orbach process 243
 - ovph complex 246
 - polynuclear system 245
 - QTM 213
 - quantum tunneling process 243
 - radical bridge 247, 248
 - spin topology 242
 - μ SQUID technique 245, 246
 - structures 239
 - superparamagnets
 - long range order 199
 - nanoparticles 200
 - Néel temperature 200
 - saturation value 200
 - short range order 199
 - superparamagnetism 201
 - thermal motion block 199
 - systematic approach 241
 - tunneling
 - equilibrium magnetization 208
 - low temperature relaxation 208
 - LZS model 208
 - Monte Carlo simulation 208
 - multispin model 209
 - propeller-like core 208
 - rhombic anisotropy 209
 - two-well model 197
 - uses 239
 - U_{eff} , Δ , τ_0 , and blocking temperature 198
 - valdien complexes 245
 - wavelength dependence 196
 - XMCD tool 211, 212
 - Y/Dy ratio 245
 - ZFS 195, 197, 198
- single-walled carbon nanotube (SWCN) 318
- SMM-SWCNT interaction 324
- spin crossover systems (SCO) 10
- spin dynamics 17
 - angular frequency 189
 - Cole-Cole plot 190
 - density 179
 - differential susceptibility 190
 - external and internal fields
 - dipolar interaction 187, 188
 - hyperfine interaction 187, 188
 - orbit-lattice parameters 189
 - in-phase susceptibility 190
 - isomorphous series 180
 - localized/delocalized ion 179
 - magnetization 189
 - measured susceptibility 190
 - out-phase susceptibility 190
 - phonons
 - continuum limit waves 182
 - Debye temperature 182
 - direct process 182, 183
 - frequency $h\nu$ 186
 - $Ho(EtS)_3$ 183
 - $Ho(EtS)_3$ 184
 - Kramers and non-Kramers ion 182
 - linear dependence 184

- spin dynamics (*contd.*)
 – – Orbach process 186
 – – orbit lattice coupling 183
 – – power dependence 185
 – – Raman process 185, 187
 – – relaxation rate 184
 – – relaxation time 186
 – relaxation process 181
 – SIM 180
 – spin lattice relaxation time T_1 , 181
 – spin–spin relaxation time T_2 , 181
 – – dipole–dipole interaction 191, 192
 – – Fermi’s golden rule 192
 – – Raman mechanism 192, 193
 – – Suhl–Nakamura model 191
 – τ distribution 190
 – τ distribution 191
 spin glass-like materials (SGL) 200
 spin Hamiltonian (SH) approach 74
 spin polarized scanning tunneling microscopy (SPSTM) 298
 spintronics 18
 state interaction (SI) matrix 134
 strong coupling 299
 superparamagnetism 200
 superposition model 40
- t**
 TbNPcPc double-decker molecules 317
 TbPc molecules 319
 TbPc2 molecules 319
 TbPc-pyrene 318
 temperature independent paramagnetism (TIP) 73
 tetracyanoethylene (TCNE) 150
 – 3d metal ions 165
 tetracyanoquinodimethane (TCNQ) 151
 – 3d metal ions 165
 tetra-phenyl-porphyrine (TPP) 149
 thioradicals 147
 time-of-flight secondary ion mass spectrometry (ToF-SIMS) 308
 time-resolved luminescent (TRL)
 immunoassays 432
 tunnel magnetoresistance (TMR) 297
- v**
 Van Vleck equation
 – anisotropy
 – – Bleaney scheme 81
 – – Dy(III) ion 79
 – – Ising type 74
 – – Jahn Teller effect 77
 – – lanthanide ion 79, 81
 – – non-degenerate wave function 76
 – – octahedral cobalt(II) complexes 78
 – – orbital moment 77
 – – spin orbit coupling 75
 – – tetragonal distortion 78
 – – unquenched orbital contribution 77
 – – Zeeman effect 76, 77
 – – zero-field splitting 76
 – aquo ion of Ce(III) 73
 – magnetic susceptibility 72
 – octahedral Co(III) compounds 73
 – spin Hamiltonian approach 74
 – TIP 73
 – Yb(III) energies and eigenfunctions 74
 vibronic coupling 136
 Vienna Ab initio Simulation Package (VASP) 319
- w**
 weak coupling 299
- x**
 X-ray absorption spectroscopy (XAS) 308
 X-ray based spectroscopies (XASs) 308
 X-ray magnetic circular dichroism (XMCD) 308, 310
 X-ray photoelectron spectroscopy (XPS) 308
- z**
 Zeeman effect 76, 77
 Zeeman perturbation 73
 Zero-field splitting (ZFS) 76, 89, 195, 411