

Index

a

- achiral hydrogenation
 - acetophenones 209–210
 - alkenes 209
 - arenes 213
 - ketones and aldehydes 209–212
 - levulinic acid 212
 - transfer hydrogenation 213–215
 - unsaturated polymers 212–213
 - water-soluble half-sandwich catalyst 214
- achiral monophosphine/Ru/chiral diamine-catalyzed asymmetric hydrogenation 168–171
- acid-base bifunctional heterogeneous catalysts
 - acidic and basic organic groups 2–7
 - coexistence, incompatible catalytic species 1
 - dual activation, electrophile and nucleophile 1, 2
 - nucleophilic addition reactions 1
 - OH H⁺ acid sites, silica-alumina surfaces 19
 - organic groups and acid sites 7–19
- acid-base interaction, solid surface
 - 1,4-addition reaction, nitroethane to methyl vinyl ketone 17
 - ¹³C CP/MAS NMR 15, 16
 - normalized intensity, ¹³C CP/MAS NMR 15, 16
 - SA(500) and triethylamine mixture 18
 - SA(500)-Net₂, preparation step 15, 16
 - silica-alumina 13
 - tertiary amines 13
- additives, reverse micelles
 - ionic liquids 521–525
 - nanomaterials 525–528
- alcohol oxidation
 - aliphatic diols 241, 242
 - amphiphilic resin particles 242–244
 - benzylic alcohols 241
 - HBr-H₂O₂ 242, 243
 - palladium(II) complex of sulfonated bathophenanthroline 239, 240
 - water-compatible oxidants 239
 - water-soluble Pd(II)-biquinoline 239
- alcohols, selective oxidation
 - O₂ by Au:PVP, activation 415
 - PVP-protected gold clusters 414
 - TOF values, Au 414
- Aldol and Mannich condensations, asymmetric
 - glycine imine ester with α-amino ester 449, 451
 - glycine imine ester with aldehyde 449, 450
 - synthesis of (+)-nemonapride 450, 451
- Aldol reaction
 - acyclic ketones 304
 - amphiphilic chiral organocatalyst 307
 - amphiphilic L-proline derivative 305–306
 - amphiphilic protonated diamine 304
 - asymmetric cascade reaction 309
 - bifunctional organocatalyst 304
 - cyclohexanone 305
 - direct cross-aldol reaction 306
 - dynamic light scattering 305
 - emulsions, optical micrograph 306
 - enantioselective direct aldehyde cross-aldol reaction 304
 - fluorescence microscope images 308
 - hexafluorophosphate salt 306–307
 - hydrophilic–lipophilic balance values 305
 - optically active isotetronic acids 307
 - organocatalytic reaction 303
- alkanes, selective oxidation 410–411
 - Au₅₅ and gold NPs 411
 - Au_{6–10} clusters 411–412

- alkanes, selective oxidation (*contd.*)
 - epoxidation of *trans*-stilbene 413
 - propylene epoxidation 413
 - propylene oxide (PO) selectivity 412
 - *in situ* UV-vis spectroscopy 412
 - *tert*-butyl hydroperoxide 411
 - Ti-OOH species 412–413
 - alkylation, SCFs
 - benzene 481
 - catalysts in scCO_2 481–482
 - isobutane and isopentane 481
 - alkylations, asymmetric
 - (–)-antofine 444
 - (–)-*cis*-clavicipitic acid 444–445
 - double alkylation of glycine imine ester 446
 - Maruoka-type chiral *N*-spiro quaternary ammonium salt 443–444
 - α -methyl serine derivatives 446
 - *N*-(diphenylmethylene) glycine *tert*-butyl ester 443
 - oxazoline system 446
 - synthesis of Selfotel 445
 - amide ligands 400, 403, 404
 - amine groups
 - amine-modified trialkoxysilane 92, 93
 - co-condensation method (one-pot synthesis) 91–92
 - functional groups 91, 92
 - grafting (post modification) 91
 - M41S materials 92
 - SBA-15 93
 - sol-gel processing 91
 - amine-catalyzed reactions
 - cyano-ethoxycarbonylation, carbonyl compounds 13, 14
 - Michael reaction 12, 13
 - SA- NEt_2 11, 12, 15
 - ^{29}Si MAS NMR analysis 12
 - tertiary amine immobilization, SA surface 11, 12
 - amphiphilic ligands 504
 - Anderson–Schulz–Flory (ASF) product distribution 560, 564, 585, 608
 - Anderson-type amphiphilic catalyst, 296
 - application, reverse micelles
 - biotransformation 538–541
 - enzyme immobilization 541–543
 - organic solvents/hydrophobic ILs 537
 - phase transition recovery 538
 - ASF product distribution. *See* Anderson–Schulz–Flory (ASF) product distribution
 - asymmetric aldol reaction 276–277
 - asymmetric carbonyl–ene reaction 178
 - asymmetric hydrogenation
 - acetophenone 215
 - aromatic ketones 216–217
 - aryl ketones 217
 - cyclic imines and iminium salts 217, 218
 - hydrophilic BINAP derivatives 215–216
 - methyl 2-acetamidoacrylate reduction 215
 - nitroalkenes 218–219
 - Noyori-Ikariya type catalysts 216
 - asymmetric Michael addition reaction 278
 - asymmetric PTC, organic synthesis
 - Aldol and Mannich condensations. *See* Aldol and Mannich condensations, asymmetric
 - asymmetric alkylations. *See* Alkylations, asymmetric
 - chiral anion PTC 441–442
 - chiral cation. *See* Chiral cation PTC
 - chiral crown ethers. *See* Chiral crown ethers, PTC
 - conjugate additions 447–448
 - cyanation reactions 459
 - cyclization reactions 457–458
 - 1,3-dicarbonyl derivatives 450–453
 - fluorination and trifluoromethylation reactions 458–459
 - glycine imine derivatives 443–446
 - hydrolysis 460
 - Neber-rearrangement 460
 - nitroalkanes 455, 457
 - oxindoles 454–455
 - ring-opening reaction 460
 - asymmetric Suzuki–Miyaura coupling reactions 142, 145
- b**
- BEP principle. *See* Brønsted–Evans–Polanyi (BEP) principle
 - bifunctional catalysts. *See* Acid-base bifunctional heterogeneous catalysts
 - BINAP (2,2-O-bis(diphenylphosphino)-1,1'-binaphthyl) derivatives 215–216
 - 1,1'-Bi-2-naphthol (BINOL), 426
 - BINOLate/La(III)-catalyzed asymmetric epoxidation
 - bridging spacers, multitopic ligands 181, 182
 - heterogenized catalysts, chalcone epoxidation 181
 - α,β -unsaturated ketones, enantioselective epoxidation 182, 183
 - BINOLate/Ti-catalyzed asymmetric carbonyl–ene reaction 156–158

- BINOLate/Ti(IV)-catalyzed asymmetric carbonyl–ene reaction
 - (R)-BINOL–Ti(OiPr)₂/(R)-BINOL complex 178
 - chiral titanium-bis-BINOL polymers 178
 - enantioselective carbonyl–ene reaction 178, 180
 - generation 178, 179
- BINOLate/Ti(IV)-catalyzed asymmetric hetero-Diels–Alder reaction 154–156
- BINOLate/Ti(IV)-catalyzed asymmetric sulfoxidation reaction
 - enantioselective sulfoxidation 179, 181
 - ICP-AES analyses 179
 - linker-bridged BINOL ligands 178
 - self-supported Ti catalysts 179, 180
- BINOLate/Zn(II)-catalyzed asymmetric epoxidation 183–184
- bromide poisoning of active sites, Au/TiO₂ catalysts
 - CO oxidation reactions 344–345
 - NaBr poison 344, 345
 - WGS reaction 345–346
 - XANES measurements 344
- Brønsted–Evans–Polanyi (BEP) principle 559, 586, 587
- C**
- carbide mechanism
 - definition 555
 - initiation 557–558
 - propagation 558–559
 - relative rate requirements 556, 557
 - termination 559
- carbon nanotubes (CNTs) 352
- catalysis. *See* Supercritical fluids (SCFs), catalysis
- catalyst recycling
 - “catalyst in a tea bag” strategy 136
 - heterogeneous catalysis 111
 - research, chiral molecular complexes 111
- catalytic acetalization reaction 279
- catalytic asymmetric fluorination reaction 280–281
- catalytic carbonylation reaction
 - activation CO₂ 70
 - with CO 68–70
 - hydroformylation reactions. *See* Hydroformylation reactions
- catalytic condensation reaction 279–280
- catalytic hydrogenation
 - alkyne 58–59
 - olefin 58
 - α,β -unsaturated aldehydes 60–61
- catalytic nanoreactor engineering
 - homogeneous and heterogeneous properties 390
 - macroreactors and microreactors 390
 - porous materials 390
- catalytic oxidation reaction 278–279
- catalytic reactions, RTILs. *See* Room-temperature ionic liquids (RTILs)
- C–C bond formation
 - alcohol oxidation 238–244
 - Diels–Alder reaction 220–224
 - Friedel–Crafts alkylation 224–226
 - Heck reaction 234–238
 - Suzuki–Miyaura coupling 226–234
- C–C coupling reactions
 - Mizoroki–Heck reaction 45–47
 - Sonogashira reaction 48
 - Stille reaction 47
 - Suzuki reaction 42–44
 - Ullmann reaction 48–49
- cetyltrimethylammonium bromide (CTAB) 500–502
- cetyltrimethylammonium chloride (CTAC) 205
- chemical reduction, gold clusters
 - amide ligands 400–404
 - gold NPs, formation 401
 - LaMer model 399–400
 - organic ligands 400, 401
 - phosphorus ligands 400–402
 - sulfur ligands 400–403
- chiral anion PTC
 - “anion metathesis” 442
 - chiral phosphate salt 442
 - C₈-TRIP 63b and PhDAP 442
 - silver salt of TRIP 63a 442
- chiral binaphthyl and biphenyl, quaternary ammonium salts
 - alkyl groups 434, 436
 - asymmetric Darzens reaction 438
 - conformational rigidity and flexibility 438
 - C₂-symmetric BINOL 434, 435
 - free OH group 437, 438
 - N-(diphenylmethylene) glycine tert-butyl ester, alkylation 434, 435
- chiral catalysis
 - asymmetric catalysis 105
 - (R)-(+)-BINOL 105
 - chiral bis(cyclohexyldiamine)-based Ni(II) complexes 106
 - chiral bis(oxazoline) ligand 106
 - chiral vanadyl salen complex 106
 - heterogeneous catalyst 105
 - mesoporous silica, L-proline 105

- chiral catalysis (*contd.*)
 - Rh and Ru catalysts 106
 - SBA-15 105
- chiral cation PTC
 - chiral quaternary ammonium salts 428–440
 - chiral quaternary phosphonium salts 440–441
- chiral crown ethers, PTC
 - BINOL 426
 - (+)-camphor-based chiral aza-crown ether 428
 - chiral spiro crown ether 426–427
 - C₂-symmetric chiral crown ethers 427
 - D-glucose-based crown ethers 442
- chiral dendritic catalysts
 - advantages, polymer resins 126
 - BINAP ligands, Ru-catalyzed asymmetric hydrogenation 127
 - carbosilane dendrimers 136
 - catalysis, core and periphery 126
 - diamine ligands, Ru-catalyzed asymmetric transfer hydrogenation 131–132
 - folded prolinamide, asymmetric aldol reactions 137, 139
 - modular chiral monodentate phosphoramidite ligands 128–130
 - modular PHOX ligands 130, 131
 - NOBIN ligands, Ti-catalyzed hetero-Diels-Alder reactions 132, 133
 - PAMAM-bound [Co(salen)] catalysts, epoxides 133, 134
 - phosphorus dendrimer-bound phosphoramidites 137, 138
 - phosphorus dendrimer-immobilized azabis(oxazoline) ligands 137
 - PPI/PAMAM-bound diphosphine ligands 133, 134
 - PPI-supported BINAP, Cu-catalyzed asymmetric hydrosilylation 135
 - quinoline derivatives, BINAP ligands 127, 128
 - Ru-catalyzed and thermomorphic asymmetric hydrogenation 128, 129
 - supramolecular monophosphite ligands 130, 131
 - TsDPEN-functionalized Janus dendrimers 138, 140
- chiral guanidinium salts 438–439
- chiral linear polymeric catalysts
 - bis(oxazoline)-based coordination polymers 125
 - covalent binding strategy 123
 - cross-linked insoluble polymers 114
 - diethyl zinc, aldehydes 120, 121
 - main-chain quaternary ammonium cinchonidine salts 124, 125
 - PEG-bounded TsDPEN 117
 - PEG-supported chiral monophosphite 117, 118
 - and PEG-supported ligands 116
 - PEG-supported Ru-BINAP catalyst 118
 - PEG-supported tartaric acid ester ligands 116, 117
 - polyester-supported BINAP ligands 121, 123
 - quaternary ammonium cinchonidine salts, asymmetric allylation 124
 - salen catalysts, olefins asymmetric epoxidation 123
 - soluble polymer-supported salen (Co/Mn) 119
 - structure, BINOL-BINAP copolymer 121, 122
 - water-soluble and pH-sensitive pendant copolymer, aldol reactions. 119, 120
- chiral onium salts 428
- chiral quaternary phosphonium salts 440–441
- 3-(chloropropyl)triethoxysilane (CPTES) 95
- cinchona, quaternary ammonium salts
 - bifunctional quaternary ammonium salt 430–431
 - cinchonine- and cinchonidine 428, 429
 - dihydrocinchonidine- and quinidine 429
 - dimeric and trimeric cinchona 433
 - 9-fluorinated quinidine-derived catalyst 432
 - free 9-OH group 431
 - N-anthracenylmethyl function 428
 - N-2',5'-bis(trifluoromethyl)benzyl quinidinium salt 430
 - N-(diphenylmethylene) glycine tert-butyl ester, alkylation 428
 - N-p-trifluoromethylbenzylcinchoninium bromide 428
 - O-methyl chinconidinium salt 430
 - phenolic OH group 431
 - quinine-derived quaternary ammonium salt 428
 - steric properties 429–430
 - (thio)urea H-bonding donors 432–433
- clustering 471
- CO binding, imaging
 - edge sites 338–339
 - electron richness of gold 337–338
 - infrared spectra, gold clusters 338, 339
 - oxidation 337

- STM techniques 338, 339
 - structure-sensitive 337
 - CO₂ cycloaddition to epoxide, SCFs
 - cross-linked-polymer-supported ionic liquid 483
 - cyclic carbonates 482
 - glycerol 483
 - propylene carbonate (PC) production 483
 - scCO₂-soluble catalysts 482
 - styrene carbonate 483
 - CO₂, hydrogenation
 - to formic acid 477
 - in methanol 477
 - *N,N*-diethylformamide production 477–478
 - scCO₂ and scCO₂-IL systems 476–477
 - scCO₂/[bmim][PF₆] system 477
 - CO oxidation
 - Au₈ clusters 418
 - Au₁₃ clusters 415
 - Au/Mg(OH)₂ 415, 416
 - Au/TiO₂ 417
 - MgO film 415, 416, 417
 - moisture 418
 - TOF 417–418
 - Co-condensation method (one-pot synthesis)
 - amine groups 91–92
 - organic–inorganic hybrid materials 88–89, 91–96
 - COFs. *See* Covalent organic frameworks (COFs)
 - combinatorial approach, homogeneous
 - asymmetric catalysis. *See* Homogeneous asymmetric catalysis
 - conformation, reverse micelles 528–530
 - conjugate additions
 - glycine imine esters 447, 448
 - synthesis of (+)-cylindricine C 448
 - synthesis of (+)-monomarine 448, 449
 - cooperative activation effect
 - epoxide hydration 381–382
 - isolation effect 382–385
 - kinetic resolution of epoxides 377–379
 - water oxidation reactions 380
 - cooperative catalysis
 - nitroaldol reactions 18, 19
 - *p*-formylbenzoic acid with nitromethane reaction 19
 - SA-NH₂-NEt₂. surface structure 18
 - coordinatively unsaturated (undersaturated) sites (CUS) sites 343
 - covalent bonded bridging ligands
 - continuous-flow reactor 188, 190
 - heterogeneous catalysis, enantioselective hydrogenations 188, 189
 - ICP-AES analysis 188
 - monodentate phosphorus ligands 187–188
 - covalent organic frameworks (COFs) 352
 - covalent-bonding methods
 - asymmetric catalysis 360
 - chiral ligand/catalyst 359–360, 362
 - chiral MOFs 360–362
 - chiral PMOs 360
 - D-POST-1 and L-POST-1 360
 - photocatalyst and chiral catalyst 360
 - post-synthetic modification 362–363
 - CTAB. *See* Cetyltrimethylammonium bromide (CTAB)
 - cyanation reactions
 - electron-deficient alkenic acceptors 459, 461
 - Strecker reaction 459, 460
 - cyclization reactions
 - asymmetric PTC synthesis 457–458
 - clusiaphenone B 457
 - synthesis of (-)-hyperibone K 457, 458
 - synthesis of isoxazolines and pyrazolines 458
- d**
- 1-decene 206, 207
 - density functional theory (DFT) analysis 335–336
 - DFT. *See* Density functional theory (DFT) analysis
 - 1,3-dicarbonyl derivatives
 - asymmetric PTC 450–452
 - conjugate addition of dimethyl malonate 452
 - synthesis of (-)-paroxetine and (-)-isonitramine 452, 453
 - Diels–Alder reaction
 - asymmetric 224
 - 2-cyclohexen-1-one and benzaldimines 221, 223
 - cyclopentadiene with buten-2-one 220–221
 - cyclopentadiene with enones 221, 223
 - diene carboxylate salts 221, 222
 - 2,4-hexadienyl acetate cycloaddition 224, 225
 - 2,6-dimethyl- β -cyclodextrin (DMCD) 504
 - DLCs. *See* Double long-chain cationic surfactants (DLCs)
 - DMCD. *See* 2,6-Dimethyl- β -cyclodextrin (DMCD)
 - 1-dodecene 205

- double long-chain cationic surfactants (DLCSS) 502
- dual reaction center site model, FT reaction
 - advantages 593
 - calculation, kinetics 592–593
 - expressions, α and CO consumption 594
 - generation, reducible oxides 594
- e**
- edge X-ray absorption fine structure (EXAFS) measurements 346
- electrospray ionization (ESI) 405
- emulsion catalysis
 - amphiphilic catalyst 286
 - Bancroft rule 285
 - DBT removal vs. reaction time 297
 - definition 285–287
 - emulsion droplets, optical photographs 297
 - H_2O_2 as oxidant. *See* Oxidative desulfurization (ODS), H_2O_2 as oxidant
 - incompatible reactants 284
 - liquid–liquid biphasic reaction system 287
 - micelles, macroemulsion and emulsion 284
 - model 286
 - O_2 as oxidant 296–297
 - organocatalysts. *See* Organocatalysts
 - O/W and W/O surfactant emulsifier 285
 - polymer-bounded catalysts. *See* Polymer-bounded catalysts
 - water as solvent 283–285
 - water in chemistry 283
- emulsion/sol-gel transport system (EST) 208
- enzyme catalysis
 - advantages 511
 - asymmetric acylation 1-phenylethanol 77
 - *Candida rugosa* lipase 77
 - desorption experiments 77, 78
 - disadvantages 511
 - DKR 1-phenylethanol, 78, 79
 - immobilization, lipase 77
 - kinetic resolution, aromatic secondary alcohols 78
 - molecular organic solvent-based reverse micelles. *See* Molecular organic solvent-based reverse micelles
 - reusability, penicillin G acylase 79
- epoxide hydration, nanoreceptors
 - DFT calculation 381
 - $\text{H}_2\text{O}/\text{EO}$ ratio 381
 - S/C ratio 382
- EXAFS. *See* Edge X-ray absorption fine structure (EXAFS) measurements
- extended x-ray absorption fine structure spectroscopy (EXAFS) 405, 406
- f**
- Fischer–Tropsch (FT) reaction
 - analysis, selectivity and conversion 559–560
 - behavior, chain growth parameter 601, 602
 - carbide mechanism. *See* Carbide mechanism
 - cutoff behavior, parameter set 560, 561
 - equilibrium distribution, self-assembling system 564
 - expression, methane selectivity 563, 599, 600
 - expressions, R_{CO} and α 599, 600
 - incorporation, reversible C–C bond formation 560
 - infinite chain length 560, 562
 - kinetic Monte Carlo approach 554
 - lumped kinetics model. *See* Lumped kinetics model, FT reaction
 - methane formation 603, 604
 - modeling 553
 - molecular microkinetics simulations. *See* Molecular microkinetics simulations, FT reaction
 - monomer formation limited kinetics vs. chain growth limit 561
 - Pichler–Schulz mechanism 555
 - reverse isotope effect 603
 - Sabatier principle behavior, R_{CO} 601
 - simulated values, α and R_{CO} 561, 563
 - transient kinetics. *See* Transient kinetics, FT reaction
- fluorinated dendrimer
 - fluorous HPLC separation 259
 - fluorous mixture synthesis 262–263
 - G2 260
- fluorinated diphosphine ligand (dfppp) complexes 266
- fluorous aminosulfonamide-catalyzed aldol reaction 276–277
- fluorous catalysis
 - benzotrifluoride 254
 - dendrimer. *See* Fluorinated dendrimer
 - ^{19}F MRI (magnetic resonance imaging) agent 258
 - fluorine content differences 258
 - fluorous molecule, structure 254
 - fluorous tags 254
 - FMS, solution-phase reaction 257–258
 - G0–G3 dendrimers 261
 - history 253–254

- HPLC separation 256–257
 - intermediates synthesis 263
 - metallic catalysis. *See* Fluorous metallic catalysis
 - Mitsunobu reaction mixture 255
 - organocatalysis. *See* Fluorous organocatalysis
 - Pd-catalyzed hydrogenolysis 261
 - separation methods 254–255
 - solid-phase extraction 256
 - fluorous Grubbs' catalysts 272–273
 - fluorous hydrazine-carbothioate-catalyzed reaction 280
 - fluorous immobilized nanoparticles catalysts
 - acridines 269
 - Diels–Alder reaction 267–268
 - fluorous alkyl iron(III) porphyrin 268
 - fluorous nanoparticles and related reaction 269
 - FSG.Hf(N(SO₂C₈F₁₇)₂)₂-catalyzed reactions 269–270
 - FSG-supported copper(II) nanoparticles 268
 - FSG-supported palladium nanoparticles 267–268
 - nano-palladium catalyst 267
 - olefin oxidation 268
 - perfluorinated bis(dihydrooxazole) copper(II) (PBDC) complexes 267–268
 - fluorous metallic catalysis
 - fluorinated diphosphine ligand (dfppp) complexes 266
 - fluorinated ytterbium-catalyzed condensation 275
 - fluorous Grubbs' catalysts 272–273
 - fluorous immobilized nanoparticles catalysts 267–270
 - fluorous palladacycle catalysts 264–265
 - fluorous palladium-NHC complexes 270–271
 - fluorous phosphine-based palladium catalyst 271–272
 - fluorous pincer ligand-based catalysts 265–266
 - fluorous silver catalyst 273
 - fluorous trizole-based ligand 274
 - fluorous Wilkinson catalyst 273–274
 - immobilization 264
 - perfluoroalkylated PCP pincer ligand 266
 - uses 263
 - fluorous *o*-iodoxybenzoic acid-catalyzed oxidation reaction 279
 - fluorous organocatalysis
 - asymmetric Aldol reaction 276–277
 - asymmetric Michael addition reaction 278
 - catalytic acetalization reaction 279
 - catalytic asymmetric fluorination reaction 280–281
 - catalytic condensation reaction 279–280
 - catalytic oxidation reaction 278–279
 - fluorous aminosulfonamide-catalyzed aldol reaction 276–277
 - Morita–Baylis–Hillman reaction 277–278
 - fluorous palladium-NHC complexes
 - dimethylformamide (DMF) 271
 - *N*-heterocyclic carbenes (NHCs) 270
 - Suzuki coupling 270–271
 - fluorous phosphine-based palladium catalyst 271–272
 - fluorous thiourea-catalyzed oxidation reaction 279
 - Fourier transform infrared (FTIR) 493
 - Fréchet-type dendritic catalyst 128–129, 131–132
 - Friedel–Crafts alkylation
 - 2-acyl imidazoles 226, 227
 - aldehydes, amines, and indoles 225
 - C-Glycosylation 226
 - nucleophilic substitution of alcohols 226, 228
 - 3-substituted indoles 224–225
 - FT reaction. *See* Fischer–Tropsch (FT) reaction
 - FTIR. *See* Fourier transform infrared (FTIR)
- g**
- gold clusters, heterogeneous catalysis
 - Au₃₀₉ clusters 406
 - characterization 405–406
 - chemical reduction. *See* Chemical reduction, gold clusters
 - d-Band spin-orbital splitting 397, 398
 - definition 397
 - electrical reduction 404
 - features 397
 - vs. gold NPs 398
 - organic synthesis 419–420
 - physical vapor deposition 403
 - preparation 399–405
 - selective hydrogenation. *See* Selective hydrogenation
 - selective oxidation. *See* Selective oxidation
 - surface states and valence band structures 397, 398
 - titanosilicalite-1 (TS-1) 405
 - grafting (post modification) method
 - amine groups 91

- grafting (post modification) method (*contd.*)
 - organic–inorganic hybrid materials 88, 91–96
- green solvents, 469. *See also* Supercritical fluids (SCFs), catalysis

h

- HAADF-STEM. *See* High-angle annular dark-field scanning transmission electron microscopy (HAADF-STEM)
- Heck reaction
 - aryl halides with olefins 235
 - aryl iodides with olefins 236–238
 - *m*-triphenylphosphane-3,3',3''-tricarboxylate, 236
 - nonionic amphiphile 235–236
 - oxime-based ligands 238
 - prolinol vinyl ether 235
 - *ris*(4,6-dimethyl-3-sulfonatophenyl)-phosphane trisodium salt 236
 - *tert*-butyl acrylate/styrenes 234
- helical polymeric catalysts
 - asymmetric hydrosilylation 142, 144
 - asymmetric Suzuki–Miyaura coupling reactions 142, 145
 - chiral phosphorus ligand 140, 143
 - synthesis, ligands and applications 140, 142
- heterogeneous asymmetric catalysis, self-supporting approach
 - asymmetric carbonyl–ene reaction 178
 - BINOLate/La(III)-catalyzed asymmetric epoxidation 180–183
 - BINOLate/Ti(IV)-catalyzed asymmetric carbonyl–ene reaction 178
 - BINOLate/Ti(IV)-catalyzed asymmetric sulfoxidation reaction 178–180
 - BINOLate/Zn(II)-catalyzed asymmetric epoxidation 183–184
 - heterogenization, chiral catalysts 175, 176
 - homochiral microporous metal-organic materials 175
 - immobilization, chiral catalysts 172–173, 174
 - metal-organic coordination polymers 174
 - metal-organic polymers 175–176
 - modes, heterogenization of homogeneous catalyst 176, 177
 - MonoPhos/Rh(I)-catalyzed asymmetric hydrogenation reactions 187–194
 - Noyori-type Ru(II)-catalyzed asymmetric hydrogenation 185–187
 - polymeric ligands 173
- heterogeneous catalysis
 - catalyst development pathway 90, 91
 - description 89
 - and homogeneous catalysis, characteristics 89–90
- heterogeneous catalysis with homogeneous performance
 - alkylation and esterification reactions 73–74
 - asymmetric catalysis 74–77
 - catalytic carbonylation reaction. *See* Catalytic carbonylation reaction
 - catalytic hydrogenation 57–61
 - enzyme catalysis 77–79
 - IL-phase catalysis 51–55
 - isomerization and oligomerization 72–73
 - selective oxidation 61–63
 - silica gel-confined IL catalysts 55–57
 - water-gas shift reaction 70–72
- heterogeneous catalytic reactions
 - hydrogenation of CO₂ 473–474
 - SCFs reactions 472
- 1-Hexene 203–204
- high-angle annular dark-field scanning transmission electron microscopy (HAADF-STEM), 342, 405–406
- higher olefins, hydroformylation
 - aqueous-organic biphasic system 497
 - aqueous-phase catalysts 498–499
 - cosolvent 499–500
 - cyclodextrins 503–505
 - RCH/RP process 497–498
 - surfactants 500–503
 - TRPTC 505–506
 - unmodified Rh-TPPTS system 498
 - water-soluble substrates 498
- high-resolution transmission electron microscopy (HRTEM) 328
- homogeneous asymmetric catalysis
 - catalyst separation methods 114, 115
 - chiral catalyst discovery 152
 - chiral dendritic catalysts. *See* Chiral dendritic catalysts
 - chiral linear polymeric catalysts. *See* Chiral linear polymeric catalysts
 - helical polymeric catalysts. *See* Helical polymeric catalysts
 - “one-phase catalysis/two-phase separation” 112
 - Ru complex-catalyzed enantioselective reactions 168–172
 - soluble polymeric supports 112–113
 - Ti(IV)-catalyzed enantioselective reactions 153–159

- Zn complex-catalyzed enantioselective reactions. *See* Zn complex-catalyzed enantioselective reactions
 - homogeneous catalytic reactions
 - CO₂ Cycloaddition to Epoxide 483
 - hydroformylation reactions 478–479
 - hydrogenation of CO₂ 477
 - organic reactions 475
 - SCFs reactions 472
 - HPMO. *See* Hybrid periodic mesoporous organosilica (HPMO)
 - HRTEM. *See* High-resolution transmission electron microscopy (HRTEM)
 - hybrid periodic mesoporous organosilica (HPMO)
 - organic–inorganic hybrid materials 89
 - PMO 98–99
 - hydroformylation
 - alkenes 202–203
 - aqueous-biphasic hydroformylation 205
 - aryl phosphines 489
 - atom-economical and clean reaction 489
 - biphasic system 508
 - cationic surfactants 205–206
 - cyclodextrin 206–207
 - data 489, 490
 - 1-hexene and styrene 204–205
 - higher olefins. *See* Higher olefins, hydroformylation
 - immobilizing catalysts 207
 - internal olefins 506–508
 - lower olefins. *See* Lower olefins, hydroformylation
 - mechanism 491–493
 - methyl 3-pentenoate 204
 - nonionic surfactants 206, 207
 - olefins 203–204, 207–208, 489, 490
 - “OXO” process 489
 - RCH/RP process 489–490
 - rhodium-phosphine complex catalytic process 489
 - switchable additives 208
 - technical and economic advantages 508
 - water-miscible organic cosolvent 205
 - water-soluble rhodium-phosphine complex catalytic systems 490–491
 - hydroformylation reactions
 - alkenes 479–480
 - C₄ olefins 63–65
 - C₂–C₂ olefins 65–67
 - 1-hexene 480
 - 1-octene 480
 - olefin 479
 - propylene 479
 - unsaturated alcohol and ester 67–68
 - hydrogenation
 - achiral hydrogenation 209–215
 - asymmetric hydrogenation 215–220
 - C–C bond formation. *See* C–C bond formation
 - gaseous hydrogen 208
 - reactions 31–32
 - hydrogen-bonded bridging ligands
 - homochiral supramolecular metal-organic assembly 190, 191
 - ICP-AES spectroscopic analyses 192
 - Ureido-4[1*H*]-ureidopyrimidone 190
- i**
- ILs. *See* Ionic liquids (ILs)
 - IL-supported/mediated metal nanoparticles
 - C–C coupling reactions. *See* C–C coupling reactions
 - hydrogenation reactions 31–32
 - molecular dynamics simulations 30–31
 - nanocatalysis 22
 - NMPs preparation, ILs. *See* NMPs preparation, ILs
 - Pd NPs 32, 34–35
 - Pt and Ir NPs 36–37
 - Ru NPs. *See* Ru NPs, IL-supported
 - SEM and TEM 29–30
 - XPS and NMR 28–29
 - immobilized catalyst separation methods 114, 115
 - immobilized chiral catalysts. *See* Homogeneous asymmetric catalysis
 - In situ* infrared spectroscopy, 346
 - ionic liquid-based reverse micelles
 - microemulsification 531–537
 - surfactants 537
 - ionic liquids (ILs). *See also* Room-temperature ionic liquids (RTILs)
 - BPy-functionalized ILs 41, 42
 - CN-functionalized 32
 - “control performance” 22
 - “heterogenized (ionic) liquids” 94
 - IL-supported/mediated NMPs. *See* IL-supported/mediated metal nanoparticles
 - immobilization via anion 94, 95
 - immobilization via cation 94, 95
 - immobilization via supported liquid phase 94, 95
 - Knoevenagel reactions 93
 - MCM-41 93
 - molecular dynamics simulations 30–31

- ionic liquids (ILs). *See also* Room-temperature ionic liquids (RTILs) (*contd.*)
 - NMPs preparation. *See* NMPs preparation, ILs
 - organic–inorganic hybrid materials 93–95
 - SBA-15 93
 - Sonogashira reaction 48
 - Stille reaction 47
 - and supercritical carbon dioxide 34–35
 - Suzuki reaction 42
 - Ullmann reaction 48
- iridium clusters, ethylene hydrogenation
 - MgO-supported clusters 348
 - structure–function relationships 346
 - Wilkinson’s catalyst $\text{Ir}(\text{PPh}_3)_3\text{Cl}$ 347
 - zeolite-supported catalyst 346, 347, 348
- isolation effect, nanoreceptors
 - alkenes, DCA- and HA-photosensitized oxidation 383, 384
 - molecular catalysts, dimerization 385–386
 - phenyl phenylacetates, structure 384
 - selective oxidation 383
 - selectivity control 382–385
 - zeolites, shape selectivity 382–383

k

- kinetic Monte Carlo model, FT reaction 554, 564–565, 567
- kinetic resolution (KR), epoxides
 - chiral Co(Salen) catalyst 378
 - Cr(Salen)py, concentration 379
 - DFT calculation 379
 - metal Salen complexes 377–378
 - nanocages, SBA-16 379
- Knoevenagel condensation
 - ILs 93
 - organic–inorganic hybrid materials 99, 103
- KR. *See* Kinetic resolution (KR), epoxides

l

- LASC. *See* Lewis acid surfactant combined catalyst (LASC)
- Lewis acid surfactant combined catalyst (LASC) 298–302
- Lewis acid-catalyzed organic reactions
 - aldol reactions, asymmetric version 301
 - α -amino phosphonate 299
 - asymmetric aldol reactions 302
 - LASC 12 and benzaldehyde mixtures 299
 - LASC-catalyzed aldol reaction 298
 - LASC-catalyzed allylation reactions 301
 - LASC-catalyzed Friedlnder annulation 302

- meso-epoxides, asymmetric ring opening 303
 - Michael reactions 299
 - scandium–bipyridine complex 301–303
 - Sc(DS)₃-catalyzed three-component 300
 - tetraallyltin 301
 - tetrahydrofuran (THF) 297
 - lignin peroxidase (LiP)
 - AOT head group 513
 - catalytic activities 513, 514
 - GGDE/TX-100 and AOT reverse 515
 - reverse micelles 540
 - LiP. *See* Lignin peroxidase (LiP)
 - lower olefins, hydroformylation
 - butene 496–497
 - description 493
 - ethylene 493–494
 - propene 494–496
 - L-proline-functionalized mesoporous silica, direct synthesis 95–96
 - lumped kinetics model, FT reaction
 - definition 554
 - dual reaction center site 592–594
 - single reaction center site. *See* Single reaction center site model, FT reaction
- m**
- MALDI. *See* Matrix-assisted laser desorption ionization (MALDI)
 - matrix-assisted laser desorption ionization (MALDI) 405
 - MBG. *See* Microemulsion-based organogel (MBG)
 - mesoporous silicas (MSs)
 - co-condensation 356–358
 - grafting method 355–357
 - hydrothermal method 353
 - periodic mesoporous organosilicas 357
 - PMOs 354
 - SAB-16 and FDU-12 353–354
 - SBA-15 353
 - surface modification 355–358
 - metal cluster catalysts
 - bromide poisoning of active sites, Au/TiO₂ catalysts 344–346
 - catalyst pretreatment temperature 328
 - CH₄ turnover rates 326
 - chromia gel pretreatment temperature 327
 - CO binding, imaging 337–339
 - copper cluster cores 329
 - electrochemical reduction of CO₂ 328
 - enzymes 329
 - Fe-S clusters 331
 - g-Al₂O₃ system 328

- gold core 329, 330
- Ir₄ clusters. *See* Iridium clusters, ethylene hydrogenation
- metal-carbonyl clusters via ligand effects. *See* Metal-carbonyl clusters via ligand effects
- 2-NT probe molecule 330–331
- open sites. *See* Open catalytic active sites
- structure-sensitive or insensitive reactions 325–326
- “varying degrees of saturation” 325
- well-characterized systems 331
- metal-carbonyl clusters via ligand effects
- Blyholder model of metal-CO bonding 333–334
- calix[4]arene phosphine (L) 332, 333
- CO and O binding, gold clusters 336
- DFT analysis 335–336
- gold binding energy 332
- Ir₄ carbonyl clusters 332–337
- Pauli repulsion 336
- π -backbonding interaction 334
- single-crystal X-ray diffraction 334–335
- metal-coordinated bridging ligands
- α -dehydroamino acid esters, asymmetric hydrogenation 192
- heterogeneous asymmetric hydrogenations 192, 194
- homochiral bimetallic coordination assemblies 192, 193
- metal-organic frameworks (MOFs)
- covalent bonding 358
- dative bonding 358–359
- definition 354
- functionalization 358
- porous structure, control 354
- post-synthetic modification 358
- structural and chemical properties 355
- surface modification 358–359
- methane formation *vs.* FT kinetics
- lumped microkinetics expression, production 584
- rate constants 583
- temperature maximum, C_n production 585
- volcano curve, production 584
- methyl 3-pentenoate, hydroformylation 204
- Michael addition, organocatalysts
- aldehydes 310
- diamine organocatalyst 309
- malonate 312
- nitroalkenes 309
- nitroolefins 310
- β -nitrostyrene 310
- Rolipram 310
- SATO catalysts 310
- surfactant-type asymmetric organocatalysts 310
- α,β -unsaturated aldehydes 311
- microemulsion-based organogel (MBG)
- ethyl valerate 542
- IL-based 543
- immobilization 541, 543
- microstructure 542
- phenylethyl acetate 541
- microenvironment engineering, nanoreactors
- Bu₄NBr, addition 387
- (1R,2R)-diaminocyclohexane 386
- diffusion of reactants and products 387–388
- factors influencing 385–386
- NaCOO-H₂O system 387
- Ru-TSDPEN 386
- surface modification 386
- TOF of Fe(salan) 387
- Mizoroki–Heck reaction
- iodobenzene and ethyl acrylate reaction 45
- Pd NPs 46, 47
- Pd/chitosan nanocomposite 45, 46
- MOFs. *See* Metal-organic frameworks (MOFs)
- molecular catalysts, nanoreceptors
- covalent-bonding methods 359–363
- noncovalent bonding methods 363–366
- reducing pore entrance size 366–369
- molecular dynamics simulations
- DFT methods 30
- Ru NPs 31, 33
- spatial distribution function, imidazolium cation 31, 33
- molecular microkinetics simulations, FT reaction
- chain growth limit 569, 573
- chain length cutoffs 567, 570
- elementary reaction rate data 565
- H₂ and CO partial pressure dependence 576–578
- high C₂+ yield 574, 575
- *vs.* methane formation. *See* Methane formation *vs.* FT kinetics
- oxygenate formation 571
- polymerization *vs.* chain growth model. *See* Polymerization *vs.* chain growth model, FT reaction
- reaction energy, propylene and methane formation 565–567
- relative rate, CH_x formation 567, 572
- simulated conversion data, parameters 567, 568

- molecular microkinetics simulations, FT reaction (*contd.*)
 - surface composition 574
 - temperature dependence, α 567, 570
 - value, α convergence 567, 571
 - molecular organic solvent-based reverse micelles
 - electrical property 513–516
 - hydrophobic group 513
 - surfactant head group 516–521
 - MonoPhos/Rh(I)-catalyzed asymmetric hydrogenation reactions
 - covalent bonded bridging ligands 187–190
 - hydrogen-bonded bridging ligands 190–192
 - metal-coordinated bridging ligands 192–194
 - Morita–Baylis–Hillman reaction 277–278
 - MSs. *See* Mesoporous silicas (MSs)
- n**
- nano-metal particles (NMPs)
 - heterogeneous solid-supported catalysts 22, 23
 - and ILs 23
 - IL-supported/mediated 22, 23, 28–31
 - preparation 23–28
 - nanoreactors, catalysis
 - catalytic nanoreactor engineering 390
 - cooperative activation effect. *See* Cooperative activation effect
 - heterogeneous catalysis 351
 - homogeneous catalysis 351
 - hydrothermal/solvothermal synthetic approach 353
 - microenvironment engineering 385–388
 - MOFs 354–355
 - molecular catalysts. *See* Molecular catalysts, nanoreceptors
 - MSs 353–354
 - pore confinement effect 369–377
 - porous structure, influence 388–389
 - 2-naphthalenethiol (2-NT) probe molecule 330–331
 - nitroalkanes
 - aldol reaction of nitroacetates 456, 457
 - conjugate addition of 2-nitropropane 455, 456
 - PTC conditions 455–457
 - NMPs. *See* Nano-metal particles (NMPs)
 - NMPs preparation, ILs
 - Au NPs 27
 - IL as reducing agent 24
 - molecular hydrogen, reducing agent 24–26
 - NaBH_4 , reducing agent 26
 - rod-like gold NPs 27, 28
 - silver nanostructures 27, 28
 - noncovalent bonding methods
 - chiral Co(Salen) complex 365
 - chiral ligand/catalyst 364
 - immobilization methods 364
 - ordered MSs 365
 - porous matrix 364–365
 - zeolites, ship-in-a-bottle synthesis 365–366
 - nonionic surfactants
 - AOT reverse micelles and LiP 513
 - catalytic activity 515, 516
 - GGDE 514
 - Noyori–Ikariya type catalysts 216
 - Noyori-type Ru(II)-catalyzed asymmetric hydrogenation
 - aromatic ketones 186, 187
 - generation 185
 - hetero-coordination, achiral bridged diphosphine 186
 - 2-NT. *See* 2-Naphthalenethiol (2-NT) probe molecule
- o**
- 1-octene 205
 - OMSs. *See* Ordered mesoporous silicas (OMSs)
 - open catalytic active sites
 - Amplex Red to resorufin, conversion 340
 - atomic layer deposition Al_2O_3 341–342
 - dependence of reaction rate 341
 - HAADF-STEM 342–343
 - kinetic poisoning experiment 341
 - pseudo-first-order rate constant 341
 - reactivity of edge and corner sites 343
 - resazurin reduction reaction 340–341
 - single nanoparticle catalysis 340
 - turnover frequency (TOF) 341–342
 - ordered mesoporous silicas (OMSs)
 - MCM-41 phase 85
 - M41S synthesis 85
 - MSU-1 85
 - Na ion exchange reaction 86
 - organic–inorganic hybrid materials, types 87–88
 - PMO 87
 - pore sizes and surface areas 85
 - SBA-1 86
 - SBA-15 85–86
 - surfactant interactions, types 86–87

- organic base-catalyzed reactions, SiO₂ enhanced
 - amine-catalyzed aldol reaction 7
 - amine-functionalized silicas 8
 - cyclic carbonate synthesis 9, 10
 - heterogeneous and homogeneous acid and base catalysts 8
 - Knoevenagel reaction 8, 9
 - MCM-41 catalysts, amine-functionalized 9
 - nitroaldol reactions 9, 10
 - propylene oxide 10, 11
 - silica-supported phosphonium salts 10, 11
 - organic substances, hydrogenation
 - citral 473–474
 - cyclohexene to cyclohexane 472
 - imines 475–476
 - levulinic acid 474
 - *N*-(1-phenylethylidene)aniline 475–476
 - phenol 474–475, 476
 - α -pinene to pinane 472–473
 - prochiral α -enamides 475
 - organic synthesis 419–420
 - organic–inorganic hybrid materials
 - advantages 91
 - aldol condensations 99, 103
 - amine groups 91–93
 - catalysis 100–102
 - co-condensation method (one-pot synthesis) 88–89, 91–96
 - esterification of alcohol 103–104
 - grafting method 88, 91–96
 - HPMO 89
 - ionic liquids (ILs) 93–95
 - Knoevenagel condensation 99, 103
 - PMOs 88, 89
 - organocatalysts
 - aldol reaction. *See* Aldol reaction
 - Michael addition 309–312
 - organometallics
 - heterogeneous asymmetric catalysis. *See* Heterogeneous asymmetric catalysis, self-supporting approach
 - homogeneous asymmetric catalysis. *See* Homogeneous asymmetric catalysis
 - oxidations, SCFs
 - alcohols in scCO₂ 480
 - cyclohexane 481
 - propylene oxide (PO) 480–481
 - styrene oxide 480
 - oxidative desulfurization (ODS), H₂O₂ as oxidant
 - alcohols and sulfides, selective oxidation 293
 - Brønsted acid ion liquids (BAILs) 294–295
 - BT oxidation, in water/oil emulsion droplets 290
 - catalytic oxidative system advantages 293–294
 - diesel, sulfur removal 291
 - emulsion catalysis 287
 - emulsion system, fluorescence optical micrograph 292
 - “green” solvent 287
 - hybrids 295
 - hydrodesulfurization (HDS) 287
 - IL emulsion system, preparation 294
 - lanthanide-containing POMs 293–294
 - metastable emulsion droplets 289, 292
 - microemulsion droplets 289
 - phosphotungstic compounds 290
 - polyoxometalate (POM) 288
 - quaternary ammonium cations 288
 - straight-run diesel 291, 292
 - sulfones 291
 - sulfur-containing compounds 288–289
 - sulfur-specific GC-FPD chromatograms 291
 - transmission electron microscopy (TEM) 295, 296
 - turnover number (TON) 289
 - oxindoles
 - asymmetric PTC 454–455
 - synthesis of (-)-esermethole 454
 - synthesis of tricyclic diamines 454–455
 - oxygen activation 409–410
- p**
- Pd NPs, IL-supported
 - acetophenone hydrogenation reaction 34
 - hydrogenation, alkynes 32, 34
 - *in situ* formation and stabilization 34
 - isomerization reaction 34, 35
 - phenanthroline (Phen) ligand-protected Pd NPs 32
 - supercritical CO₂ extraction 35
 - perfluoroalkylated PCP pincer ligand 266
 - periodic mesoporous organosilica (PMO)
 - aliphatic 97–98
 - aromatic 98
 - catalytic application 104
 - ethane-bridged 96
 - HPMO 98–99
 - mild pH conditions 97
 - OMSs 87
 - organic–inorganic hybrid materials 88, 89
 - synthesis with surfactants 96–97
 - phase-transfer catalysis (PTC)
 - advantages 425

- phase-transfer catalysis (PTC) (*contd.*)
 - asymmetric PTC. *See* Asymmetric PTC, organic synthesis
 - phosphine ligands 490, 491
 - phosphorus ligands 400, 401, 402
 - photocatalysis
 - biphenyl (Bp)-PMO catalyst 107
 - impregnation method 106
 - nano-structured metallic catalyst 107
 - nano-structured titania (TiO₂) 106
 - organic–inorganic mesoporous materials 107
 - porphyrin-bridged PMO 107
 - Pichler–Schulz mechanism 555
 - PMO. *See* Periodic mesoporous organosilica (PMO)
 - poly(ethylene glycol) (PEG)
 - definition 114, 116
 - PEG-bounded TsDPEN 117
 - and PEG-supported chiral ligands and catalysts 116
 - PEG-supported chiral monophosphite 117, 118
 - PEG-supported Ru-BINAP catalyst 118
 - PEG-supported tartaric acid ester ligands 116, 117
 - uses, chiral catalysts immobilization 114
 - poly(propyleneimine) (PPI) 133–135
 - polyacrylamide gel electrophoresis (PAGE) 402, 403
 - polymer-bounded catalysts
 - aldehydes, transfer hydrogenation 318
 - amphiphilic polymer-immobilized nanoparticle 314
 - amphiphilic polystyrol/polyethylene glycol resins 316
 - C–C coupling reactions 314
 - cinnamaldehyde 314
 - emulsion system, microscope images 319
 - ethyl pyruvate 313, 314
 - Heck reactions 314–315
 - heterogeneous catalysts 313
 - hydrogenated cinnamaldehyde 313
 - Kharasch reaction 316–317
 - *N*-heterocyclic carbene (NHC) palladium complexes 316–317
 - 1-octene, hydroformylation 318
 - O/W emulsion catalytic system 317–318
 - palladium–phosphine complex catalysts 315
 - poly(amido-amine) (PAMAM) dendrimer 315
 - polymer–metal hybrid materials 312–313
 - polymer-stabilized metal emulsion catalysis 313–314
 - polyvinylpyrrolidone (PVP) 313
 - selective catalytic reaction 313
 - Suzuki coupling 315
 - polymerization vs. chain growth model, FT reaction
 - lumped microkinetics 576, 578
 - mechanistic model, C–C bond formation and CO activation 581
 - rate of CO consumption 579, 582
 - simulated coverages, monomer formation 580
 - pore confinement effect
 - achiral cluster Os₃(CO)₁₂ 374
 - asymmetric epoxidation 372–373, 376
 - asymmetric reactions 370–371
 - chiral bis(oxazoline) ligand 373
 - chiral Mn(Salen) catalysts 369–370, 372
 - *cis*- and *trans*-methylstyrene, oxo-Mn(Salen) 377
 - conversion and enantioselectivity 375, 377
 - MCM-41 and MCM-48 372
 - MCM-41 and SBA-15 371
 - olefins 375
 - pores influencing chiral catalysis 375, 376
 - Pt nanoparticles in CNTs 374
 - Zn-Al layered double hydroxides 372, 373
 - pore entrance size, molecular catalysts
 - chiral catalysts 367
 - encapsulation strategy, advantages 369
 - FDU-12-BINAPO 367, 368
 - MSs 366
 - nanocages 366
 - porous structure, nanoreactors
 - Binol-based MOFs 388–389
 - carbonyl–ene reaction 388
 - chiral PMOs 389
 - Friedel–Crafts reaction 388
 - PPI. *See* Poly(propyleneimine) (PPI)
 - propene 202
 - Pt and Ir NPs, IL-supported
 - acetone hydrogenation 36
 - enantioselective hydrogenation 36
 - size distribution, Ir(0) NPs 36, 38
 - PTC. *See* Phase-transfer catalysis (PTC)
 - PVP (poly(*N*-vinyl-2-pyrrolidone)) 402
- q**
- quat salts
 - definition 425
 - mechanism of PTC 425, 426

r

- RCH/RP process. *See* Ruhrchemie/Rhone-Poulenc (RCH/RP) process
- reverse micelles
- additives. *See* Additives, reverse micelles
 - application. *See* Application, reverse micelles
 - biodiesel preparation 544
 - conformation 528–530
 - enzyme catalysis. *See* Enzyme catalysis
 - IL-based microemulsions 544
 - ionic liquid-based. *See* Ionic liquid-based reverse micelles
 - micellar enzymology 543
 - microemulsifications 512
 - microemulsions 511–512
 - molecular organic solvent-based. *See* Molecular organic solvent-based reverse micelles
 - “polar pool” 512
 - pseudophase model and enzyme-catalyzed reaction kinetics 530–531
 - room-temperature ILs 512
 - “water pool” 512
 - W/IL microemulsions 544
- Rh NPs, IL-supported
- (*E*)-2-(benzoylamino)-2-propenoic acid derivatives hydrogenation 42
 - bipyridine coordination modes, NMPs 41
 - bipyridine-functionalized ILs 41, 42
 - BPys behavior, styrene hydrogenation 40–41
 - styrene hydrogenation 40
 - substituted aromatic compounds hydrogenation 40
- rhodium-phosphine complex catalytic systems
- activity and selectivity 495
 - aqueous-organic biphasic transition metal catalysis 491
 - 1-dodecene 500
 - P-C bond 495
 - phosphine ligands 490, 491
 - RCH/RP advantages 490–491
 - TPPTS 490, 494
- room-temperature ionic liquids (RTILs)
- common cations and anions 21
 - Friedel–Crafts reactions 21
 - IL-supported/mediated metal nanoparticles. *See* IL-supported/mediated metal nanoparticles
 - solid-supported IL. *See* Heterogeneous catalysis with homogeneous performance
- RTILs. *See* Room-temperature ionic liquids (RTILs)

Ru complex-catalyzed enantioselective reactions

- achiral bisphosphine/Ru/chiral diamine-catalyzed asymmetric hydrogenation 171–172
 - achiral monophosphine/Ru/chiral diamine-catalyzed asymmetric hydrogenation 168–171
- Ru NPs, IL-supported
- catalytic hydrogenation 1,3-cyclohexadiene and cyclohexene 37
 - phosphonium and imidazolium ILs 39, 40
 - Rh NPs. *See* Rh NPs, IL-supported
 - [Ru(COD)(2-methylallyl)₂] reduction 37
 - size distributions 37, 39
 - SYBYL representations, CYD 37, 39
 - toluene hydrogenation 37
- Ruhrchemie/Rhone-Poulenc (RCH/RP) process
- large-scale manufacturing method 497
 - *m*-OHCC₆H₄SO₃Na and C₃H₇P(*m*-C₆H₄SO₃Na)₂ 494
 - TPPTS 490
 - traditional cobalt process 497
 - water-soluble rhodium catalyst 489–490
- Ruhrchemie/Rhône-Poulenc process 202

s

- scanning tunneling microscopy (STM) 338
- SCFs. *See* Supercritical fluids (SCFs), catalysis
- Schiff base/Ti(IV)-catalyzed asymmetric hetero-Diels–Alder reaction
- acid additives 153, 154
 - tridentate Schiff base ligands 153
- selective hydrogenation
- Au₂₅(SR)₁₈ clusters 407–409
 - H₂-D₂ exchange reaction 407
 - turnover frequencies (TOFs) 407
- selective oxidation
- alcohols. *See* Alcohols, selective oxidation
 - alcohols aerobic oxidation 61–62
 - alkanes 410–411
 - alkenes. *See* Alkenes, selective oxidation
 - CO oxidation. *See* CO oxidation
 - olefins aerobic oxidation 62–63
 - oxygen activation 409–410
- silica gel-confined IL catalysts
- confinement effect, supercage of silica gel 55
 - covalently supported IL catalysts 56–57
 - emission behavior 56
 - preparation and model 55

- single reaction center site model, FT reaction
 - BEP relationships 586, 587
 - catalysts performance 590, 591
 - influence, CO activation energy 588, 590
 - M–O and M–C bond energies 586
 - predicted RCO and α 588, 589
- size-exclusion chromatography (SEC) 402, 403
- solid-supported IL catalyzed reactions. *See* Heterogeneous catalysis with homogeneous performance
- soluble polymeric supports 112–113
- Sonogashira reaction
 - carbpalladacycle complex 48
 - description 48
 - iodobenzene and phenylacetylene 48
- SSITKA. *See* Steady-state isotopic transient kinetic analysis (SSITKA)
- steady-state isotopic transient kinetic analysis (SSITKA) 594
- Stille reaction 47
- sulfonated Xantphos 204
- sulfonic/carboxylic acid-amine bifunctional catalyst
 - cooperative catalysis, sulfonic acid and primary amine groups 4
 - heterogeneous and homogeneous acid and base catalysts 3, 4
 - initial turnover frequency, aldol reactions 6, 7
 - nitroaldol reactions 5
 - SBA-15-AB 3, 4
 - SBA-G-AP0.5 6
 - surface silanol groups 6
 - surface structures, amine and carboxylic acid grafted silica catalysts 6
- sulfur ligands
 - ligand exchange 401–402
 - reduction of Au(I)SR 401, 402
 - thiolation 402–403
- supercritical fluids (SCFs), catalysis
 - alkylation 481–482
 - clustering 471
 - CO₂ cycloaddition to epoxide 482–483
 - CO₂, hydrogenation. *See* CO₂, hydrogenation
 - cycloaddition in scCO₂ 470
 - description 469–470
 - ethylene carbonate 470
 - features of reactions 471–472
 - hydroformylation reactions 479–480
 - organic substances, hydrogenation. *See* Organic substances, hydrogenation
 - oxidations 480–481
 - pure substance 470
 - thermophysical properties 470, 471
- supported IL catalyst
 - alkylation and esterification reactions 73–74
 - asymmetric catalysis 74–77
 - BMImPF₆ 52, 53
 - [BMIm][Tf₂N] behavior 54
 - catalytic carbonylation reaction. *See* Catalytic carbonylation reaction
 - catalytic hydrogenation. *See* Catalytic hydrogenation
 - enzyme catalysis 77–79
 - heterogenized homogeneous IL catalyst system 51
 - immobilizing organometallic complexes 53, 54
 - isomerization and oligomerization 72–73
 - Lewis acidic IL-phase catalysis 51
 - magic-angle spinning (MAS) 53
 - para-hydrogen induced polarization (PHIP) NMR spectroscopy 53
 - selective oxidation 61–63
 - two-dimensional nano-IL catalyst 51, 52
 - water-gas shift reaction 70–72
- Suzuki reaction
 - description 42–43
 - ionic polymer-stabilized Pd NPs/IL system 44
 - ligands, norborn-5-ene-2, 3-dicarboxylic anhydride 44
 - nitrile-functionalized IL 43
 - Pd NPs 43, 44
- Suzuki–Miyaura coupling
 - aryl bromides 227, 230
 - aryl iodides and bromides 233
 - biaryls synthesis 226–227
 - bulky phosphine ligands 228
 - catalyst recycle 229, 230
 - drawback 233
 - heterocyclic halides with boronic acids 228, 230, 231
 - ligand-free catalysts 231–232
 - microwave promoted 234
 - Pd(II)-(di-2-pyridyl)methylamine 229–230
 - pH-responsive palladium catalyst 232–233
 - silica-supported catalyst 232
 - water-soluble phosphine ligands 227
- t**
- tartaric acid-derived quaternary ammonium salts 438, 439
- thermoregulated phase-transfer catalysis (TRPTC) 206, 505–506

- Ti(IV)-catalyzed enantioselective reactions
- BINOLate/Ti-catalyzed asymmetric carbonyl–ene reaction 156–158
 - BINOLate/Ti(IV)-catalyzed asymmetric hetero-Diels–Alder reaction 154–156
 - BINOLate/Ti-catalyzed asymmetric ring-opening aminolysis, epoxides 158–159
 - Schiff Base/Ti(IV)-catalyzed asymmetric hetero-Diels–Alder reaction 153–154
- Ti-catalyzed hetero-Diels–Alder reactions 132, 133
- transient kinetics, FT reaction
- analysis, isotope switch experiments 597
 - reported data vs. interpretation, simulations 598
 - residence times, methane 595, 597
 - simulations 595, 596
 - SSITKA measurements 594
- transition-metal-catalyzed asymmetric reactions 133, 134
- TRPTC. *See* Thermoregulated phase-transfer catalysis (TRPTC)
- u**
- UCC. *See* Union Carbide Corporation (UCC)
- Ullmann reaction
- “green” pathway 49, 50
 - ionic liquids 48, 49
- union Carbide Corporation (UCC) 489, 490
- urea-amine bifunctional catalyst
- acetone with 4-nitrobenzaldehyde aldol reaction 2, 3
 - aldol reaction, urea and amine groups 2, 3
 - surface structures, immobilized MSNs 2
- v**
- VOCs. *See* Volatile organic compounds (VOCs)
- volatile organic compounds (VOCs) 469
- w**
- water catalysis
- heat capacity 201
 - heat of vaporization 201
 - hydroformylation. *See* Hydroformylation
 - hydrogenation. *See* Hydrogenation
 - properties 201–202
- water oxidation reactions
- molecular catalysts 380
 - SBA-16, nanocage 380
 - TOF 380
- water-gas shift (WGS) reaction 345–346
- water-in-oil (W/O) microemulsions
- cetyltrimethylammonium-based surfactants 521
 - counterion binding 522
 - interfacial region 515, 518
 - nonionic surfactant 514
 - *p*-nitrophenyl-*n*-hexanoate 517, 520
 - reverse micelles 512
 - straight-chain alcohols 513
- water-soluble catalyst, rhodium-phosphine complex systems 490–491
- WGS. *See* Water-gas shift (WGS) reaction
- W/O microemulsions. *See* Water-in-oil (W/O) microemulsions
- x**
- X-ray absorption near edge structure (XANES) measurements 344
- X-ray photoelectron spectroscopy (XPS) 28–29, 328
- z**
- Zn complex-catalyzed enantioselective reactions
- BINOLate/Zn/diimine-catalyzed asymmetric diethylzinc addition 162–164
 - BINOLate/Zn/diimine-catalyzed asymmetric hetero-Diels–Alder reaction 165–167
 - chiral amino alcohol/Zn/racemic amino alcohol-catalyzed asymmetric diethylzinc addition 159–162

