

Index

a

absorption 20–22
 coefficient 64
 spectra of exciplexes 338
 spectrum 102
 strength analysis 21

acridine orange 277

ACRXTN and ACRSA 282–285

adiabatic electronic energies 299

Ag₂Cl₂(dppb)₂ 34

Ag(dbp)(P₂-nCB) 38, 44
 characterization of 40–45
 emission quenching 36–38

Ag(I)-based TADF compounds
 Ag₂Cl₂(dppb)₂ 34
 Ag(phen)(P₂-nCB) 34–38
 characterization of Ag(dbp)(P₂-nCB)
 40–45
 emission quenching in
 Ag(phen)(P₂-nCB) 36–38
 energetically lower-lying 4d-orbitals
 34
 ligand-centered (LC) orbitals 34
nido-carborane-bis-
 (diphenylphosphine) (P₂-nCB)
 34
 sterical hinderance 38–40

Ag(I) complexes 1–46, 84–85

Ag(phen)(P₂-nCB) 34–38, 40

aggregation-induced emission (AIE)
 403, 472, 563

angular dependence_TADF 215

anionic bidentate ligand
 diphenyl-bis(pyrazol-1-yl)borate
 (pz₂Bph₂⁻) 142

anthracene crystal 199

atomic force microscope (AFM),
 ITO/PEDOT 484

Au(I) complexes 84, 150, 159

Avogadro number 21

b

B3LYP/def2-SVP level of theory 15,
 20, 21, 23, 24

benzothiazole 403, 404

benzoxazole 403, 404

bidentate binap 84

bilayer-type exciplex 416

2,2'-bipyridyl based ligands 181–182

1,3-bis(carbazol-9-yl)benzene (mCP)
 143, 170, 517

1,3-bis{3-[3-(9-carbazolyl)phenyl]-9-
 carbazolyl}benzene (CPCB)
 514

2,7-bis(9,9-dimethyl-acridin-10-yl)-9,9-
 dimethylthioxanthene-S,S-
 dioxide (DDMA-TXO2) 449,
 457–460

bisdiimine-type copper(I) complexes
 ([Cu(NN)₂]⁺) 131

bis(diisobutylphenylphosphino)amido
 (PNP) ligands 157

bis(diphenyl-phosphino)methane
 (dppm) 181

bis(diphenylphosphino)ethane (dppe)
 181

bis[2-(diphenylphosphino)phenyl]ether
 (POP) 131, 168, 181, 214, 449

6,6'-bis(2-4diphenylquinoline) (B1PPQ)
 338

- 2,7-bis(phenoxazin-10-yl)-9,9-dimethylthioxanthene-*S,S*-dioxide (DPO-TXO2) 431, 432, 436, 441–443
- bis-4,6-(3,5-di-3-pyridylphenyl)-2-methylpyrimidine (B3PYMPM) 332, 365, 414
- 1,3-bis(9H-carbazol-9-yl) benzene (mCP) 474, 475, 478, 484, 485, 487, 495
- 2-(bis(trimethylsilyl)methyl)-6-fluoropyridine (3 L) 100
- 2-(bis(trimethylsilyl)methyl)-6-methylpyridine (2 L) 100
- 2-(bis(trimethylsilyl)methyl)pyridine (1 L) 100
- 2-(bis(trimethylsilyl)methyl)quinoline (4 L) 100
- blended single-layer exciplex 414–416
- blue fluorescent emitter 486–490
- blue OLEDs
academia 552–554
efficient, stable 550–552
- blue PHOLED emitters 257
- Boltzmann constant 2
- Boltzmann distribution 4, 109, 121, 235, 263
- Boltzmann statistics 274, 301, 318
- Boltzmann thermal distribution 4
- Born–Oppenheimer approximation 299, 304, 305
- buffer hole-injection layer (Buf-HIL) 515
- tert*-butylcarbazole donor 526
- C**
- canonical partition function 259
- carbazolyl-modified 1-(2-pyridyl)-pyrazole based ligands 191–192
- cationic Cu(I)-bis-phen complexes 275, 276
- cationic Cu(I) complexes 62
- cationic [Cu(NN)₂]⁺ and [Cu(NN)(PP)]⁺ complexes 178
- charge-neutralized Cu(I) complex 140
- charge-neutral three-coordinate Cu(I) complexes 146–155
- charge-separated state 347
- charge separation versus charge recombination 343–345
- charge transfer (CT)
emitter 528
exciplex 332
- chelating diphosphine POP ligand 180
- chemical structures, of host 517
- closed quantum system 312
- combined steady-state and decay data analysis 250–252
- Commission Internationale de l'Éclairage (CIE) 504, 551
- comprehensive efficiency analysis of OLEDs 206–209
- computation of ISC and RISC rate constants 264–265
- Condon approximation 260
- electronic spin-orbit coupling matrix elements 261–262
- overlap of vibrational wavefunctions 262–263
- confined exciplex systems 356
- conformational rigidity 270
- contact radical ion pair (CRIP) 342–343
- copper (I) complexes
four-coordinated
Cu(I)-bis-phenanthroline complexes 275–277
three-coordinated
Cu(I)-NHC-phenanthroline complex 270–275
- Coulomb energy 434, 435
- crystalline Cu(I) complexes 128
- [Cu(CH₃CN)₄(POP)]BF₄ 141
- Cu₂Cl₂(N[^]P)₂ 27
- [Cu(dbp)P₂]₂BF₄ 133
- [Cu(dbp)(POP)]BF₄ 133
- [Cu(dmp)₂]⁺ (dmp = 2,9-dimethyl-1,10-phenanthroline) complex 123
- [Cu(dmp)(dtbp)]⁺ (dtbp = 2,9-di-*tert*-butyl-1,10-phenanthroline) 179

- Cu(dmp)(phanephos)⁺ 15–17, 20, 23, 28–30
 [Cu(dmp)(phanephos)](PF₆) 18, 19, 134
 [Cu(dmp)(POP)]BF₄ 131
 [Cu(dppbz)₂]BF₄ (**48**) and [Cu(dppbz)(POP)]BF₄ 145
 [Cu(dppbz)₂]⁺ (dppbz = 1,2-bis(diphenylphosphino)benzene complex 123
 Cu(dppb)(pz₂Bph₂) 9, 22–25, 29, 30
 Cu₂(4L)₂ 111
 Cu(I) complexes
 with large ΔE(S₁–T₁)
 absorption 20–22
 DFT and TD-DFT calculations 16
 flattening distortions and non-radiative decay 16–18
 radiative S₁→S₀ rate 20–22
 Strickler-Berg relation 20–22
 TADF properties 18–20
 molecular and electronic structure on emissive properties 178–181
 for OLEDs
 energy levels of molecular orbitals in tetrahedral geometries 122, 123
 ligand variation 123–125
 with small ΔE(S₁–T₁)
 DFT and TD-DFT calculations 22–23
 emission spectra and quantum yields 23
 temperature dependence of the emission decay time and TADF 28–30
 triplet state T₁ and spin-orbit coupling 23
 Cu(I)-NHC-phenanthroline complex 270–275
 Cu(I)-phenanthroline complex, ultrafast dynamics of a 313–316
 Cu(L_{iPr})(SPh) 152
 Cu(L_{Me})(SPh) 152
 [Cu(μ-CAN)]₂ (CAN = 2-(bis(trimethylsilyl)methyl)pyridine derivatives)
 absorption spectrum 102
 DFT calculation 103, 104, 112
 emission decay kinetic analysis 105, 106, 108, 109
 emission properties 104, 105
 measurement 111
 OLED device 110, 112
 outline 100
 planer 8-membered cyclic ring 112
 synthesis 111
 X-ray crystallographic study 101, 102
 X-ray structure analysis 112
 Cu₄(μ-I)₄ cubane unit 81
 Cu₂(μ-X)₂ cores 155, 156
 [Cu(NN)(PP)]⁺ complexes 132, 133, 142
 with NN ligands 134–142
 with phen or bipy derivatives as ligands 130–134
 [Cu(N[^]N)(POP)] complexes 78
 [Cu(N[^]N)(P[^]P)] 61, 77–79
 [Cu(N[^]N)(P[^]X)] 77–79
 [Cu(phen)(POP)](BF₄) 189
 [Cu(P[^]P)₂] 61, 78
 cuprous emissive complexes 177
 Cu(pympz)(POP)]⁺ 78
 [Cu(pytfmpz)(POP)]⁺ 78
 Cu₂(3L)₂ 111
 [Cu(tmbpy)(POP)]BF₄ 133
 [CuX(PPh₃)₂(4-Mepy)] (X = Cl, Br, I) complexes 79
 cyano-based TADF molecules 391–396
 cyclometalated iridium(III) complexes 93, 120, 122
- d**
- DCz-DBTO₂, temperature dependent emission of 434
 decay data analysis 249–252
 decay time τ 6
 delayed fluorescence (DF) 199, 203–204, 300–301
 kinetics /of the ¹CT prompt state 449
 intensity 249

- delayed fluorescence (DF) (*contd.*)
 lifetime 233
 TADF 467
 time resolved emission in solid state
 440, 446
 delayed $S_1 \rightarrow S_0$ fluorescence 5
 $\Delta E(S_1-T_1)$ minimization 7
 density functional theory (DFT) 16,
 64, 267, 469, 555
 density matrix formalism of MCTDH
 311–312
 Dexter energy transfer 383, 487,
 495
 DFT and TD-DFT calculations 16,
 22–23
 DFT-based multireference
 configuration interaction 267
 diazafluorene 404–405
 1,4-diazatriphenylenes (ATP) 402, 403
 dibenzothiophene-*S,S*-dioxide acceptors
 531
 9,10-dicyanoanthracene (DCA) 336
 1,4-dicyanobenzene (DCB) 343
 diimine ligands ($[Cu(NN)_2]^+$) 177
 dilute methyl cyclohexane (MCH) 431
 dimeric Cu(I) chlorido NHC-picolyl
 complexes 82
 dimethylacridine (DMAC) 552, 564
 2,9-di-*n*-butyl phenanthroline ligand
 38
 dinuclear Cu(I) complex
 $[Cu_2(NN6)_2P_4](BF_4)_2$ 137
 dinuclear Cu(I) complexes 80–83
 $[(dppbz)CuX]_2$ 146
 for OLEDs
 other dinuclear Cu(I) complexes
 157
 possessing $Cu_2(\mu-X)_2$ cores
 155–157
 dinuclear Cu(I) halide complexes 83
 dinuclear TADF-type Cu(I) emitters
 122
 diphenyl ketone 408–410
 2-Di-phenylphosphinopyridine
 (PyrPHOS) 83
 diphenyl sulfoxide 405–407
 diphosphine ligands 147
 dipole moment operator 64
 dipyrido[3,2-*a*:2',3'-*c*]phenazine (dppz)
 137
 donor-acceptor (D-A) molecular
 systems, TADF
 cyano group 391–396
 design principles of 384–386
 intramolecular or intermolecular
 architectures 391
 molecular designs and properties of
 380
 nitrogen heterocycle 396
 down-hill process 259, 260
 DPEPO (bis[2-(diphenylphosphino)
 phenyl]ether) 214
 $[(dppbz)Cu(\mu-X)]_2$ (X) 155
 DPTZ-DBTO2
 emission spectra of 437
 phosphorescence spectra of 438
 steady-state fluorescence spectra of
 439
 Direct Singlet Harvesting 46
 Duschinsky rotation effect (DRE) 308
 DV-CDBP 511, 512, 518
 DV-MOC-DPS 511
- e**
- electrical excitation 240
 conditions for efficient
 electroluminescence 241–244
 steady-state 240–241
 electric dipole radiation 201
 electrochemical cyclo-voltammetry
 measurements 437
 electroluminescence (EL) 476, 482,
 501
 devices 490
 efficiency 242
 emitters 257–258
 4P-NPB 489
 properties, of emitters 507
 quantum efficiency 202–203
 electroluminescence (or external)
 quantum efficiency (EQE)
 202–203, 282, 382, 390
 electron-accepting moieties 365–368,
 391, 396
 electron-donating 368
 moieties 360

- electron donor molecules 360
 electronic excitation 2, 267, 271, 277, 286, 331, 347, 471
 electronic spin-orbit coupling (SOC) 259
 matrix elements 261–262
 electronic spin-spin coupling (SSC) 259
 electron paramagnetic resonance (EPR) spectroscopy 302
 electron-rich pyridylpyrazole ligands NN20–NN22 141
 electron transfer (ET) 333, 437, 471
 electron transport layer (ETL) 202, 392, 393, 458, 481
 El-Sayed rules 261, 262, 268, 345
 El-Sayed's law 302
 emission decay kinetic analysis 105, 106, 108, 109
 emission decay time 5–7, 28–30, 41, 45, 46, 151, 158, 177, 182, 213, 274
 emission layer (EML) 1, 2, 4, 201, 202, 481, 484, 486, 561
 emission properties 22, 23, 29, 34, 40, 52, 81, 99, 104–105, 132, 133, 143, 148, 273, 277, 428–438
 emission quenching in Ag(phen)(P2-nCB) 36–38
 emission spectra and quantum yields 23
 emissive charge-neutral three-coordinate Cu(I) complexes 154
 emissive Cu(I) complexes 123, 160, 177, 178, 180–182, 184
 emissive [Cu(NN)(PP)]⁺ complexes 194
emitter host interactions 258, 457–459
 emitters 2, 36, 61–86, 119–170, 353
 energetically lower-lying 4d-orbitals 34
 energy diagram for TADF 459
 energy gap law 76, 131, 140, 262, 280, 436, 524
 energy level diagram 426
 energy separation $\Delta E(S_1-T_1)$ 30–34
 eosin dyes 6, 19, 28, 29, 44, 321, 426, 485, 493
 E-type delayed fluorescence 62, 121, 199, 258, 426
 evaporation chamber 548
 excimer 104, 232, 331, 386
 exciplex 332
 bilayer-type 416
 blended single-layer 414
 decay processes of
 charge-separation vs charge recombination 343–345
 CRIP and SSRIP 342, 343
 fluorescence rate 340–341
 ISC 345–346
 electron acceptor molecule 360
 electron donor molecule 360
 electron-accepting moieties 365–368
 electron-donating moieties 360–365
 electronic structures of
 charge-transfer (CT) 332
 formation 332–335
 HOMO and LUMO 332
 local-excitation (LE) 331
 molecular orbitals 332
 stabilization 335, 337
 OLEDs. *See* organic light-emitting diodes (OLEDs)
 optical properties of
 absorption spectra 338–339
 photoluminescence 336–338
 organic solid films
 prompt vs. delayed fluorescence 347–350
 spectral shift 350–352
 triplet LE state 346
 excitation energies and radiative rate constants
 DFT-based multireference configuration interaction 267
 fluorescence and phosphorescence rates 268–269
 time-dependent density functional theory 266

- excited states 450
 cycles 235–238
 properties 452–455
- f**
- fac*-Ir(ppy)₃ derivatives 93
 FASAE, chemical structure of 433
 FDQPXZ 524
 Fermi's Golden rule (FGR) approach
 259, 278, 302, 303
 first order Fermi's golden rule 320
 first order mixing coefficient 297
 first-order perturbation theory 260
 flattening distortion dynamics of the
 MLCT excited state 76–77
 flattening distortions and non-radiative
 decay 16–18
 fluorescence (FLUO) 544
 blue 199
 efficiency 239
 materials, TADF 563–564
 OLED 119
 and phosphorescence decays 232
 and phosphorescence rates 268–269
 quantum yield 234
 rate constant 340
 formed 335
 Förster resonant energy transfer (FRET)
 258
 Förster energy transfer 383
 Förster ENT 358
 four-coordinated
 Cu(I)-bis-phenanthroline
 complexes 270, 275–277
 Fourier transform photocurrent
 spectroscopy (FTPS) 338
 Franck–Condon factors 2, 16, 36, 76,
 414
 Frank–Condon weighted density of
 states 446
 full wave at half maximum (FWHM)
 482
 fullerenes 245, 379, 380
- g**
- gated iCCD detection system 440
- generic Jablowski diagram for TADF
 460
 green and blue emitters 77–79
 grid based quantum nuclear wavepacket
 dynamics 305
 ground and excited-state wavefunctions
 298
 gyromagnetic factor 261
- h**
- halogen-bridged binuclear copper
 complex [(dppbz)Cu(μ-X)]₂
 147
 halogen-bridged Cu(I) complexes 81
 halogen-bridged dinuclear Cu(I)
 complexes 61, 95, 156
 heavy atom effect 159, 313, 425
 heptazine 401
 heteroatom lone pair orbitals 427
 heteroleptic Cu(I) complexes 77, 83,
 133, 134, 180, 182–188, 191–194
 heteroleptic [Cu(N[^]N)(P[^]P)] complexes
 61, 78
 heteroleptic [Cu(NN)(PP)]⁺ complexes
 177, 178, 181
 heteroleptic diimine/diphosphine
 [Cu(NN)(PP)]⁺ complexes
 with 2,2'-bipyridyl based ligands 181
 with carbazolyl-modified
 1-(2-pyridyl)-pyrazole based
 ligands 191–192
 with 1-phenyl-3-(2-pyridyl)pyrazole
 based ligands 192–193
 with 3-phenyl-5-(2-pyridyl)-
 1H-1,2,4-triazole based ligands
 193–194
 with 2-(2'-pyridyl)benzimidazole and
 2-(2'-pyridyl)imidazole based
 ligands 182–185
 with 1-(2-pyridyl)-pyrazole based
 ligands 189–190
 with 2-(2-pyridyl)-pyrrolide based
 ligands 188
 with 5-(2-pyridyl)tetrazole based
 ligands 185–187
 with 3-(2'-pyridyl)-1,2,4-triazole
 based ligands 187–188

- heteroleptic PyrPHOS dinuclear complexes 83
- highest occupied molecular orbital (HOMO) 3, 63, 123, 178, 200, 270, 298, 332, 382, 469, 470, 491, 545
- highly efficient thermally activated delayed fluorescence device 214–218
- highly emissive d^{10} metal complexes
 advantages 61
 Ag(I) complexes 84–85
 Au(I) complexes 84–85
 Cu(I) complexes 61
 dinuclear Cu(I) complexes 80–83
 flattening distortion dynamics of the MLCT excited state 76, 77
 green and blue emitters 77, 79
 molecular structures of Cu(I) complexes 65
 molecular structures of d^{10} metal complexes 84
 Pd(0) complexes 84–85
 phosphorescence and TADF mechanisms 62, 64
 photophysical properties 72
 Pt(0) complexes 84–85
 structure-dependent photophysical properties 64–76
 three-coordinate Cu(I) complexes 79, 80
- hole transport layer (HTL) 201, 202, 547
- HOMO-LUMO excitation 8, 40, 45
- homoleptic $[\text{Cu}(\text{NN})_2]^+$ complexes 177–179
- homoleptic $[\text{Cu}(\text{phen})_2]^+$ complex 179
- homoleptic $[\text{Cu}(\text{PP})_2]^+$ complexes 134
- homoleptic complex $[\text{Cu}(\text{POP})_2]^+$ 134
- homoleptic copper(I) bisdiimine complex $[\text{Cu}(\text{dtp})_2]\text{BF}_4 \mathbf{3}$ (dtp = 2,9-di-*tert*-butyl-1,10-phenanthroline) 131
- i**
- internal quantum efficiency (IQE) 93, 156, 199, 202, 218, 257, 258, 279, 286, 332, 353, 382, 466, 486, 487, 501
- intersystem crossing (ISC) 1, 345–346, 425, 467, 501
- rate constants
 beyond the Condon approximation 263–264
 computation of ISC and RISC rate constants 264
 Condon approximation 260–263
 static approaches 265
- intraligand charge-transfer (ILCT) 125
- intramolecular charge transfer (ICT)
 426, 469
 emission 121
 processes 277
- intramolecular D-A molecules
 bilayer-type exciplex 416
 blended single-layer exciplex 414–416
- ionisation potential 434
- Ir-based phosphorescent materials 120
- isoemissive point 442, 445
- isolated heteroleptic complexes
 $[\text{Cu}(\text{dmp})(\text{PP})]^+$ 134
- i*-Pr (propyl) substituted DPTZ-DBTO2 456
- j**
- Jahn-Teller distortion 80, 150, 313
- k**
- Kasha's rule 300
- ketones 245
- kinetics of the ^1CT prompt state 449
- l**
- $^3\text{LE}_D$ phosphorescence (PH) 441, 444
- ligand-centered (LC) orbitals 34
- ligand-to-ligand charge transfer (LL'CT) 3, 139
- light-emitting dendrimers 502, 528
- light emitting electrochemical cells (LEEC) 1, 133, 177
- light-outcoupling efficiency 203, 501

- light-outcoupling factor 202, 205, 218, 222
- linear vibronic coupling (LVC) model 307
- liquid crystal displays (LCDs) 466, 501
- lock-in technique 452
- low efficiency roll-off triplet-triplet-annihilation device 218–222, 405, 407, 411, 416, 417, 490, 495
- lowest unoccupied molecular orbital (LUMO) 3, 63, 123, 200, 270, 298, 332, 382, 469, 470, 491, 546
- low order Taylor expansion 306
- low-priced pure blue-phosphorescent OLEDs 120
- luminescence decay curves of
Ag(dbp)(P₂-nCB) 42
- luminescent Cu(I) complexes 83, 122, 134, 156, 179, 191, 270
- luminescent cyclometalated Ir(III) complexes 61, 62
- luminescent dinuclear copper complexes
[Cu(μ-CAN)]₂ (CAN = 2-(bis(trimethylsilyl)methyl)pyridine derivatives) 100
- cyclometalated iridium(III) complexes 93
- distortion 93
- fac*-Ir(ppy)₃ derivatives 93
- luminescence properties 99, 100
- 1,10-phenanthroline derivatives 93
- structure 94–99
- luminescent neutral [Cu(N[^]N)POP] complexes 78
- m**
- mCP (1,3-bis(N-carbazolyl)benzene) 214
- M06/def2-SVP level of theory 38
- measured decay time 6, 16, 19, 29, 217
- metal complexes exhibiting TADF 157–159
- metal-free TADF emitters
acridine orange 277
delayed fluorescence 278
- dibenzothiophene-S-S-dioxide
DBTO2 acceptor 278
- Duschinsky effects 278
- mechanism of the triplet-to-singlet upconversion 282–285
- phenothiazine (PTZ) donor 278
- photophysics 279
- reverse internal conversion 278
- second-order spin-orbit 260
- 1,2,3,5-tetrakis(carbazol-9-yl)-4,6-dicyanobenzene (4CzIPN) 279
- (metal + ligand L) to ligand L' charge transfer (^{1,3}MLL'CT) 34
- metal-to-ligand charge 262
- metal-to-ligand charge transfer (MLCT) 3, 17, 62, 93, 100, 103, 120, 259
- 2-methyltetrahydrofuran (2-MeTHF) 125, 450
- microscopic spin-orbit Hamiltonians 261
- m-MTDATA:PBD exciplexes 357
- molecular fluorescence 300
- monodentate halide anions 146
- mononuclear Cu(I) complexes for OLEDs
bis(diimine) type 129, 131
Boltzmann equation 127
charge-neutral three-coordinate Cu(I) complexes 146–155
[Cu(NN)(PP)]⁺ complexes
with NN ligands 134–142
with phen or bipy derivatives as ligands 131–134
- decay rate of emission 127
- electroluminescence processes 126
- electron exchange 126
- emission mechanism 127
- emission processes of phosphorescent emitters 126
- equilibrium constant 127
- free energy change 127
- intramolecular exciplex 126
- non-radiative rate constant 127
- photophysical properties 128
- radiative constant 127
- radiative rate constant 126
- rate constants 128

- simple excited state model 126
 - SOC interaction 126
 - temperature-dependent non-radiative processes 128
 - tetrahedral Cu(I) complexes with the LUMO on the PP ligand 129, 142–146
 - mononuclear Cu(I)-phenanthroline complexes 313
 - MTXSFCz 490–492
 - absorbance PL 491
 - molecular structure of 491
 - multi-configurational Ehrenfest (MCE) 310
 - multi-configurational time-dependent Hartree (MCTDH) approach 265, 310–311
- n**
- National Television System Committee (NTSC) 504, 512, 523
 - natural transition orbital (NTO) analysis 124, 144, 388
 - neutral Cu(I) complexes 62, 81, 140, 142
 - neutral mononuclear Cu(I) complexes 139
 - N*-heterocyclic carbene (NHC) Cu(I) complexes 153, 270
 - nido*-carborane-bis-(diphenylphosphine) (P_2 -nCB) 34
 - nitrogen heterocycle-based TADF molecules
 - benzothiazole and benzoxazole derivatives 403–404
 - diazfluorene derivatives 404–405
 - 1,4-diazatriphenylenes (ATP) 402–403
 - heptazine derivatives 401–402
 - oxadiazole 401
 - pyrimidine derivatives 400–401
 - quinoxaline derivatives 404
 - thiadiazole 401
 - triazine derivatives 396–400
 - triazole 401
 - nonadiabatic coupling elements 304, 309
 - non-isotropic emitter orientation 204–205, 209, 214, 216, 218, 222
 - non-polar zeonex matrix 446
 - non-radiative (NR) quenching 450
 - non-radiative exciton quenching processes 202
 - non-radiative rate constant 107, 108, 121, 126–128, 474, 476
 - nonpolar solvents 139, 343, 430, 433–435, 450, 468
 - nonradiative decay processes 122, 258, 349
 - normal fluorescence (NF) 120, 467
 - NPB, TPBi exciplexes 357
 - N*-phenylcarbazole (*N*-PhCz) 469
 - $n-\pi^*/\pi-\pi^*$ mixed state transitions 431
- o**
- optical modeling 205–206
 - optical properties of the DPTZ-DBTO2 451
 - optical transition dipole vectors (TDVs) 200, 204, 205, 208, 215–218, 220, 222
 - organic donor-acceptor systems 258, 262, 269, 286
 - organic light-emitting diode (OLED) 177, 501
 - black box 209–214
 - comprehensive efficiency analysis 206–209
 - delayed fluorescence 199, 203–204
 - device 110–112, 483
 - electroluminescence quantum efficiency 202–203
 - emitter materials 544–547
 - highly efficient thermally activated delayed fluorescence device 214–218
 - low efficiency roll-off triplet-triplet-annihilation device 218–222
 - non-isotropic emitter orientation 204–205
 - optical modeling 205–206
 - optical transition dipole vectors 200
 - TADF materials

- organic light-emitting diode (OLED) (*contd.*)
- device structures and operation
 - mechanisms 380–382
 - emitters 382
 - host materials and sensitizers 382–383
 - host-free 383
 - research of 380
 - TX 465
 - Using Exciplexes 353
 - to vacuum-deposited counterparts 526–527
 - working principle 200–202
- organic photo-voltaics (OPVs) 331, 347
- organoboron-based TADF molecules 411–412
- oxadiazole 137, 401, 402
- p**
- p-and e-type delayed fluorescence 199
 - Pd(0) complexes 84
 - perturbation theory 259, 260, 268, 269, 271, 303, 319
 - PF01 (4,4'-bis[phenyl(9,9'-dimethylfluorenyl)amino]biphenyl) for HTL (40 nm) 110
 - phase alternating line (PAL) systems 504
 - phenanthroline-based ligands 179, 180
 - 1,10-phenanthroline (phen) 34
 - phenothiazine 446
 - donors 531
 - phenothiazine donor and a dibenzothiophene-*S,S*-dioxide acceptor (PTZ-DBTO2)
 - absorption of 431
 - chemical structures of 430
 - phenothiazine donor unit (PTZ) 452
 - phenothiazine-dibenzothiophene-*S,S*-dioxide (PTZ-DBTO2) 429
 - 3-phenyl-5-(2-pyridyl)-1H-1,2,4-triazole based ligands 193–194
 - 1-phenyl-3-(2-pyridyl)pyrazole based ligands 192–193
 - 1-phenyl-3-(2-pyridyl)pyrazole diimine ligands 192–193
 - phosphorescence (PHOS) 6, 544
 - and TADF mechanisms 62–64
 - and triplet state measurements 438–439
 - decay time 6
 - lifetime 232
 - phosphorescent emitters 61, 199
 - TADF 564
 - phosphorescent OLEDs (PHOLEDs) 120, 209, 257, 297, 493
 - phosphorescent red and green emitting materials 199
 - photoexcitation
 - conditions for efficient TADF 239–240
 - excited-state cycles 235–238
 - fluorescence and phosphorescence decays 232–233
 - rate equations 232
 - steady-state fluorescence and phosphorescence intensities 233–234
 - TADF on-set temperature 238
 - photoluminescence of DPTZ-DBTO2 450
 - photoluminescence of exciplexes 336–338
 - photoluminescence quantum efficiency (PLQY) 545
 - photoluminescence quantum yield (PLQY) 177, 181, 348, 382
 - photophysical properties, of emitters 505
 - photophysics of TADF
 - absorption of PTZ-DBTO2 431
 - absorptivity 437
 - 2,7-bis(phenoxazin-10-yl)-9,9-dimethylthioxanthene-*S,S*-dioxide 431
 - broad Gaussian profile 432
 - chemical structure of FASAF 433
 - Coulomb energy 434
 - dibenzothiophene-*S,S*-dioxide 429
 - dual fluorescence 434
 - dynamical processes 455–457

- electrochemical cyclo-voltammetry measurements 437
 electron transfer 437
 emission spectra of DPTZ-DBTO2 437
 emission Stokes-shift 430
 emitter host interactions 457–459
 energy diagram 459
 energy gap law 436
 excitation power 437
 excited state properties 452–455
 excited states involved 450–451
 fluorene units 429
 Gaussian band shape 430
 intensified CCD cameras 428
 ionisation potential 434
 kinetics of the ^1CT prompt state 449–450
 Lippert-Mataga plot 433
 next generation materials 428
 nonpolar solvents 435
 $n-\pi^*/\pi-\pi^*$ mixed state transitions 431
 optical absorption spectra 431
 phenothiazine-dibenzothiophene-*S,S*-dioxide (PTZ-DBTO2) 429
 phosphorescence and triplet state measurements 438–439
 photoluminescence quantum yield 438
 picosecond and nanosecond fluorescence lifetime measurement 428
 polar solvents 435
 PTZ-DBTO2 absorption spectrum 430
 red edge absorption 436
 solvatochromism 433, 435
 solvent orientation polarizability parameter 433
 solvent polarity 436
 standard integrating sphere measurement 437
 temperature dependent emission of DCz-DBTO2 434
 time correlated singlet photon counting techniques 428
 time resolved emission in solid state 446–448
 time resolved emission in solution 440–446
 twisting mechanism 434
 X-ray crystal structure of DPO-TXO2 432
 platinum group metals (PGM) 549
 PMMA-doped emitter 36
 4P-NPB 486, 487
 polarized angular dependent emission spectroscopy 206, 207
 polar polyethylene oxide (PEO) 322
 polar solvents 280, 285, 435, 437, 471, 514
 polycyclic aromatic hydrocarbons 245
 polymer-based OLED 199
 poly(para-phenylenevinylene) 297
 polystyrene (PS) host matrix 533
 poly(vinyl carbazole) (PVK) 133
 potential energy surfaces 304–309
 p-polarized angular dependent emission pattern 211
 prompt and delayed fluorescence intensities 245–248
 prompt fluorescence (PF) 234, 300
 prompt versus delayed fluorescence 347–350
 prototype OLEDs 142
 pseudo-equilibrium constant 63
 pseudo Jahn–Teller distortion 313
 pseudo-Jahn-Teller (PJT) effect 68, 125, 313
 [Pt(0)(binap)₂] 84
 Pt(0) complexes 84
 P-type delayed fluorescence 426
 Purcell effect 202, 205, 208
 Purcell factor 205, 208, 353, 358
 2-(2'-pyridyl)benzimidazole ligands 137, 139
 2-(2'-pyridyl)benzimidazole unit 139
 2-(2'-pyridyl)benzimidazolylbenzene 183, 184
 2-(2'-pyridyl)benzimidazolyl diimine ligands 183, 185
 2-(2'-pyridyl)benzimidazolyl ligand 184

- 1-(2-pyridyl)-pyrazole based ligands 189–190
- 2-(2-pyridyl)-pyrrolide based ligands 188–189
- 1-(2-pyridyl)-pyrazole (czpypz) based diimine ligand 191
- 5-(2-pyridyl)tetrazolate (NN13) 140, 185–187
- 5-(2-pyridyl)tetrazole based ligands 185–187
- 3-(2'-pyridyl)-1,2,4-triazole based ligands 187–188
- 3-(2'-pyridyl)-1,2,4-triazole diimine ligands 187–188
- pyrimidine 360, 365, 400
- PyrPHOS dinuclear complexes 83
- q**
- quadratic vibronic coupling (QVC) model 307
- quantum mechanical considerations 6, 27, 32–34, 45
- quantum-well (QW) 481–483, 485
- quantum yield of triplet formation 230, 410
- quinoxaline 360, 365, 366, 404, 525
- r**
- radiative decay rates 18, 19, 21, 32, 33, 38, 45, 205, 274, 275, 298, 300, 323, 382, 452, 467, 468, 477, 524, 561
- radiative exciton fraction 202–204, 206, 208–213, 218, 219, 221, 222
- radiative fluorescence rate $k_r(S_1 \rightarrow S_0)$ 7
- radiative quantum efficiency (RQE) 202, 210, 211, 213, 217
- radiative rate constants 62–64, 69, 107, 108, 119, 121, 128, 266–269
- radiative singlet-singlet rate $k_r(S_1 \rightarrow S_0)$ 30
- radiative $S_1 \rightarrow S_0$ rate 20
- rate equations 232, 240, 263, 347
- Rayleigh-Schödinger perturbation 269
- refractive index 21, 33, 64, 208, 340, 341, 390
- reverse intersystem crossing (rISC) 5, 63, 120, 258, 302, 426, 467, 502
- PTZ-DBTO2 316–322
- triplet-to-singlet transformation 377, 378
- roll-off effect 3, 120
- Rosenberg-Parker method 245, 249
- $[Ru(bpy)_3]^{2+}$ (bpy = 2, 2'-bipyridine) 120
- s**
- saturation/triplet-polaron quenching 3
- sensitizers 331, 356–360, 377, 382–383, 399, 410, 416, 417
- single particle density operators (SPDO's) 312
- singlet and triplet excitons 2, 93, 117, 120, 121, 178, 298, 377, 381, 382, 466, 486, 487, 495, 502, 560
- singlet exciton sensitizers 383, 416
- singlet harvesting mechanism 2, 5
- slow spin-lattice relaxation (SLR) 26
- small energy separation $\Delta E(S_1-T_1)$ 2, 3, 32
- small-molecule OLEDs 257
- small splitting $\Delta E(S_1-T_1)$ 8
- small zero-field splittings (ZFS) 26
- solid state lighting (SSL) 501, 502
- solution-processed blue TADF materials and devices 504–512
- solution-processed green TADF materials and devices 512–523
- solution-processed TADF polymers and dendrimers
- non-TADF monomer, TADF macromolecule 532
- phenothiazine donors 531
- TTA 528
- solution-processed yellow-to-red TADF materials and devices 523–526
- solvatochromism 341, 433, 435, 471
- solvent orientation polarizability parameter 433
- solvent-separated radical ion pair (SSRIP) 342–344

- spectral shift 19, 64, 79, 150, 350–352, 414
- spin angular momentum 63
- spin-orbit coupling (SOC) 1, 23–27, 62, 63, 70, 81, 83, 85, 120, 177, 202, 259–262, 297, 302, 332, 381, 466
- spin-orbit coupling matrix element (SOCME) 260–262, 264, 268, 269, 276, 277, 281, 286, 302, 315, 316, 318
- spin-coating or blade-coating 528
- spin-dependent interaction operator 259
- spin-lattice relaxation (SLR) 4, 28
- spin-orbit coupling (SOC) 1–4, 6, 23, 26, 27, 44, 46, 120, 131, 157, 177, 178, 259–263, 277, 284, 297, 302, 332, 345, 454, 466, 502
- $S_1 \rightarrow S_0$ fluorescence 3, 30
- steady-state data analysis 245–249
- steady-state fluorescence and phosphorescence intensities 233
- S–T energy gaps 121, 122, 214, 258, 259, 262, 270, 279, 280, 427, 495
- sterically encumbered phenanthroline ligands 179
- Stokes shift 69, 70, 80–82, 85, 140, 190, 263, 280, 430
- Strickler-Berg equation 64
- Strickler-Berg relation 20–22
- strongly green–blue emitting [Cu(pypz)(POP)]⁺ 78
- substituted phen ligands 34
- synthesized heteroleptic d¹⁰ metal complexes 79
- t**
- TADF-type heteroleptic copper(I) emitters 142
- TADF-type OLEDs 121, 144, 152, 157
- Tamm–Dancoff approximation (TDA) 266, 316
- TB-1PXZ 510, 511, 522
- temperature dependence, emission decay time and TADF 28–30
- temperature-dependent radiative constants 64
- 2,4,5,6-tetra(carbazol-9-yl)-1,3-dicyanobenzene (4CzIPN) 512
- 1,2,4,5-tetracyanobenzene (TCB) 342, 343
- tetrahedral Cu(I) complexes 122, 123, 127
- tetrahedral Cu(I) complexes with the LUMO on the PP ligand 129, 142–145
- tetrahedral d¹⁰ copper (I) complexes 160
- tetrahedral four-coordinate Cu(I) complexes 79–80
- tetrahedral TADF-type Cu(I) complexes 129
- 1,2,3,5-tetrakis(carbazol-9-yl)-4,6-dicyanobenzene (4CzIPN) 279–282
- tetra-ter-butylperylene (TBPe) 285
- thermally activated delayed fluorescence (TADF) 199, 377, 502
- approaches 559–566
- blue OLEDs
- academia 552–554
- efficient, stable 550–552
- chromophore 529
- Cu-complex 379
- data analysis
- combined steady-state and decay data 250–252
- decay data 249–250
- steady-state data 245–249
- development of 377, 379
- diphenyl ketone 408–410
- diphenyl sulfoxide 405–407
- donor-acceptor (D-A) molecular systems 377
- emitter materials, OLED 544–547
- emitters 554, 555
- copper (I) complexes 269
- emitter-host interactions 258
- excitation energies and radiative rate constants 266
- intersystem crossing 259
- metal-free TADF emitters 277

- thermally activated delayed fluorescence (TADF) (*contd.*)
- metal-to-ligand charge-transfer (MLCT) 259
 - singlet and triplet-coupled
 - open-shell configuration 258
 - spin-orbit coupling (SOC) 259
 - thermally stable dyes 258
- energy diagram of 467
- energy transfer 561
- eosin 379
- fluorescent materials 563–564
- intramolecular D-A molecules 413
- kinetics
- Arrhenius equation 229
 - Boltzmann-weighted average 230
 - complex schemes 244
 - direct ISC 229
 - electrical excitation 240–244
 - excitation types 231
 - photoexcitation 232
 - quantum yield of triplet formation 230
 - radiative rate constants 229
 - rISC 229
 - three-state scheme 229
- luminescent efficiency of 389–391
- material design
- Ag(I)-based TADF compounds 34–36
 - Boltzmann constant 2
 - Cu(I) complex with large $\Delta E(S_1-T_1)$ 15–22
 - Cu(I) complex with small $\Delta E(S_1-T_1)$ 22–30
 - energy separation $\Delta E(S_1-T_1)$ 30
 - Franck-Condon factors 2
 - intersystem crossing 1
 - ligand-to-ligand charge transfer (LL'CT) 3
 - metal-to-ligand charge transfer 3
 - molecular parameters, and
 - diversity of materials 4–15
 - photocatalysis 1
 - photoluminescence quantum yields 2
 - photon generation 2
 - photophysical principles 1
 - relaxation properties 1
 - saturation/triplet-polaron quenching 3
 - $S_1 \rightarrow S_0$ fluorescence 3
 - $S_1 \rightarrow S_0$ fluorescence rate 30–34
 - singlet harvesting mechanism 2
 - spin-orbit coupling 1
 - triplet emitters 2
 - triplet harvesting effect 2
 - zero-field splitting 3
- OLEDs 379, 380, 467
- on-set temperature 238
- organoboron 411–412
- phosphorescent emitters 490–495, 564–566
- polymers 412
- processing aspects 547–549
- with quantum-well structure 481–486
- singlet-triplet energy splitting 386–389
- sustainability aspects 549–550
- TADF emitters 562–563
- x-bridged diphenyl ketone 410–411
- x-bridged diphenyl sulfoxide 407–408
- thermally stable dyes 258
- thiadiazole 401, 402
- thiones 245
- thioxanthone (TX) 465
- OLEDs 465–467
 - phosphorescent emitters 490–495
 - TADF emitters with QW 481–486
 - TX-based TADF emitters 468–490
 - WOLED based 4P-NPB 486–490
- 9-H-thioxanthene-9-one-10,10-dioxide (TXO) 469
- third generation-emitter materials 199, 222
- three-coordinate Cu(I)
- complexes 79, 80, 147, 155
 - phenanthroline complexes 80
- three-coordinated
- Cu(I)-NHC-phenanthroline complex 270–275

- time correlated singlet photon counting techniques 428
- time-dependent density functional theory (TDDFT) 16, 64, 81, 266–270, 276–283, 286, 316, 318, 475
- time-dependent Hartree (TDH) method 310
- time-dependent Schrödinger equation 303
- time-dependent self-consistent field (TDSCF) 310, 311
- time-independent electronic Schrödinger equation 303
- time-resolved EL and PL spectroscopy 207
- time-resolved electroluminescence (EL) 207, 221
- time resolved emission in solid state 446–448
- time resolved emission in solution 440–446
- time resolved fluorescence decay of DPO-TXO2 441, 442
- time resolved gated emission measurements 428
- time resolved normalized emission spectra of DDMA-TXO2 449
- transition dipole moment 23, 32, 62, 63, 204, 205, 210, 219, 222, 269, 276, 285, 297, 340, 358, 384, 390, 525
- transition dipole vector (TDV) 200, 201
- transition electric dipole moment (M), 64
- triazine derivatives 396–400
- triazolylpyridine ligand N24–N27 141
- 4,4',4''-tri(9-carbazoyl)triphenylamine (TCTA) 481
- triphenylamine (TPA) 395, 401, 403, 412, 469
- triphenylphosphine (PPh₃) 69, 79, 180–182
- triplet emitters 2
- triplet exciplex 127, 345–348, 358, 386, 414, 415
- triplet exciton-polaron quenching (TPQ) 383
- triplet fusion (TF) 278, 425
- triplet harvesting effect 2
- triplet lifetime 120, 313, 443, 444
- triplet relaxation processes 230
- triplet state T₁ and spin-orbit coupling 23–27
- triplet–triplet annihilation (TTA) 8, 104, 120, 178, 199, 356, 383, 425, 502, 528
- 4,4',4''-tris(*N*-carbazolyl)-triphenylamine (TCTA) 332, 414
- 1,3,5-tris(*N*-phenylbenzimidazole-2-yl)benzene (TPBI) 477, 478
- tris- or bis(cyclometalated) iridium(III) complexes 120
- T₁→S₀ and S₁→S₀ decay processes 4
- Tully's Trajectory Surface Hopping (TSH) 309
- twisted intramolecular charge transfer (TICT) 63, 471
- twisting mechanism 434
- two-emitting states model 121
- TX-based TADF emitters
- aggregation-induced emission 472
 - delayed component 477
 - high radiative decay rate 469
 - mCP 474
 - OLED structure 475
 - TCTA 481
 - temperature dependence 474
 - TICT 471
 - TPBI 477
 - TXO-TPA and TXO-PhCz 469, 471
- TXO-PhCz 471
- emission characteristics 470
 - mCP 485
 - molecular structure 478
 - TADF 482
 - transient PL 478
- TXO-PhCz:TPBI 477–480
- TXO-TPA
- absorption spectra 471
 - AIE 472
 - blue fluorescent emitter 486

- TXO-TPA (*contd.*)
- emitting layer 475
 - mCP 474, 486
 - molecular structures 468
 - phosphorescent 471
 - single crystal structure 473
 - TCTA 481
 - temperature dependence 474
 - TPBI 477
 - transient PL 475
 - triplet energy 471
 - UV-vis absorption 472
 - TXO-TPA:mCP 474, 476, 486
- u**
- ultrafast spin-vibronic dynamics 315
- v**
- vacuum thermal evaporation (VTE) 547
 - vapor-deposited OLEDs 81, 155, 157
 - variational multi-configurational Gaussian wavepacket (vMCG) 310
 - vibrational thermalisation 300
 - vibrational wavefunctions, overlap of 262
 - vibronic coupling
 - adiabatic electronic energies 299
 - Born–Oppenheimer approximation 299
 - delayed fluorescence 300–301
 - density matrix formalism of MCTDH 311–312
 - electron exchange interaction 298
 - electronic ground state 298
 - Fermi's Golden rule (FGR) approach 303
 - first order mixing coefficient 297
 - ground and excited-state wavefunctions 298
 - model Hamiltonian 306–309
 - multi-configurational time-dependent Hartree approach 310–311
 - perturbation theory 303
 - poly(para-phenylenevinylene) 297
 - potential energy surfaces 304–306
 - radiative rate 297
 - rISC 302–303
 - rISC of PTZ-DBTO2 316–322
 - singlet and triplet excitons 298
 - systematic material design 299
 - time-dependent Schrödinger equation 303
 - time-independent electronic Schrödinger equation 303
 - Tully's Trajectory Surface Hopping (TSH) 309
 - ultrafast dynamics of a Cu(I)-phenanthroline complex 313–316
- w**
- white organic light emitting diodes (WOLED) 486, 488, 490
 - work function (WF) 200, 515
- x**
- xanthene dyes 245
 - X-ray crystallographic study 101–102
 - X-ray crystal structure of DPO-TXO2 432
- z**
- Zeonex 237, 238, 245–251, 438, 446, 448, 450, 452, 457, 458
 - zero-field splitting (ZFS) 3, 6, 7, 26, 70, 120, 230, 260