

Errata for “Magnetic Resonance Elastography - Physical Background and Medical Applications” by S. Hirsch, J. Braun, and I. Sack, Wiley-VCH.

Corrections as of March 29, 2017.

If you find any errors, the authors kindly ask you to report them to mre-book@charite.de so that they can be corrected here.

Chapter 4

- Page 91: in the top row of fig. 4.7, S' should be S^* instead.

Chapter 10

- Section 10.3.1, Page 207, 4th line: Romano et al published the first inversion method capable of retrieving multiple model parameters from an MRE scan with a sufficient number of independent measurements [see the three references listed below]. The method is based on a variational formulation of the wave equation for determining both Lamé parameters. More details to the variational formulation are given in chapter 10.7. Similar to the work of Romano et al., a multi-parametric direct inversion method termed AIDE (algebraic inversion of the differential equation) was introduced by Oliphant et. al [215].
 - Romano, A. J., Shirron, J. J., and Bucaro, J. A. (1998). On the noninvasive determination of material parameters from a knowledge of elastic displacements theory and numerical simulation. *IEEE Transactions on Ultrasonics, Ferroelectrics, and Frequency Control*, 45(3), 751–9, doi: 10.1109/58.677725
 - Romano, A. J., Bucaro, J. A., Ehman, R. L., and Shirron, J. J. (2000). Evaluation of a material parameter extraction algorithm using MRI-based displacement measurements. *IEEE Transactions on Ultrasonics, Ferroelectrics, and Frequency Control*, 47(6), 1575–81, doi: 10.1109/58.883546
 - Romano, A. J., Bucaro, J. A., and Ehman, R. L. (2000). The application of a material parameter extraction algorithm to MRI-based displacement measurements. *The Journal of the Acoustical Society of America*, 107(5), doi: 10.1121/1.428931

Chapter 13

- Section 13.1.2: all instances of ε without subscript should be typeset as a tensor: $\boldsymbol{\varepsilon}$

References

The following references need to be corrected:

- [47]: Guo, J., Hirsch, S., Scheel, M., Braun, J., and Sack, I. (2016). Three-parameter shear wave inversion in MR elastography of incompressible transverse Isotropic media: application to in vivo lower leg muscles. *Magnetic Resonance in Medicine*, 75(4), 1537–45. doi: 10.1002/mrm.25740
- [69]: Dittmann, F., Hirsch, S., Tzschätzsch, H., Guo, J., Braun, J., and Sack, I. (2016). In vivo wideband multifrequency MR elastography of the human brain and liver. *Magnetic Resonance in Medicine*, 76(4), 1116–26, doi: 10.1002/mrm.26006
- [210]: Barnhill, E., Kennedy, P., Johnson, C. L., Mada, M., & Roberts, N. (2015). Real-time 4D phase unwrapping applied to magnetic resonance elastography. *Magnetic Resonance in Medicine*, 73(6), 2321–31, doi: 10.1002/mrm.25332
- [304]: Kolipaka, A., Woodrum, D., Araoz, P. A., and Ehman, R. L. (2012). MR elastography of the in vivo abdominal aorta: A feasibility study for comparing aortic stiffness between hypertensives and normotensives. *Journal of Magnetic Resonance Imaging*, 35(3), 582–86. doi: 10.1002/jmri.22866
- [314]: Kolipaka, A., Aggarwal, S. R., McGee, K. P., Anavekar, N., Manduca, A., Ehman, R. L., and Araoz, P. A. (2012). Magnetic resonance elastography as a method to estimate myocardial contractility. *Journal of Magnetic Resonance Imaging*, 36(1), 120–7. doi: 10.1002/jmri.23616
- [353]: Romano, A. J., Guo, J., Prokscha, T., Meyer, T., Hirsch, S., Braun, J., Sack, I., and Scheel, M. (2014). In vivo waveguide elastography: effects of neurodegeneration in patients with amyotrophic lateral sclerosis. *Magnetic Resonance in Medicine*, 72(6), 1755–61, doi: 10.1002/mrm.25067
- [585]: Sahebjavaher, R. S., Baghani, A., Honarvar, M., Sinkus, R., and Salcudean, S. E. (2012). Transperineal prostate MR elastography: Initial in vivo results. *Magnetic Resonance in Medicine*, 69(2), 411–20. doi: 10.1002/mrm.24268