

Index

a

- achiral iron porphyrin catalysts,
 165–172
 adamantane, 377, 381
 1-adamantanol, 377
 aerobic dehydrogenative coupling
 reaction, 74
 AH *see* asymmetric hydrogenation (AH)
 AHS *see* asymmetric hydrosilylation
 (AHS)
 alcohols, hydrogen generation
 energetic application, 469–470
 organic compound synthesis,
 470–473
 alkane hydroxylation, 381
 alkene cyclopropanation mechanism,
 166
 alkenyl-heteroarenes, 196–205
 alkyl–alkyl cross-coupling, 279
 alkyl bis(imino)pyridine cobalt complex,
 108–109
 alkylboronic acids coupling, 332
 α - β -unsaturated hydrazones, 336
 aluminum-(5,10,15,20-
 tetraphenylporphyrin) (Al(TPP)),
 538
 amino-bis(phenolate) Fe^{III} catalyst
 precursor, 138
 aminophenoxide-based Co complexes,
 542–544
 aminopyridine manganese complexes,
 357
 aminosalen-type *O,N,N*-pincer
 nickel(II) complexes, 119–120
 ammonia borane, 476
 ammonia, dinitrogen reduction
 catalytic ammonia formation,
 559–569
 Haber–Bosch-process, 551–555
 iron–sulfur systems, 557–559
 metal complexes, 556–557
 nitrogenase-inspired systems,
 555–556
 annulation, of alkene, 397
 anti-Markovnikov addition, 392
 anti-Markovnikov hydrosilylation
 reaction, 151–154, 157, 159
 aromatic hydroxylation method, 382
 2-arylpiperidines, 392, 394, 412
 asymmetric borohydride reduction
 cobalt catalysts, 226–227
 nickel catalysts, 230
 asymmetric conjugate addition (ACA)
 reactions, 192, 195, 198
 asymmetric hydrosilylation (AHS)
 cobalt catalysts, 225–226
 copper catalysts, 232–235
 iron catalysts, 220–223
 nickel catalysts, 229, 230
 asymmetric intramolecular
 cyclopropanation, of indoles, 183
 asymmetric olefin epoxidation, 360
 asymmetric transfer hydrogenation
 (ATH)
 cobalt catalysts, 224
 copper catalysts, 232
 iron catalysts, 213–218
 Mn(I) catalysts, 211–212
 nickel catalysts, 228–229

ATH *see* asymmetric transfer
hydrogenation (ATH)
1,2-azido alcohols, 139
azines, 73

b

Baran hydroamination reaction, 138
base metal cooperative catalysis, 18–21
 β -diketiminato ligand, 554
 β -diketiminatocopper(II) complexes,
383
Bell–Evans–Polanyi (BEP) principle, 34,
35
bent tri-iron complexes, 500
benzamide substrates, 396
 β -hydride elimination, 465
1,1'-bibenzimidazole, 72
bidentate bis(phosphine)cobalt
catalysts, 117
bidentate dimethylglyoxime ligands, 541
bidentate diphosphine nickel(II)
complex, 121, 122
bimetallic cobalt(II) diacetato
complexes, 542
bimetallic reaction catalysis
alkyne cycloadditions, 58–59
binding and activation, 50–51
C=O cleavage products, 57
C–X activation reactions, 56
E–H addition and elimination
reactions, 54–55
oxidative addition and reductive
elimination reactions, 54
bioinspired Cu^{II}–thiophenol catalytic
system, 17
bipyrrolidine ligand, 378
bis(amino)amido pincer-ligated Ni^{II}
catalyst, 154, 155
bis(arylimidazol-2-ylidene)pyridine
cobalt methyl, 112
bis(2-dimethylaminoethyl)ether
(BDMAE), 408
bis(imino)pyridine (BIP) ligand, 8–14
bis(phosphino)boranecobalt(I) complex,
114

Born–Oppenheimer (BO)
approximation, 34
borohydridonickel(II) hydride, 120–121
boronates, 332
boronic acids, 332

c

carbene radicals, 22–24
carbene transfer reactions, 164, 172, 186
carbon dioxide (CO₂)
Cu-catalyzed homogeneous
hydrogenation, 84–86
hydroboration, 259–260
hydrosilylation, 258–259
catalytic ammonia formation
cyclic aminocarbene iron system,
567–568
diphosphine iron system, 568–569
iron and cobalt PNP systems, 566–567
tripodal systems, 560–566
catalytic hydride reduction, 83
catalytic hydrogenation, 83
catalytic transfer hydrocyanation
reaction, 148, 150
catechol-derived redox non-innocent
ligands, 14–15
C=C bond catalytic oxygenation
cobalt, nickel, and copper catalysts,
372–375
iron catalysts, 363–372
manganese catalysts, 356–363
C=C hydrogenations
cobalt catalysts, 107–118
iron catalysts, 99–107
nickel catalysts, 118–122
Chalk–Harrod mechanism, 151, 153
C–H amination
activated amine sources, 409–412
with unactivated amines, 409
C–H bond oxidation
cobalt catalysts, 380–381
copper catalysts, 383–384
iron catalysts, 377–380
manganese catalysts, 376–377
nickel catalysts, 381–383

- chelation-assisted C—H
 - functionalization
 - alkylation, 407–409
 - alkynylations, 403
 - C—C bond formation via C—H
 - activation, 392–409
 - C—C multiple bonds, 392–393
 - C—H allylation, 397
 - C—halogen formation, 412–415
 - C—H arylation, 404–406
 - C—H cyanation routes, 404
 - C—heteroatom formation, 409–415
- chemoselective iron-catalyzed
 - hydrosilylation
 - of carboxamides, 252–253
 - of carboxylic acids, 257–258
 - of carboxylic esters, 254, 256
 - of primary amides, 253–254
 - of ureas, 253–255
- C—heteroatom multiple bonds,
 - 393–396
 - aldehydes, addition to, 394–396
 - imines, addition to, 393–394
 - isocyanates, addition to, 396
- C—H functionalization, 299–308,
 - 312–319
- chiral [3.1.0]bicycloalkane lactones,
 - 182–183
- chiral bis(imino)pyridine methyl cobalt,
 - 109, 110
- chiral iron porphyrin catalysts, 172–176
- chiral spiro-*bis*oxazoline ligands, 181,
 - 183
- cis*- β -methylstyrene, 367
- cis*-3,4-dihydroisoquinolines, 400
- cis*-1,2-dimethylcyclohexane, 377
- cobalt catalysts
 - C=C bond catalytic oxygenation,
 - 372–375
 - C—H bond oxidation, 380–381
 - coupling reactions, of CO₂
 - aminophenoxide-based Co
 - complexes, 542–544
 - cobalt–salen complexes, 530–537
 - Co–porphyrins, 537–540
 - N₄ ligands, 540–541
 - cross-couplings
 - C—H functionalization, 299–308,
 - 312–319
 - preactivated substrate approach,
 - 308–312, 320–322
 - hydroazidation reaction, 130–132
 - hydrochlorination reaction, 133–134
 - hydrocyanation reaction, 133
 - hydrohydrazination reaction,
 - 128–129
 - hydrosilylations, of aldehydes and
 - ketones, 247–248
 - Markovnikov hydroalkoxylation
 - reaction, 135–136
 - radical hydrofluorination reaction,
 - 135
 - silylamines, reduction of N₂, 572–575
- cobalt–salen complexes, 530–537
- cobalt water oxidation catalysts,
 - 440–443
- CO₂ hydrogenation
 - DLPNO-CCSD(T) free energy profile,
 - 40–42
 - hydricity *vs.* reactivity, 43–45
 - mechanistic studies, 38–39
 - motivation behind studying, 35
 - precious metal catalysts, 38
 - reaction equations, 37
- computational chemistry, 33
- controlled catalytic radical-type
 - reactions, 27
- cooperative catalysis, 15–21, 26–27
- Co–porphyrins, 537–540
- copper–boxmi complex, 346
- copper catalysts
 - conjugate addition reactions
 - alkenyl-heteroarenes, 196–205
 - α -substituted α,β -unsaturated
 - carbonyl compounds, 192–196
 - homogeneous hydrogenation
 - carbonyl compounds, 86–89
 - catalytic cycle, 83
 - CO₂, 84–86
 - vs.* heterogeneous catalysts, 82
 - olefins and alkynes, 89–91
 - hydroamination reaction, 143–147

- copper water oxidation catalysts, 445–446
- CO₂ reduction challenges, 35–37
- [Cp(CO)₂Fe^{II}(THF)]BF₄ catalyst
(Cp=cyclopentadienyl;
THF=tetrahydrofuran), 184–186
- cross-coupling reactions, 297
- C*-silyl cyclopropanes, 177, 178
- C*₂-symmetrical iron(III) chiral porphyrins, 176, 177
- Cu-promoted N–N bond formations cyclizations, 74–82
- double bond, 72–74
- single bond, 71–72
- cyclic(alkyl)(amino)carbene (CAAC), 480
- cyclic aminocarbene iron system, 567–568
- cyclohexane, 381
- cyclopentadienyl compounds, 498
- cyclopentadienyl iron complexes, 107
- d**
- dehydrogenation, from ammonia borane and amine boranes, 474–480
- dehydrogenative cyclizations (DCs), 75, 77, 79–80
- density functional theory (DFT) methods, 34, 426, 459, 532
- 1,2-diamino-cyclohexane fragment, 532
- 2,6-dichloroanilinium tetrafluoroborate, 510
- dichloroisobutane (DCIB), 404
- dicobalt(II) macrocycles, 510
- diethylene glycol dimethyl ether, 466
- 1,3-diketone-type ligands, 374
- dimethoxyethyl ether (DME), 572
- dimethylaminoborane (DMAB), 476
- dimethylformamide (DMF), 461
- N, N'*-dimethylpropyleneurea (DMPU), 194
- dinitrogen reduction ammonia, 551–555
- N₂ activation, 500–551
- nitrogenase-inspired systems, 555–556
- silylamines, 569–575
- 1,4-dioxane, 466
- 3,5-diphenyl-1,2,4-triazole, 75, 76
- direct hydrogenation (AH) cobalt catalysts, 223–224
- copper catalysts, 231–232
- iron catalysts, 218–220
- Mn(I) catalysts, 211–212
- nickel catalysts, 228
- directing groups (DGs), 391
- direct reductive amination (DRA), 251–252
- D*₂-symmetrical iron(III) chiral porphyrin, 175
- dual-catalytic Markovnikov hydroarylation, 139, 140
- DuBois system, 20–21
- e**
- eco-friendly catalytic procedures, 163
- electrochemical water oxidation studies, 429
- eliminative cyclizations (ECs), 75, 80–81
- eliminative dehydrogenative cyclization (EDC), 75, 81–82
- enamines, electrophilic trifluoromethylation of, 335
- enantioselective cobalt-catalyzed hydrogenation, 117
- extended X-ray absorption fine spectroscopy (EXAFS), 426
- f**
- Fe(F₂₀TPP)(CPh₂), 167
- Fe and Co pivalate isocyanide-ligated catalyst systems, 157–159
- Fe^{II}(TTP)-catalyzed alkene cyclopropanation (TTP=dianion of *meso*-tetrakis-(4-tolyl) porphyrin), 165
- ferrocene, 492
- fluorinated diglyoxime complexes, 498
- fluorine, 413

formic acid, hydrogen generation from
 aluminum, 467
 iron, 461–466
 nickel, 466–467

g

galactose oxidase (GOase), 16–17
 Grignard reagents, 192, 193, 205, 268,
 285, 392
 group 6 metals, 514–518

h

Haber–Bosch process, 549
 Heck-type, cobalt-catalyzed
 cross-coupling reaction, 310–311
 heteroaromatics, 392
 heterobimetallic catalysis, 60
 heterobimetallic [NiFe] clusters, 505
 heteroleptic arene/alkene cobaltate
 complexes, 115, 116
 hexamethylphosphoramide (HMPA),
 193
 high-valent cobalt catalysis, 302–308
 hindered amine drug intermediates,
 137–138
 homogeneous CO₂ hydrogenation, 35,
 37
 hydrogen atom transfer (HAT), 128,
 129, 137, 138
 hydrogen generation
 from alcohols, 469–473
 carbon and boron nitrogen-based
 substrates, 453–458
 from formic acid, 460–468
 hydrogen storage, in liquid organic
 hydrogen carriers, 473–474

i

imines, 393–394
 1*H*-indazoles, 317
 indoles, 394
 iridium complexes, 339
 iron- and nickel-based catalysts
 bioinspired di-iron molecules,
 490–495

bioinspired [NiFe] complexes and
 [NiMn] analogs, 501–508
 mono- and poly-iron complexes,
 496–501

iron catalysts

aminohydroxylation, 141, 143–145
 C=C bond catalytic oxygenation,
 363–372
 C–C cross-coupling
 NHCs, 276–283
 phosphines, 283–291
 simple ferric salts, 266–273
 TMEDA, 273–276
 C–H bond oxidation, 377–380
 hydromethylation reaction, 137
 hydrosilylations, of aldehydes and
 ketones, 242–247
 silylamines, reduction of N₂, 570–572
 iron corroles, 179, 180
 iron phthalocyanine complexes, 177
 iron–pincer complex system, 19–20
 iron polypyridyl monophenolate
 complexes, 496
 iron(II) tetraaza macrocyclic complexes,
 180

k

kinetic isotope effect (KIE), 441

l

light-harvesting complex 1 (LH1), 426
 light-responsive ligands, 7–8
 local coupled-cluster (L-CC) method, 34
 low-valent cobalt catalysis, 299–302

m

macrocyclic ligand, 543
 maleonitrile dithiolate, 497
 manganese catalysts
 aerobic oxidative hydroxyazidation,
 139–143
 C=C bond catalytic oxygenation,
 356–363
 hydration reaction, 130, 131
 hydrosilylations, of aldehydes and
 ketones, 248–250

- manganese water oxidation catalysts,
430–435
 bioinspired Mn_4O_4 models, 430–432
 biomimetic models, Lewis acid,
 432–433
 Fe–TAML complexes, 436–440
- Markovnikov hydroboration reaction,
157, 159
- Markovnikov hydrosilylation reaction,
155, 157, 158
- meta*-chloroperoxybenzoic acid
(mCPBA), 367
- metal-pendant nitrogen-based system,
20–21
- methylamine borane (MAB), 478
- Michael acceptors, 191, 192, 195, 196
- molybdenum–polypyridyl complexes,
516
- monoanionic polyamine ligands, 437
- Mukaiyama hydration, 128
- multi-iron catalyst, 440
- multimetallic cooperation catalysis,
64–65
- n**
- N_2 activation, 500–551
- N*-chloroamines, 412
- N*-halosuccinimides, 413
- N*-heterocyclic carbenes (NHCs),
276–283, 479
- nickel catalysts
 C=C bond catalytic oxygenation,
 372–375
 C–H bond oxidation, 381–383
 Grignard addition, 203
 Lewis-acid-assisted carbocyanation
 reaction, 147–149
 water oxidation catalysts, 443–444
- nitrene radicals, 25–26
- nitride- and carbide-centered tetrairon
clusters, 501
- nitrogenase-inspired systems, 555–556
- N_4 ligands, 540–541
- noble metals, 2
- non-innocent ligands, 1–2, 26
- N*-(trifluoromethylthio)phtalimide, 347
- nucleophilic trifluoromethylthiolation
copper complexes, 342–344
nickel-based catalytic systems,
344–345
- o**
- olefins
 aerobic epoxidation of, 374
 cis-dihydroxylation, 356
 epoxidation, 359
 stoichiometric cis-dihydroxylation of,
 362
- osotriazoles, 81
- oxazoline iminopyridine cobalt
complex, 110, 111
- oxidants, in water oxidation reactions,
428–429
- oxidative addition (OA), 395
- oxidative C–H olefination, 396–397
- oxygen-evolving complex (OEC),
425–428
- p**
- pair natural orbitals (PNOs), 34
- P,B,P* nickel(I) hydride, 121
- peracetic acid (PAA), 358
- perfluoroalkylation reactions, 348–350
- phenylated iron(II) ferrates, 269, 270
- 2-phenylpyridines, 406
- phosphines, 283–291
- pH-responsive ligands, 5–7
- piano-stool cyclopentadienyl phosphine
iron complexes, 245, 246
- P,N,P*-nickel(II) hydride, 120
- P,N,P* nickel(II) methyl complex, 120,
121
- polar double bond reduction
 AH (*see* asymmetric hydrogenation
 (AH))
 AHS (*see* asymmetric hydrosilylation
 (AHS))
 ATH (*see* asymmetric transfer
 hydrogenation (ATH))
- polyethylene glycol (PEG), 395
- polynuclear catalysis, 64–65
- polypropylene carbonate (PPC), 530

polypyridyl ruthenium, 339
 potassium bis(anthracene) cobaltate,
 115
 potassium bis(anthracene)ferrate, 106
 potential energy surface (PES), 34
 power-to-gas techniques, 453
 preactivated substrate approach,
 308–312, 320–322
 propargylic hydrazines, 129, 130
 proton-coupled electron transfer
 (PCET) mechanism, 36–37, 314
 proton exchange membrane (PEM), 453
 pumiliotoxin C, 192
 pyrazolines, 71

r

Raney-nickel catalysts, 118–119
 rate-determining step (RDS), 34, 441
 reactive bis(perfluoroalkyl)zinc
 reagents, 348
 redox-active ligands, 8–15, 26
 redox-active phosphole ligand, 492
 redox noninnocent ligands, 341
 redox-responsive ligands, 3–5
 regiodivergent hydrosilylation reaction,
 155, 156
 responsive ligands *see*
 stimuli-responsive ligands
 ring-opening cross-coupling, 346
 rivastigmine, 146, 147

s

Sandmeyer trifluoromethylation, 330
 Schiff bases (SB), 184
 sequential asymmetric
 conjugation–enolate trapping,
 195, 196
 silicon electrophiles, 563
 silylamines, reduction of N₂
 cobalt, 572–575
 iron, 570–572
 silyl enol ethers, electrophilic
 trifluoromethylation of, 335
 simple iron salt catalysts, 266–273
 single-electron transfer (SET)
 mechanism, 298, 313

single-site catalysis, 49–50
 binding and activation, 50–51
 oxidative addition and reductive
 elimination reactions, 54
 Sonogashira–Hagihara coupling
 reaction, 403
 steroidal substrates, 380
 stimuli-responsive ligands, 2–8, 26
 stoichiometric hydrofunctionalization
 reaction, 134
 stoichiometric reaction pathways, 50
 substrate radicals, 21–27
 Suzuki–Miyaura coupling reaction, 291
 synthetic models, for water oxidation,
 428

t

tetra-anionic tetra-amido macrocyclic
 ligands (TAML's), 436–437
 tetrabutyl ammonium bromide (TBAB),
 531
 tetrahydrofuran (THF), 560
 tetrakis-Schiff base macrocycles, 511
 tetrameric B-(cyclotriborazanyl)amine-
 borane, 478
 tetrameric cobalt(I) complex, 116
 tetrameric nickel(I) complex, 122
 N,N,N',N'-tetramethylethylenediamine
 (TMEDA), 273–276
 tetra-O-acetyl-β-D-glucopyranoside,
 495
 Togni's reagent II, 339
 transition state theory (TST), 34
 transmetalated iron(II) ferrates, 269, 270
 1,5,7-triazabicyclo-[4,4,0]dec-5-ene
 (TBD), 534
 1,3,5-triaza-7-phosphaadamantane
 (PTA), 492
 2,2,2-trifluoroethanol (TFE), 395
 trifluoromethylated heteroarenes, 332
 trifluoromethylation reactions
 CF₃• radicals, using Langlois' reagent,
 332–333
 copper and electrophilic CF₃⁺
 sources, 333–341
 electrophiles, reaction with, 330–331
 oxidative coupling, 331–332

- trifluoromethylthiolation reactions
 - electrophilic reaction, 345–348
 - nucleophilic reaction, 342–345
 - tri-iron complexes, 500
 - trimeric B-(cyclodiborazanyl)amine-borane, 478
 - tris(2-pyridylmethyl)amine (TPA) ligand, 383, 442
- u**
- Umemoto's reagent, 334–336
- v**
- vinyl-heteroarenes, 200–202
 - vinyl-substituted cyclic carbonates, 544
- w**
- water oxidation
 - cobalt catalysts, 440–443
 - copper catalysts, 445–446
 - manganese catalysts, 430–435
 - molecular iron catalysts, 435–440
 - nickel catalysts, 443–444
 - oxidants, 428–429
 - oxygen-evolving complex (OEC), 425–428
 - synthetic models, 428
- x**
- X-ray crystallography, 426
- z**
- Z-diastereoselective
 - bis-pentafluoroethylthiolation, 349
 - Ziegler-type Co/Al hydrogenation catalysts, 107–108
 - Zn–Salen catalyst, 18