

Index

a

- ABA triblock copolymers 26
 by disulfide formation 787–788
 quasi-triblock copolymers 1054–1055,
 1059–1060
 by radical-radical termination 940
 by sequential monomer addition
 1059–1060
 using symmetrical trithiocarbonates
 382–386, 684, 936, 939, 1055
 using Z-connected bis-RAFT agents
 235–237, 263, 387–389, 936, 1055
- ABC triblock copolymers 26, 632
 by photoinduced energy or electron
 transfer (PET)-RAFT 632
 quasi-ABC triblock copolymers 1057
- AB diblock copolymers, *see* diblock
 copolymers 26
- acenaphthyl chromophore 807
 donor in light-harvesting polymers
 807
- acenaphthylene 242, 287, 314, 398, 422,
 559, 807
- 2-(acetoacetoxy)ethyl methacrylate
 (AAEMA),
 2-(methacryloyloxy)ethyl
 3-oxobutanoate) 240, 716, 847,
 849
- acetone oxime (AO) group 830, 836–837
 active ester 830, 836–837
 functional monomer 837
 reaction with amino groups 836–837
- acrylamide(s) 43–46, 280, 284–285,
 388–389, 391–412, 419–421, 503,
 510, 511, 572, 624, 650, 684, 691,
see also acrylamide (Am),
 2-acrylamido-2-methylpropanesulfonate sodium salt
 (AMPS), *N*-acryloylmorpholine
 (NAM), *N,N*-diethylacrylamide,
N,N-dimethylacrylamide
 (DMAm), *N*-(2-hydroxyethyl)
 acrylamide (HEAm),
N-isopropylacrylamide (NIPAm)
N-(2-(acetoacetoxy)ethyl)acrylamide
 (AAEAm) 849
- aqueous RAFT 684, 686, 701
- di(BOC)-acrylamide 838
- with dithiocarbamates 569, 572
- with dithioesters 241, 244, 249, 260,
 280, 284–285
- experimental procedures 48
 acrylamide in solution 46
N,N-diethylacrylamide is solution
 46
N,N-dimethylacrylamide is solution
 46
- functional acrylamides
 active ester 426
 alkyne 426
 azide 289
 betaine 45
 bisphosphonate 45
 hydroxyl, *see N*-(2-hydroxyethyl)
 acrylamide (HEAm)

- acrylamide(s) (*contd.*)
- protected thiol 427
 - protected amine 45, 293, 427, 428
 - with functional trithiocarbonates 402–412
 - N*-isopropylacrylamide in solution 46
 - photoinduced energy or electron transfer (PET)-RAFT 691
 - photoRAFT 686
 - RAFT agent selection 27–28, 43–45, 112
 - redox RAFT 648, 650
 - with thiocarbonates 383, 384, 385, 394, 398, 399, 400, 404, 650
 - with trithiocarbonates 388–389, 391–412, 419–420, 421
 - with xanthates 496, 503, 510, 511, 516
- 2-acrylamido-2-methylpropanesulfonic acid sodium salt (AMPS) 128, 244, 284, 317, 385, 386, 391, 394, 421, 511, 532, 681–682
- acrylates 38, 41, 240, 281, 685, 70, *see also* butyl acrylate, 2-ethylhexyl acrylate, methyl acrylate, octadecyl acrylate (oligo(ethylene glycol) monomethyl ether) acrylate, (poly(ethylene glycol) monomethyl ether) acrylate
- copolymers 48
 - with dithiocarbamates 559–565, 572
 - with dithioesters 240, 281–283
 - experimental procedures
 - acrylic acid in solution 43
 - butyl acrylate in solution 41
 - t*-butyl acrylate in solution 43
 - butyl acrylate in surfactant-free emulsion 42
 - 3,5-dibromobenzyl acrylate in solution 42
 - 2-ethylhexyl acrylate in solution 42
 - methyl acrylate in solution 51
 - methyl acrylate block *N*-vinylcarbazole in solution 51
 - octadecyl acrylate in solution 42
 - (oligo(ethylene glycol) monomethyl ether) acrylate in solution 43
 - functional acrylates
 - active ester 290, 426
 - alkyne 290, 426
 - ATRP inimer 295, 429
 - isocyanate 294
 - NMP inimer 296, 429
 - protected thiol 292, 427
 - protected amine 292, 293, 427
 - with functional trithiocarbonates 402–412
 - RAFT agent selection 27–28, 112
 - surfactant-free emulsion
 - polymerization 42
 - with trithiocarbonates 388–389, 391–412, 419–421
 - with xanthates 503, 510
- acrylic acid (AA) 43, 147, 209, 711, 787, 831
- experimental procedure 43
 - NHS ester, *see N*-acryloyloxysuccinimide
- acrylonitrile (AN)
- blocks 241, 260
 - copolymers 209, 241, 247, 249, 260, 383–385, 391, 397, 399, 400, 408, 569
 - acrylonitrile/butadiene 209, 249, 264, 385, 393, 399, 408
 - acrylonitrile/maleic anhydride 883
 - acrylonitrile/methacrylic acid 209
 - acrylonitrile/styrene 241, 249
 - with dithiocarbamates 552
 - with dithioesters 112, 234, 242, 244, 245, 247
 - homopolymer 242, 244, 245, 247, 383–385
 - propagating radicals 666
 - RAFT agent selection 28, 112
 - rate coefficients for radical addition (k_i, k_p) 107
 - with trithiocarbonates 112, 380, 383–385

- 4-acryloxyphenyldimethylsulfonium triflate (APDMST) 837
- N*-(2-acryloyloxyethyl)pyrrolidone (NAP) 685
- acryloyl chloride (AC) 780, 785, 839
- N*-acryloylmorpholine (NAM) 280, 421, 509, 522, 572, 653, 693, 715, 782, 832, 849, 1054
- with dithiocarbamates 563, 572
- with dithioesters 242, 245, 248, 260, 280
- end-group removal 444
- with trithiocarbonates 384, 391, 393, 396, 398, 399, 410, 421, 444
- 2-(acryloyloxy)ethylisocyanate, *see* 2-isocyanatoethyl acrylate
- N*-acryloyloxysuccinimide (acrylic acid, NHS ester) 242, 244, 245, 249, 290, 383, 396, 400, 404, 426, 831
- with dithioesters 242, 244, 245, 249
- with trithiocarbonates 383, 396, 400, 404, 407
- activators re-generated by electron transfer (ARGET)-ATRP
- concurrent RAFT-ATRP 658–689
- active ester 830–836, *see also* acetone oxime (AO) group, *p*-dialkylsulfonium phenoxy ester (DASPE), hexafluoroisopropanol ester, *N*-hydroxysuccinimide (NHS) ester, nitrophenyl ester, pentafluorobenzyl (PFB) ester, pentafluorophenyl (PFP) ester, salicylic acid ester
- acyl chloride 580, 830, 839
- acryloyl chloride 780–781, 839
- functional monomers 288, 290, 425, 426, 830–836, 1004
- functional RAFT agents 251, 252, 402, 404–405, 761
- intermediate in ketoform reaction 580
- methacryloyl chloride 839
- 4-vinylbenzoyl chloride 839
- addition-fragmentation coupling 766–768
- addition rate coefficient, *see* rate coefficients
- adipic acid dihydrazide (ADH) 716, 724
- aliphatic dithioesters 32, 38, 223, 236–238, 313–314, 325, *see also* 2-phenylpropan-2-yl 2-phenylethanedithioate
- alkoxyamines 205, 296, 308–310, 403, 438, 429, 717, 737, 768
- addition-fragmentation coupling 768
- ESARA 310, 438, 768
- NMP inimer 429
- RAFT-F-NMP 308–310, 438
- alkyl diethyldithiocarbamate (DC) 656–657
- initiators 668
- alkyl halide 308–309, 413, 436, 502, 579, 656, 659–660, 839–840, 964
- alkyl pseudohalides 656–658, 668–669
- alkyne group 830, 847, 848, 857–858, *see also* yne
- alkyne-functional monomer 288–289, 426, *see also* propargyl acrylate, propargyl methacrylate
- alkyne-functional RAFT agent 235, 253–255, 407–408, 758–759
- copper-catalyzed azide–alkyne cycloaddition 288, 717, 757–759, 847, 857–858
- strain-promoted azide–alkyne cycloaddition 848
- thiol–yne reactions 776–780
- allyl ether 775
- O*-allyl hydroxylamine 724
- allyl isocyanate 840–841
- allyl methacrylate 38, 416, 779
- aluminium naphthalocyanine (AlNc) 621–622
- aluminium phthalocyanine (AlPc) 612, 621, 623
- 2-aminoethanethiol 857
- N*-(2-aminoethyl)acrylamide hydrochloride (AEAm) 724

- N*-(2-aminoethyl)methacrylamide
hydrochloride (AEMAm) 240, 244, 293, 420, 428
- (2-aminoethyl) methacrylate
hydrochloride (AEMA) 129, 242–244, 253, 293, 383, 428, 846
copolymers 846
- 3-amino-1,2-propanediol 853
- N*-(3-aminopropyl)acrylamide hydrochloride (APAm) 385, 428
- N*-(3-aminopropyl)methacrylamide
hydrochloride (APMAm) 44, 45, 240, 244, 293, 383, 392, 399, 420, 428, 437, 683, 846, 850
- 3-aminopropyltriethoxysilane (APTES) 723
- 4-amino-2,2,6,6-tetramethylpiperidin-1-oxyl 724
- ammonium persulfate (APS) 508, 650, 787, 907
thermal initiation 907
redox initiation 508, 650, 787
- amphiphilic block copolymers (ABCs) 518–519, 709–711, 714, 740, 783, 851, 859, 944, 1078, 1176
- anhydride group 842–843, 878, *see also* maleic anhydride, *N*-carboxyanhydrides
- anionic polymerization 438, 709, 755
RAFT mechanism transformation (RAFT-*T*-anionic) 438
- anthracen-9-ylmethanol 780
- anti-Markovnikov orientation 775
- antimicrobial polymers 629–633, 859, 1064–1065, 1109
against *acinetobacter baumannii* 631–632
against *escherichia coli* 631–632
against *mycobacterium smegmatis* 631, 632
against *pseudomonas aeruginosa* 631–632
against *staphylococcus aureus* 631, 632
- aqueous PISA 509, 696
- aqueous RAFT (aRAFT) polymerization
deoxygenation methods 690–696
grafting from biomolecules 699–701
photoinitiated 685–690
polymerization induced self-assembly (PISA) 696–699
RAFT agents 679–684
thermally initiated (azo-compound initiators) 684–685
- 4 arm PEG-dibenzocyclooctyne (tetra-DBCO-PEG) 634
- 3-arm star polymers 191, 200, 205, 631, 785
- aromatic dithioesters 37, 223, 227, 230, 233–238, 241, 280, 313–314, 325, *see also* benzyl benzodithioate, benzyl pyridine-2-carbodithioate, cyanomethyl benzodithioate, 4-cyano-4-((phenylcarbonothioyl)thio)pentanoic acid, 2-cyanopropan-2-yl benzodithioate, 2-cyanopropan-2-yl naphthalene-1-carbodithioate, dithioesters, 1-phenylethyl benzodithioate, 2-phenylpropan-2-yl benzodithioate
- S*-aroylthiohydroxylamine 846
- aryl(pyridin-4-yl)carbamodithioates (*N*-aryl-*N*-(pyridin-4-yl)dithiocarbamates 567, 583
monomers 567
- ascorbic acid (AsAc) 43, 48, 508, 514, 620–621, 624–625, 652, 689–691, 695, 701, 713
- atom transfer radical polymerization (ATRP) 61, 118, 176, 178, 297, 308–309, 312–313, 421, 438, 502, 554, 579, 588, 617, 656–660, 708, 755, 815, 905, 983, 1160
- ATRP inimers 295
- concurrent ATRP/RAFT 308, 657
- RAFT mechanism transformation (RAFT-*T*-ATRP) 308–309, 438

- azide group
- azide-functional monomer 252, 253, 426, 649, 847, *see also* *N*-(3-azidopropyl)methacrylamide
 - azide-functional RAFT agent 408, 412, 517, 522, 757–759
 - bis-azide RAFT agent (bis- N_3 -TTC) 634
 - carbonyl azide-functional RAFT agent 409
 - azide installation post polymerization 412, 515, 839, 847, 1009
 - azide reduction to amine 835
 - azide-thioacid reaction 835
 - copper-catalyzed azide–alkyne cycloaddition 288, 502, 506, 726, 758–759, 847, 857–858, 991, 1009
 - RAFT agent synthesis 502, 506
 - Star polymers synthesis 991, 1009
 - strain-promoted azide–alkyne cycloaddition 848
- N*-(3-azidopropyl)methacrylamide (AzMA) 847
- azlactone group 46, 406, 830, 842, 859, 863
- functional monomer, *see* 2-vinyl-4,4-dimethylazlactone 46, 777, 824, 842
 - functional RAFT agent 406
- azo-compound initiators (dialkyldiazenes)
- 2,2'-azobis(*N*-(2-carboxyethyl)-2-methylpropionamide) 684
 - 4,4'-azobis(4-cyanopentanoic acid) (ACPA) 32, 36, 37, 42, 45, 320, 579, 589, 684
 - 1,1'-azobis(1-cyclohexanenitrile) (ACHN) 29, 39, 40, 46, 49, 52, 320, 321
 - 4,4'-azobis[2-(imidazolin-2-yl)propane] dihydrochloride (VA-044) 685, 991, 1002
 - 2,2'-azobis(4-methoxy-2,4-dimethylpentanenitrile) 684
 - 2,2'-azobis(2-methylbutyronitrile) (AMBN) 579
 - 2,2'-azobis[2-methyl-*N*-(2-hydroxyethyl)propionamide] 684
 - 2,2'-azobis(2-methylpropionitrile) [azobisisobutyronitrile, (AIBN)] 29, 32, 40–43, 45–49, 51–53, 99, 105, 280, 313, 318, 320–322, 324, 579, 589, 590, 647, 991, 1173, 1175–1176
 - dimethyl 2,2'-azobis(2-methylpropionate) [azobis(methyl isobutyrate, AIBMe) 589
 - azole-*N*-carboxamide methacrylates 841
 - 3,5-dimethylpyrazole-*N*-carboxamide methacrylate 841
 - imidazole-*N*-carboxamide methacrylate 841, 842
 - 1,2,4-triazole-*N*-carboxamide methacrylate 841
- b**
- BAB triblock copolymers
- using R-connected bis-RAFT agents 235–236, 260–262, 267, 936
- balanced activity RAFT agents 551, 561–564, 582–583, 593, *see also* 1*H*-pyrazole-1-carbodithioates
- barbiturate-linked poly(*n*-butyl acrylate) 784, 785
- Barton–McCombie reaction 765
- base catalyzed thiol-Michael reaction 779
- benzil dimethyl ketal (Irgacure 651) 779
- benzodithioates, *see* dithiobenzoates, dithioesters
- 5,6-benzo-2-methylene-1,3-dioxepane (BMDO) 243, 244, 297, 302, 396, 404, 431, 505, 515–516, 725
- ring-opening copolymerization-induced self-assembly (rROPISA) 725
- benzothioic *S*-acid 788–789

- benzoyl peroxide (BPO, dibenzoyl peroxide) 32, 443, 507, 648–649, 670–671, 765, 911, 914
- benzoyl peroxide–amine redox system 507, 648–649
- benzoyl peroxide–*N,N*-dimethylaniline redox system 648–649
- in eRAFT 670–671
- in RAFT end-group removal 443–446, 765
- benzoyloxy radicals 648–649
- benzyl benzodithioate (benzyl dithiobenzoate) 27, 248
- application scope 27, 248
- experimental procedure
- styrene polymerization in emulsion 36
- industrial use 30
- benzyl (diethoxyphosphoryl)methane-dithioate (DEPMD) 769–770
- benzyl dithioacetate, *see* benzyl ethanedithioate
- benzyl ethanedithioate (benzyl dithioacetate) 104, 266
- application scope 266
- in emulsion polymerization 104, 314
- O*-benzylhydroxylamine 846
- benzyl (2-hydroxyethyl) carbonotrithioate [2-(benzylsulfanylthiocarbonylsulfanyl)ethanol (BTSE)] 404, 832
- ROP-*T*-RAFT initiator 404
- benzyl mercaptan 760, 761
- benzyl methacrylate
- experimental procedures
- benzyl methacrylate in solution 41
- poly(benzyl methacrylate)-*block*-poly(*N*-(2-hydroxypropyl) methacrylamide) in solution 53
- benzyl pyridine-2-carbodithioate (BPCD) 249, 769–771
- application scope 249
- benzyl 1*H*-pyrrole-1-carbodithioate 48, 559, 894
- application scope 559
- crosslinking polymerization 894
- experimental procedure
- N*-vinylphthalimide solution
- polymerization 48
- 3-(((benzylthio)carbonothioyl)thio)propanoic acid (BCPA) [3-(benzylsulfanylthiocarbonylsulfanyl)propionic acid] 195, 400, 832, 907–908
- bioapplications of RAFT 4–5, 412, 420, 624–625, 629–635, 699–701, 850–859, 1027–1030, 1059–1065, 1103–1106, 1108–1110, 1112–1114, 1117, 1119, 1121–1122, 1125–1132
- antimicrobial polymers 629–633, 859, 1064–1065, 1109
- bioconjugate synthesis 412, 527–528, 624–625, 634–635, 699–701, 855–859, 1027–1030, 1104, 1106, 1109, 1110, 1113, 1117, 1121, 1125, 1126, 1128, 1130
- cosmetic applications 1114, 1119
- drug delivery 851–855, 1104, 1117, 1125–1130
- high throughput library synthesis 629–633, 1059–1065
- ophthalmic applications (contact lens) 1104, 1105, 1117, 1131
- patents 1103–1106, 1108–1110, 1112–1114, 1117, 1119, 1121–1122, 1125–1132
- therapeutic applications 851–855, 1059–1064, 1105, 1110, 1125–1130
- bioconjugation 412, 527–528, 624–625, 634–635, 699–701, 855–859, 1027–1030, 1104, 1106, 1109, 1110, 1113, 1117, 1121, 1125, 1126, 1128, 1130

- 2,2'-bipyridine (bpy), *see* copper activators
- biotin 691, 789, 790
- bis(2-acrylamido)ethyl disulfide (disulfide diacrylamide, DSDAm) 242, 303, 432
- N,N'*-bis(acryloyl)cystamine 400, 432, 882, 996
- bis(2-acryloyloxy)ethyl disulfide (disulfide diacrylate, DSDA) 303, 393, 408, 432
- bis-azide RAFT agent (bis-N₃-TTC) 634
- (3,5-bis((6-butylamidopyridin-2-yl) carbamoyl)phenoxy)methyl 2-bromopropanoate 785
- S,S'*-bis(α,α' -dimethyl- α' -acetic acid)-trithiocarbonate), *see* 2,2'-(thiocarbonylbis(sulfanediy))bis(2-methylpropanoic acid)
- bis-maleimide linker 780, 845, 947
- bis-maleimido(triethylene glycol) 780
- bis(2-methacryloyloxy)ethyl disulfide (disulfide dimethacrylate, DSDMA) 244, 297, 303, 432, 718, 908–909
- bis-RAFT agents 235–237, 260–263, 382, 387, 388, 718, 879, 895–896, 936, 1007
- bis-aliphatic dithioesters 237, 267
- bis-aromatic dithioesters 235–236, 260–263, 718
- bis-trithiocarbonates 382, 387, 388, 936
- disulfide functional 718
- R-connected 235–236, 260–262, 267, 879, 895–896, 936
- triblock copolymer synthesis, *see* ABA triblocks, BAB triblocks
- Z-connected 235–236, 263, 879, 936
- bis(thioacyl) disulfides 238, 239, 308, 502, 553, 578–580
- thiuram disulfides 553, 578–580
- xanthogen disulfides 502
- block copolymers, *see also* ABA triblock copolymers, ABC triblock copolymers, BAB triblock copolymers, diblock copolymers, multiblock copolymers, quasi-block copolymers
- aliphatic dithioesters 264–266
- aromatic dithioesters 241–249
- cationic RAFT polymerization 1171–1176
- dithiocarbamates 558–565, 567–571, 582–585
- DNA conjugates 625, 634, 679
- experimental procedures 51–53
- flow chemistry 691
- functional dithioesters 250–259
- mechanistic considerations 50–51, 103–105, 933–948
- protein conjugates 701, 761
- redox initiation 649–656, 669, 671
- switchable RAFT agents 116–118, 567–571, 583–585
- trithiocarbonates 391–401
- xanthates 509–515, 518–521
- BOC protecting group, *see* *t*-butyloxy-carbonyl protecting group
- bond dissociation energy 177
- carbon-halogen in ATRP initiator 177
- carbon-oxygen in alkoxyamine 177
- carbon-sulfur in RAFT agent 177, 615
- boronic acid/diol condensation 849
- bottlebrush polymers 437, 957–961, 963–964
- branched wormlike micelles 697
- 4-bromobenzenediazonium tetrafluoroborate (BrPhN₂⁺) 670–672
- bromoethyl acrylate (BEA) 840
- N*-[2-(4-bromophenyl)ethyl]acrylamide (BPEAm) 848
- 2-bromopropanoate 783, 785–786
- brush polymers 956–964, 1020–1025, *see also* graft polymers
- bimodal molar mass 1020, 1021
- block copolymer 1020, 1023
- Janus 1020, 1022–1024
- looped 1020, 1022

- brush polymers (*contd.*)
 mixed composition (mikto-arm)
 1020
 patchy 1020, 1022
- buffer
 acetate buffer 45
 phosphate buffered saline (PBS)
 691–693, 695
- butadiene (Bd)
 blocks 264, 265, 399
 copolymers 209, 247, 249, 264, 265,
 385, 393, 399, 408
 butadiene/styrene 249, 264, 265
 acrylonitrile/butadiene 249, 264,
 385, 393, 399, 408
 homopolymer 265, 280, 393, 399
 with dithioesters 247, 249, 264, 265,
 280
 with trithiocarbonates 385, 393, 399,
 408, 421, 424
- S*-(2-((*t*-butoxycarbonyl)amino)ethyl)
 methanesulfonothioate 790
 thiol-methanethiosulfonate reaction
 790
- N*-*t*-butylacrylamide (tBAM) 736
- butyl acrylate (BA, *n*-butyl acrylate) 41,
 42, 51, 505, 657, 708, 759, 765,
 772, 773, 785, 810
- t*-butyl acrylate (tBA) 39, 43, 421, 615,
 671, 777, 780, 810, 845, 986,
 1007
- butyl (1-(((3,5-dioxo-1,2,4-triazolidin-4-yl)
 hexyl)amino)-1-oxopropan-2-yl)
 carbonotrithioate (URZ-TTC)
 410, 772
- butyl 2-(((dodecylthio)carbonothioyl)
 thio)-2-methylpropanoate (CTA
 ester) 396, 1101, 1132
 application scope 396
 industrial use 1101, 1132
- t*-butyl hydroperoxide 48, 652
t-butyl hydroperoxide–ascorbic acid
 redox system 648, 652
- butyl methacrylate (*n*-butyl methacrylate)
 38, 416, 557, 669, 857, 1056
- t*-butyloxycarbonyl (BOC) primary amine
 protecting group 129, 290, 292,
 293, 297, 767, 790
 in monomer 292, 293, 427, 838, *see*
also di(BOC)-acrylamide
 in RAFT agent 251, 407, 767
 in reagent 790, *see also S*-(2-((*t*-but-
 oxy carbonyl)amino)ethyl)
 methanesulfonothioate
 2-(((butylthio)carbonothioyl)thio)
 propanoic acid (BTPA, BuPAT)
 398, 614, 624, 628, 686, 690,
 701
 application scope 398
 aqueous RAFT (aRAFT) 686, 690,
 701
 functional derivatives
 ATRP initiator 410
t-butyldimethylsilyl 411
 poly(dimethylsiloxane) (PDMS)
 412
 protected maleimide 410
 photoinduced energy or electron
 transfer (PET) RAFT 614, 624,
 628
- C**
 carbonotrithioates, *see* trithiocarbonates
 4-(((2-carboxyethyl)thio)carbonothioyl)
 thio)-4-cyanopentanoic acid
 818–820
 application scope 27, 393
 industrial use 30
 3-(((1-carboxyethyl)thio)carbonothioyl)
 thio)propanoic acid 27, 398,
 429
 application scope 27, 398
 2-(1-carboxyl-1-methylethylsulfanylthio-
 carbonylsulfanyl)-2-methyl-
 propionic acid (CMP), *see* 2,2'-
 (thiocarbonylbis(sulfanediy))
 bis(2-methylpropanoic acid)
 2-carboxyprop-2-yl radicals 381
 catalytic chain transfer (CCT) 793, 936,
 1077–1078, 1082, 1083

- cationic RAFT polymerization
 1161–1163
 block copolymers 1173–1176
 design 1163–1172
 end-functionalized polymers
 1172–1173
 initiators 1167–1170
 monomers 1170–1172
 RAFT agents 1163–1167
 RAFT mechanism transformation
 (cationic-*T*-RAFT) 438,
 1175–1178
 star polymers 1177–1178
- chain length distribution (CLD) 66, 98,
 199, 554, 906
- chain transfer coefficient (C_{tr}) 96–103,
 226–230, 374
 apparent chain transfer coefficient
 (C_{tr}^{app})
 dithiocarbamates 554–556, 580
 dithioesters 101, 228–229
 trithiocarbonates 375–377
 xanthates 509–510
- measurement of 97–103
 dispersity method 102, 227, 230,
 375–377
 kinetic simulation 103, 228–229,
 375–377
 ln(CLD) method 97–98
 Mayo method 97–98, 100, 228–229,
 375–377, 554–556
 $M_n(0)$ method 228–229, 375–377
 Walling method 98–100, 228–229,
 375–377, 554–556
- reverse chain transfer coefficient (C_{tr})
 97, 226–230
- chain transfer rate coefficient (k_{tr}) 96,
 125, 226–230, 374
 apparent chain transfer rate coefficient
 (k_{tr}^{app})
 reverse chain transfer rate coefficient
 (k_{tr}) 97, 226–230
 measurement of 97–103
- chain transfer agent (CTA), *see* RAFT
 agent
- chemoselective H_2S -responsive
 fluorescence probe 789
- 2-chloroethyl vinyl ether (CEVE) 1171
- 4-chloromethylstyrene (4-vinylbenzyl
 chloride) 241, 247, 262, 280,
 385, 391, 393, 395, 396, 399, 400,
 408, 421, 510, 559, 560, 572, 839,
 848, 859, 1004, 1009
 with dithiocarbamates 559, 560, 572
 with dithioesters 241, 247, 262, 280
 with trithiocarbonates 385, 391, 393,
 395, 396, 399, 400, 408, 421
 with xanthates 496, 503, 510
- chloroprene 242, 245, 249, 265, 401, 424,
 1034
 SI-RAFT 1034
- chlorophyll *a* (Chl *a*) 619–620
- cholest-5-en-3 β -yl 6-iodohexyl ether 783
- classically hyperbranched polymers 967
- click chemistry approach 517, 522, 525,
 649, 755, 757, 806, 847, 944–947,
 949, 960, 991, 992
- CMP-functionalized ssDNA 701
- comb polymers 297, 425, 956
- co-mediated radical polymerization 708
- composites 1030–1039
 anisotropic structures 1031, 1039
 SI-RAFT 1030–1039
- computational chemistry 139, 152, 153,
 156, 159, 165, 167, 172, 180
- Concanavalin A* (*ConA*) 630, 857
- confocal laser scanning microscopy
 (CLSM) 1025
- continuum models 147
- controlled/living polymerization, *see*
 reversible deactivation
 polymerization
- controlled radical polymerization (CRP),
see reversible deactivation
 radical polymerization 15, 60,
 176, 614, 625–626, 630, 695–696,
 786, 861, 964, 983, 1042
- copolymerization 48, 753–755
 branched copolymerization 209
 of multifunctional monomer 966–967

- copolymers 755, *see also* block copolymers, gradient copolymers
- desulfurization of RAFT (co)polymers 763–76
- copper activators, *see also* ligands
 - in alkoxyamine synthesis
 - CuX/Me₆TREN 517
 - in alkyne-azide coupling
 - CuX/PMDETA 759, 1024
 - in concurrent RAFT/ATRP
 - CuX/ dNbpy 657
 - CuX/Me₆TREN 657
 - CuX/PMDETA 657
 - CuX/TMPA 657
 - in radical generation
 - CuX/Me₆TREN 120, 517
 - in redox RAFT
 - Cu^{II}(DC)₂/bpy 668
 - CuX/Me₆TREN 658
 - CuX/PMDETA 313, 656–658, 669
 - CuX/TMPA 658–659, 669–670
- copper catalyzed azide-alkyne cycloaddition (CuAAC) 523, 717, 847, 857, 945, 955, 1023
- copper-mediated polymerization 708
- core-crosslinked stars (CCS) 301, 433, 874, 918, 994, 996, 998, 1006–1007
- core-crosslinked nanoparticles 718, 722–725
 - degradable core 725
 - thiol-functional vesicles 718
- coumarin chromophore 286, 613, 629, 691, 807
 - acceptor in light-harvesting polymers 242, 286, 807
 - dye 613, 629
 - in photoresponsive methacrylamide 691
 - in styrenic monomer 241, 247, 286
- cross-coupling reactions 325, 591, 848–849
- crosslink density distribution 873, 876
- crosslinked polymers 873, 910
 - crosslinking monomers, *see* divinyl monomers
- cumene hydroperoxide (CHP) 43, 652
 - cumene hydroperoxide-ascorbic acid redox system 648, 652
- cumyl dithiobenzoate (CDB), *see* 2-phenylpropan-2-yl benzodithioate
- cumyl phenyldithioacetate (CPDA), *see* 2-phenylpropan-2-yl 2-phenylethanedithioate
- curve-crossing model 167, 169–170
- cianoalkyl RAFT agents 105, 108, 110, 130, 175, 233–236, 380–381
- 2-cyanobutan-2-yl 4-chloro-3,5-dimethyl-1*H*-pyrazole-1-carbodithioate 28, 30, 40
 - application scope 28, 562, 582–583
 - industrial use 30, 561, 1099
 - synthesis of 579
- 2-cyanobutan-2-yl 3,5-dimethyl-1*H*-pyrazole-1-carbodithioate 28, 30, 119, 561, 579, 582–583
 - application scope 28, 561, 582–583
 - chain transfer coefficient 583
 - industrial use 30, 561, 1099
 - synthesis of 579
- 4-cyano-4-(((dodecylthio)carbonothioyl)thio)pentanoic acid 7, 27, 31, 39, 46, 49, 131, 393, 441, 612, 614–615, 627–628, 819, 898, 902–903, 913, 1132
 - application scope 27, 393
 - deoxygenation methods 31
 - experimental procedure
 - N*-isopropylacrylamide in solution 46
 - methyl methacrylate in solution 39
 - poly[(methacrylic acid)-*co*-(oleyl methacrylate)] in solution 49
 - industrial use 30, 1100, 1102
 - PET-RAFT 131, 614–615, 627–628
 - side reactions 441

- 4-cyano-4-(((ethylthio)carbonothioyl)thio)pentanoic acid 392, 441, 681, 693, 695
 application scope 392
 side reactions 441
- 4-cyano-4-(((propylthio)carbonothioyl)thio)pentanoic acid 392, 693
 application scope 392
- 2-cyano-5-hydroxypentan-2-yl benzo-dithioate (CHPB) 250 757
- cyanoisopropyl decyl trithiocarbonate, *see* 2-cyanopropan-2-yl decyl carbonotrithioate
- cyanoisopropyl dithiobenzoate, *see* 2-cyanopropan-2-yl benzodithioate
- cyanoisopropyl 1-dithionaphthalate, *see* 2-cyanopropan-2-yl naphthalene-1-carbodithioate
- cyanoisopropyl *N,N*-diethyldithiocarbamate, *see* 2-cyanopropan-2-yl diethylcarbamidithioate
- cyanoisopropyl dodecyl trithiocarbonate, *see* 2-cyanopropan-2-yl dodecyl carbonotrithioate
- cyanoisopropyl methyl trithiocarbonate, *see* 2-cyanopropan-2-yl methyl carbonotrithioate
- cyanoisopropyl radicals, *see* 2-cyanopropan-2-yl radicals (NIPAM)
- cyanomethyl benzodithioate 249
 polymerizations 249
- cyanomethyl 3,5-dimethyl-1*H*-pyrazole-1-carbodithioate 9, 27, 30, 40, 46, 49, 52, 119, 583
 application scope 27, 582–583
 chain transfer coefficient 583
 experimental procedures
N,N-dimethylacrylamide in solution 46
 methyl acrylate in solution 40
 poly[(methyl acrylate)-*co-N,N*-dimethylacrylamide] in solution 49
 poly(*N,N*-dimethylacrylamide)-*block*-poly(vinyl acetate) in solution 52
 industrial use 30
 polymerizations 40, 46, 49, 52, 119, 562
- cyanomethyl dodecyl carbonotrithioate 400, 429
 polymerizations 400
- cyanomethyl methyl(phenyl)carbamidithioate (*N*-methyl-*N*-phenyldithiocarbamate) 28, 30, 149, 179
 application scope 28
 industrial use 29–30
 polymerizations 565
- cyanomethyl methyl(pyridin-4-yl)carbamidithioate 30, 47, 569–570
 polymerizations 47, 569–570
 experimental procedure
N-vinylcarbazole in solution 47
 synthesis 580
- 4-cyanopentanoic acid dithiobenzoate, *see* 4-cyano-4-((phenylcarbonothioyl)thio)pentanoic acid
- 4-cyano-4-((phenylcarbonothioyl)thio)pentanoic acid (4-cyanopentanoic acid dithiobenzoate) 35, 45, 228, 243–244, 307, 312, 322, 614, 627, 628, 662, 671, 679–681, 683, 701, 757, 832, 838, 915, 1102
- chain transfer coefficient 228
- eRAFT 662–664, 667, 672
- experimental procedures
N-(2-hydroxylpropyl)methacrylamide in pH 5.2 buffer 45
N-[3-aminopropyl] methacrylamide in pH 5 buffer 45
- functional derivatives 251–252
 active ester 252
 azide 253
 dibromomaleimide 255
 ene 255
 hydroxyl 250
 NHS ester 251, 831

- 4-cyano-4-((phenylcarbonothioyl)thio)pentanoic acid (4-cyanopentanoic acid dithiobenzoate) (*contd.*)
- norbornene 256
 - protected maleimide 256–257
 - poly(dimethylsiloxane) (PDMS) 257–258
 - poly(ethylene glycol) (PEG) 258
 - thiolactone 256
 - yne 253–254
 - industrial use 1100, 1102
 - PET-RAFT 307, 312
 - polymerizations 45, 243–244
 - SUMI 307
- 2-cyanopropan-2-yl radicals
- (cyanoisopropyl radicals) 107, 443, 766–767, 810
 - rate coefficient for monomer addition 107
- 2-cyanopropan-2-yl benzodithioate (cyanoisopropyl dithiobenzoate)
- 2, 27, 30, 33, 40–41, 46, 49, 51, 67–69, 108, 109, 111, 122, 124, 149, 151, 161, 164–166, 211, 242, 662, 754, 779, 909–810, 898
 - application scope 27, 242
 - chain transfer coefficient 228, 230
 - eRAFT 662
 - experimental procedures
 - butyl acrylate polymerization in solution 41
 - methyl methacrylate polymerization in solution 40–41
 - poly(methyl methacrylate-*co*-trimethylsilyl methacrylate-*co*-*t*-butyl dimethylsilyloxystyrene) in solution 49
 - industrial use 30, 1100, 1102
 - initialization 67–69, 151
 - RAFT equilibrium coefficient 124, 232
- 2-cyanopropan-2-yl diethylcarbami-dithioate (cyanoisopropyl *N,N*-diethyl-dithiocarbamate)
- 565, 668
 - application scope 565
 - eRAFT 668
- 2-cyanopropan-2-yl decyl carbonotrithioate (cyanoisopropyl decyl trithiocarbonate) 810
- 2-cyanopropan-2-yl dodecyl carbonotrithioate (cyanoisopropyl dodecyl trithiocarbonate) 30, 165, 198, 201, 376–379, 391–392, 429, 432, 810, 813, 908, 913, 1100, 1102
- application scope 380–381
 - chain transfer coefficient 376–377
 - industrial use 1100, 1102
 - RAFT equilibrium coefficient 378
- 2-cyanopropan-2-yl methyl carbonotrithioate (cyanoisopropyl methyl trithiocarbonate) 391, 764
- application scope 391
- 2-cyanopropan-2-yl naphthalene-1-carbodithioate (cyanoisopropyl 1-dithionaphthalate) 211, 245, 616
- application scope 245
 - mathematical modeling 211
- 2-cyanopropan-2-yl 1*H*-pyrrole-1-carbodithioate 194, 559, 579
- application scope 559
 - mathematical modeling 194, 208
 - synthesis of 579
- cyclic ketene acetals 512, 515–516, 533, 725, *see also* 5,6-benzo-2-methylene-1,3-dioxepane (BMDO), 2-methylene-4-phenyl-1,3-dioxolane
- cyclic polymers 5, 517, 521–522, 761, 780, 785, 787, 949–950
- polystyrene 761, 781
- cyclic *N*-vinylamides, *see* *B*-vinylcaprolactam, *N*-vinyl-2-piperidone, *N*-vinylpyrrolidone
- [4+2] cycloaddition reaction 769

- β -cyclodextrin (β -CD) 719, 731, 737,
950–951, 985, 991, 993,
1009
- cyclohexylamine 773–774
- cyclopentadiene 769
- cyclopolymerization 5, 297, 300,
429–431, 516
- cysteine (Cys) 315, 318, 700, 720, 789,
1027
- d**
- degassing, *see* deoxygenation
- density functional theory (DFT)
140–142, 144, 157, 174, 514, 521,
660, 664
- deoxygenation in RAFT polymerization
(degassing) 31, 34, 40–42,
46–53, 442, 611, 619, 654–656,
690–696, 1052–1053, 1058,
1067
- enzyme-catalyzed deoxygenation
654–656, 691–696, 1057–1058
with enzyme-mediated initiation
654–656, 693–696
with thermal initiation 691–693
- experimental procedures 34, 40–42,
46–53
- in high throughput experiments 630,
1052–1053, 1059, 1067
- kinetics of MMA polymerization 31
- in PET-RAFT 442, 611, 619, 630,
690–691, 1058
- deoxygenation in RAFT end-group
removal 773
- desulfurization of RAFT polymers
addition-fragmentation coupling
766–768
radical-induced oxidation 768–769
radical-mediated reduction 765–766
thermolysis 763–764
- DET-RAFT polymerization 660
- diacetone acrylamide (DAAm, *N*-(1,1-
dimethyl-3-oxobutyl)acrylamide)
44, 398, 421, 716, 719, 724, 730,
734
- dialkyldiazene initiators, *see*
azo-compound initiators
- N,N*-dialkyldithiocarbamates 114, 116,
552, 553, 581, 586, 588, 590
- p*-dialkylsulfonium phenoxy ester
(DASPE) 837–838
- 4-acryloxyphenyldimethylsulfonium
triflate 837
- dibenzoyl peroxide, *see* benzoyl peroxide
(BPO)
- dibenzyl carbonotrithioate (DBTTC,
dibenzyl trithiocarbonate) 27,
42, 111, 376–377, 385, 431, 886,
1101, 1132
- application scope 27, 385
- chain transfer coefficients 376–377
- experimental procedures
3,5-dibromobenzyl acrylate in
solution 42
2-ethylhexyl acrylate in solution
42
octadecyl acrylate in solution 42
- industrial use 27, 1101, 1132
Blocbuilder DBTM 1101
- RAFT crosslinking polymerization
886
- RAFT equilibrium coefficients 378
- dibenzyl trithiocarbonate, *see* dibenzyl
carbonotrithioate
- diblock copolymer 512, 515, 518–519,
521, 679, 696–699, 709, 714, 718,
720, 727–729, 732, 734–735, 737,
740, 770, 787, 807, 857, 936, 938,
940, 945, 955, 968, 1042,
1056–1058, 1175
- mechanistic consideration 103–105
- di(BOC)-acrylamide (DBAm) 838
- 3,5-dibromobenzyl acrylate 42
- dibromotrithiocarbonate [DiBrTTTCm,
bis(5-bromo-5-methyl-4-
oxohexyl) 2,2'-(thiocarbonylbis
(sulfanediy))bis(2-methylpro-
panoate)]
- ATRP initiator 657

- dibromotrithiocarbonate [DiBrTTCm, bis(5-bromo-5-methyl-4-oxohexyl) 2,2'-(thiocarbonylbis(sulfanediyl))bis(2-methylpropanoate)] (*contd.*)
- N,N'*-dicyclohexylcarbodiimide (DCC) 37, 238, 240, 878
- Diels-Alder [2+4] cycloaddition 845
- diene monomers, *see also* butadiene, chloroprene, isoprene
- with dithioesters 280
- with trithiocarbonates 424
- (diethoxyphosphoryl)methanedithioates 769, 770, 1089, 1090, 1110, *see also* benzyl (diethoxyphosphoryl)methanedithioate
- N,N*-diethylacrylamide (DEAm) 46, 696, 781–783, 831, 832, 1058
- experimental procedures
- N,N*-diethylacrylamide in solution 46
- with dithioesters 242, 245, 248, 249, 264, 280, 317
- with trithiocarbonates 395, 398, 399, 400, 409, 421
- 4-(diethylamino)salicylaldehyde (DEAS) 649
- diethylene glycol acrylate (DEGDA) 394, 400
- diethylene glycol dimethacrylate (DEGDMA) 301, 392, 393, 432, 882, 883
- (diethylene glycol monomethyl ether) acrylate (DEGA) 761–762
- (diethylene glycol monomethyl ether) methacrylate (DEGMA) 761–762
- α,ω -difunctional polymer 789
- difunctional CTAs, *see* bis-RAFT agents 936, 948
- 2,3-dihydroxypropyl methacrylate, *see* glycerol monomethacrylate
- 2-(diisopropylamino)ethyl methacrylate (DPAEMA) 683
- N,N*-dimethylacrylamide (DMAm) 46, 49, 52, 117, 118, 280, 301, 307, 421, 511, 518, 521, 522, 572, 583, 587, 590, 618, 652, 654–656, 684, 686, 689–691, 693–696, 701, 772, 780, 791, 792, 818, 820, 849, 903, 1027, 1053, 1057, 1063
- experimental procedures
- N,N*-dimethylacrylamide in solution 46
- poly(*N,N*-dimethylacrylamide)-*block*-polystyrene by surfactant-free emulsion polymerization 36–37
- propagating radical 766
- selective initialization 108
- single unit monomer insertion (SUMI) 108
- with dithioesters 234, 241, 242, 244, 245, 248, 249, 250, 253, 259, 262, 264, 280, 301, 307, 317
- with trithiocarbonates 107, 818
- 2-(dimethylamino)ethyl methacrylate (DMAEMA) 240, 416, 557, 615, 632, 696, 698, 720, 725, 857, 1003, 1060, 1064
- block 857
- 4-dimethylaminopyridine (DMAP) 37, 878
- dimethyl(phenyl)phosphine 779
- N*-(1,1-dimethyl-3-oxobutyl)acrylamide, *see* diacetone acrylamide
- 4,4'-dinonyl-2,2'-bipyridine (dN bpy), *see* copper activators
- 2,5-dioxopyrrolidin-1-yl 4-vinylbenzoate (4-vinylbenzoic acid, NHS ester) 243, 251, 290, 426, 831
- diphenylacetylene 661
- α,ω -dipyrenyl poly(*N*-isopropylacrylamide) 783
- 2,2'-dipyridyl disulfide 788–789, 968
- discrete oligomers 813, 818, 823
- oligo(methyl acrylate) 823
- diselenocarbonyl compounds 786

- dispersion polymerization 38, 187, 195, 211, 440–442, 650, 686, 691, 696–699, 709–713, 718–721, 723–726, 729–731, 733–736, 898, 900–901, 996, 998–999, 1006
- dispersity 28, 33, 37–38, 88, 102–103, 123–124, 206, 227, 230, 375, 429, 513, 515, 532, 626, 654, 805–806, 823, 831, 834–839, 953, 967, 1041
- dissociative electron transfer (DET)-RAFT 179, 312, 439, 660
- disulfide group
- bis-RAFT agents
 - bis-dithiobenzoate (DSDB) 718
 - monomers, *see* bis(2-methacryloyl)oxyethyl disulfide
 - disulfide diacrylamide (DSDAm), *see* bis(2-acrylamido)ethyl disulfide
 - disulfide diacrylate (DSDA), *see* bis(2-acryloyloxy)ethyl disulfide
 - disulfide dimethacrylate (DSDMA), *see* bis(2-methacryloyloxy)ethyl disulfide
- RAFT agents 406
- RAFT inimers 301, 308
- reactions 787–790
- reagents, *see* S-(2-((t-butoxycarbonyl)amino)ethyl) methanesulfonothioate, bis(thioacyl) disulfide, 2,2'-dipyridyl disulfide
- dithiobenzoates (benzodithioates) 59, 76–89, 187, 223, 659, 661, 662, 753, 759, 764, 767, 781, 808, 830, 935, *see also* dithioesters
- kinetic analysis 85–87
- macroRAFT agent 807, 808
- MMA model system 88–89
- quantum-chemical calculations 87–88
- retardation mechanisms
- intermediate radical termination 77, 119–123
 - missing step reaction 77–86, 88–90, 121
 - slow fragmentation 59, 76, 77, 85, 86, 123–126
- dithiocarbamates 112–114, 662, 668, 753, 759, 830, 936, 1165, *see also* N,N-dialkyldithiocarbamates, aryl(pyridin-4-yl)carbamo-dithioates, 1H-imidazole-1-carbodithioates, 1H-pyrazole-1-carbodithioates, cyanomethyl methyl(phenyl)carbamo-dithioate, methyl(pyridin-4-yl)carbamo-dithioates, 1H-pyrrole-1-carbo-dithioates
- activity of 580–586
- application scope 556–571
- balanced activity 551, 561–564, 582–583, 593, *see also* 1H-pyrazole-1-carbodithioates
- canonical forms 114, 582
- cationic polymerization 585
- chain transfer coefficients 555–556, 580–586
- emulsion polymerization 587
- dithiocarbamate group removal/ transformation 588–591
- by copper-promoted cross-coupling 591
 - by oxidation 591
 - by radical-induced coupling 588
 - by radical-induced disproportionation 588–589
 - by radical-induced reduction 589
 - by reaction with nucleophiles 589–590
 - by thermolysis 590–591
- monomers 558–570, 567–571
- 1,1-disubstituted MAMs 557
 - methacrylates 557
 - LAMs, IAMs 552, 576
 - vinyl monomers 552, 576
- monosubstituted MAMs 572–575
- acrylamides 572, 573
 - acrylates 572, 574
 - styrenes 572, 575

- dithiocarbamates (*contd.*)
- RAFT mechanism transformation
 - 587–588
 - RAFT-*T*-ATRP 588
 - RAFT-*T*-ROMP 587–588
 - ROP-*T*-RAFT 587
 - ROMP-*T*-RAFT 587–588
 - selection of 554, 575, 582, 593
 - R substituent 585
 - switchable dithiocarbamates 30, 47, 51–52, 116–118, 583, *see also* methyl(pyridin-4-yl)carbamo-dithioates
 - synthesis of dithiocarbamates 575–580
- dithiocarbonates, *see* xanthates
- dithioesters (dithioates) 223–270, 753, 759, *see also* benzyl benzodithioate, benzyl (diethoxyphosphoryl)methanedithioate, benzyl pyridine-2-carbodithioate, cyanomethyl benzodithioate, 4-cyano-4-((phenylcarbonothioyl)thio)pentanoic acid, 2-cyano-propan-2-yl benzodithioate, 2-cyanopropan-2-yl naphthalene-1-carbodithioate, dithio-benzoates, 2-((phenylcarbonothioyl)thio)acetic acid, 1-phenylethyl benzodithioate, 2-phenylpropan-2-yl benzodithioate, 2-phenylpropan-2-yl benzodithioate, 2-phenylethanedithioate
- application scope 234
- aliphatic dithioesters 236–237, 264–267
 - aromatic dithioesters 235–236, 241–249, 250–263
 - bis-aliphatic dithioesters 237, 267
- atom-transfer radical polymerization 308–309
- bis-aromatic dithioesters 235, 260–263
 - crosslinking polymerization 301, 303–304
 - model co-networks 304
- cyclopolymerization 297, 300
- dithioester group
- removal/transformation 314–325
 - boronic acid cross-coupling 325
 - electrocyclic reaction 322–325
 - oxidation 321
 - radical-induced coupling 318
 - radical-induced reactions 315
 - radical-induced reduction 318
 - reaction with nucleophiles 315
 - thermolysis 322
- α,ω -dithiol functional polystyrene 787
- emulsion/mini-emulsion
- polymerization 313–314
- industrial use 1101–1102, 1132
- initiation, *see also* redox initiation
- photoinitiation 311–312
 - thermal initiation 310–311
- inimers 301, 305–306
- chain transfer coefficients 226–230
 - equilibrium coefficients 230–232
 - mechanism 224–232
- monomers
- 1,1-disubstituted MAMs 240
 - methacrylamides 240, 278–279
 - methacrylates 239–240, 268–277
 - 1,2-disubstituted MAMs 280
 - macromonomers 297–299
 - monosubstituted LAMs and IAMs 287–288
 - monosubstituted MAMs
 - acrylamides 280, 284–285
 - acrylates 240, 277, 281
 - acrylonitrile 112, 234, 242, 244, 245, 247
 - diene 280
 - styrenics 280, 286–287
 - dithioesters with reactive functionality 289–297
- publication statistics 224, 233, 236
- RAFT agent selection 230–237
- RAFT mechanism transformation 307–308
- cationic-*T*-RAFT 1175–1178

- nitroxide-mediated polymerization (NMP) 309–310
 - ring-opening polymerization (ROP) 308
 - ring-opening metathesis polymerization (ROMP) 308
 - RAFT-SUMI 306–307
 - ring-opening polymerization 297, 301, 302
 - self-condensing vinyl polymerization 301, 305–306
 - side reactions 126–130, 313
 - with reactive functionality 288–297
 - synthesis of 237–239
 - 1,4-dithiothreitol 316–317, 724
 - divinyl monomers, (crosslinking monomers) 301, 303, 431–433, 906–907, 910, 966, *see also*
 - bis(2-acrylamido)ethyl disulfide (DSDAm), *N,N'*-bis(acryloyl)-cystamine, bis(2-acryloyloxy)ethyl disulfide (DSDA), bis-maleimidotriethylene glycol, bis(2-methacryloyloxy)ethyl disulfide (DSDMA), ethylene glycol dimethacrylate (EGDMA), diethylene glycol diacrylate (DEGDA), diethylene glycol dimethacrylate (DEGDMA), divinylbenzene, methylene-bis-acrylamide (MBAm), triethylene glycol dimethacrylate (TEGDMA)
 - crosslinking monomers 301, 303, 431–433
 - crosslinking polymerization
 - polymer networks 301, 431–433, 873–915
 - star microgels 301, 431–433
 - divinylbenzene (DVB) 241, 249, 258, 261, 301, 391, 392, 394, 395, 399, 400, 404, 405, 432, 709, 723, 885, 887, 892, 900, 906, 911, 913–915, 918, 967
 - with dithioesters 241, 249, 258, 261, 301
 - PISA 709, 723
 - with trithiocarbonates 391, 392, 394, 395, 399, 400, 404, 405, 432
 - 1-dodecanthiol (DDT) 760, 791, 966
 - 2-(dodecylthiocarbonothioylthio)-2-methylpropionic acid (DDMAT) 37, 395–396, 420, 424, 614–615, 662–664, 668–672, 834, 837, 898, 1101, 1132
 - application scope 395–396
 - experimental procedures
 - acrylic acid in solution 41
 - t*-butyl acrylate in solution 41
 - styrene surfactant-free in emulsion 37
 - industrial use 1101, 1132
 - maleimido ethoxyethyl ester 409, 700
 - photoinitiated RAFT 614–615
 - redox initiated RAFT 662–664, 668–672
 - 2-(((dodecylthio)carbonothioyl)thio)propanoic acid (DoPAT) 27, 30, 209, 399, 1034, 1035
 - application scope 27, 399
 - industrial use 30, 1101, 1132
 - SI-RAFT 1034, 1035
 - dodecyl vinyl ether 775
 - doxorubicin (DOX) 853, 855, 1004, 1029, 1041
 - drug delivery 4, 718, 738–739, 851–855, 1003, 1004, 1029, 1056, 1059–1064
 - dual concurrent electrochemically mediated ATRP/RAFT 669
 - dual-responsive PNIPAM-*b*-PAA copolymer 855
 - dynamic polymeric nanoobjects 1041
- e**
- electrochemically-initiated RAFT polymerization (eRAFT) 312, 439, 660–673

- electrochemically-initiated RAFT polymerization (eRAFT) (*contd.*)
 - ATRP initiation 668–670
 - direct eRAFT 665–667
 - initiator reduction 670–673
 - benzoyl peroxide 670–671
 - diazonium salt 670–672
 - monomers
 - butyl acrylate 666, 667
 - butyl methacrylate 666, 669
 - ferrocenyl methacrylate 672
 - methyl methacrylate 666, 669–671
 - styrene 666, 667, 669
 - organic mediators 667–668
 - RAFT agents 661–665
 - 4-cyano-4-((phenylcarbonothioyl)thio)pentanoic acid 662–664, 667, 672
 - 2-cyanopropan-2-yl benzodithioate (cyanoisopropyl dithiobenzoate) 662
 - 2-cyanopropan-2-yl diethylcarbamo-dithioate (cyanoisopropyl N,N-diethyldithiocarbamate) 668
 - dithiobenzoates 312, 661–665, 667, 672
 - reduction potential 661
 - trithiocarbonates 439
 - surface initiated (SI-eRAFT) 672
 - electron affinity (EA) 176
 - electron paramagnetic resonance (EPR) 59–65, 70–76, 81, 86–87, 89–91, 103, 124, 149, 204, 379
 - emulsion polymerization 34–37, 42, 195, 313–314, 413–414, 587, 710, 1078–1080, 1092, 1095–1097, 1100, 1105, 1108–1112, 1114, 1121, 1125, 1129, 1132, *see also* polymerization-induced self-assembly (PISA)
 - amphiphilic macroRAFT agents 36–37, 42, 48, 1080, 1096, 1100, 1111
 - with dithiocarbamates 587
 - with dithioesters 313–314
 - experimental procedures
 - butyl acrylate 42
 - styrene 36–37
 - with macromonomer RAFT agents 1078–1079
 - mathematical modeling
 - deterministic modelling 195, 209
 - stochastic modelling 205–206
 - patents 1078–1080, 1092, 1095–1097, 1100, 1105, 1108–1112, 1114, 1121, 1125, 1129, 1132
 - sulfur-free emulsion polymerization 1078–1079
 - surfactant-free emulsion
 - polymerization 36–37, 42, 48, 1080, 1096, 1100, 1111
 - with trithiocarbonates 413–414
 - with xanthates 507–509, 512, 515
 - enaminones 846
 - end-functionalized polymers 807, 1172–1173
 - α -end-functionalized poly(*n*-butyl acrylate) 772–773
 - α,ω -end-functionalized poly(*N*-isopropyl-acrylamide) 761
 - end-group fidelity, *see* livingness
 - end-group removal, *see* RAFT end-group removal
 - Energy-Directed Tree Search 145, 152
 - enhanced green fluorescent protein (EGFP) 855, 857
 - enzymatic initiation systems 654–656, 693–696
 - glucose oxidase (GOx) 654, 696
 - horseradish peroxidase (HRP) 693–695
 - enzyme catalyzed deoxygenation
 - enzymatic initiation systems 693–696
 - initiation by thermal initiation 691–693
 - enzyme-responsive inimer 966

- Eosin Y (EY) 312, 612, 623, 624, 630, 635, 689, 690, 1027
- epoxide group 783, 843–844
- epoxy-functional monomer, *see* glycidyl methacrylate
- eRAFT, *see* electrochemically-initiated RAFT polymerization
- ester-end-functionalized polymer 783
- 2-ethanolamine 779
- 2-((ethoxycarbonothioyl)thio)acetic acid (ethylxanthogenacetic acid) 48, 497, 512, 516
 application scope 497
 experimental procedure
 vinyl acetate polymerization in solution 48
- 1-(ethoxycarbonyl)prop-1-yl dithiobenzoate (EPDTB) 836
- 1-ethoxyethyl acrylate 759–760
- 2-ethoxyethyl vinyl ether (EOEVE) 1171, 1175
- ethyl α -bromoisobutyrate (EBiB, ethyl 2-bromo-2-methylpropanoate) 659, 669, 670
- ethylene glycol diacrylate (EGDA) 301, 496, 503, 996, 999
- ethylene glycol dimethacrylate (EGDMA) 198, 241, 242, 261, 266, 249, 301, 304, 392, 395, 399, 407, 432, 718, 899, 900, 901, 906, 908, 911–913, 915, 966
- 2-ethylhexyl acrylate (EHA) 39, 42, 421, 810
- N*-ethylpiperidine hypophosphite (EPHP) 445, 531, 765, 766
- S*-ethyl-*S'*-(α, α' -dimethyl- α'' -acetic acid) trithiocarbonate, *see* 2-(((ethylthio)carbonothioyl)thio)-2-methylpropanoic acid
- O*-ethyl-*S*-(1-methoxycarbonylethyl) dithiocarbonate, *see* methyl 2-((ethoxycarbonothioyl)thio)propanoate
- ethyl 2-((phenoxycarbonothioyl)thio)propanoate 616
- ethyl 2-(((propylthio)carbonothioyl)thio)propanoate (*S'*-(1-ethoxycarbonylethyl) *S*-propyl trithiocarbonate) 72–76, 378, 379, 399
- RAFT equilibrium coefficients 72–76, 378
- 2-(((ethylthio)carbonothioyl)thio)-2-methylpropanoic acid 394, 681, 685, 886
- aqueous RAFT 681, 685
- crosslinking polymerization 886
- functional derivatives
 active ester 404
 protected thiol 406
 protected maleimide 409
- polymerizations 394
- (5-ethyl-2,4,6-trioxohexahydropyrimidin-5-yl)methyl 2-bromopropanoate 786
- ethylxanthogenacetic acid, *see* 2-((ethoxycarbonothioyl)thio)acetic acid
- f**
- fast polymerizations 624, 654, 805, 1027, 1168
- ferrocenyl methacrylate (FcMMA) 672
- flash chromatography 813, 820
- flat surfaces 1017, 1018, 1023, 1025
- Flory interaction parameter 898, 900
- Flory polymer solvent interaction parameter 875
- fluorescein-based semicarbazide 724
- fluoroalkenes 1176
- fluoroparticles 720, 722–723
- fraction of living chains, *see* livingness
- folic acid tethered polymer-modified magnetic nanoparticles (FA-MNPs) 855
- 2-(4-formylbenzoyloxy)ethyl methacrylate (FBEMA) 846

- 3-formyl-4-hydroxybenzyl methacrylate (FHMA) 725
- fragmentation rate coefficient, *see* rate coefficients
- α,ω -functionalized poly(*N*-isopropylacrylamide) 760
- ω -functionalized poly(*N*-isopropylacrylamide) 760, 777
- α,ω -functionalized poly(diethylene glycol monomethyl ether methacrylate) 789
- ω -functionalized poly(*tert*-butyl acrylate) 778
- functional monomers 240, 288, 420, 424, 425, 433, 717, 755, 954, *see also* reactive groups
- functional RAFT agents 775, 807, 1004, *see also* reactive groups
- aromatic dithioesters 235
- trithiocarbonates 390–412, 436, 759
- furan 437, 769, 853, 955, 991
- furan-protected maleimide 437, 853, 991
- reactive group 955
- g**
- galactopyranose-functionalized poly(*N*-acryloylmorpholine) macroRAFT 715, 730, 731
- galactosamine 832
- gel permeation chromatography (GPC) 35, 36, 41, 49, 50, 52, 53, 649, 651, 653, 658, 669, 695, 818, 819, 908, 909, 993, 996, 998–1000, *see also* size exclusion chromatography (SEC)
- Gilson Pipetmax 268
- liquid handling robot 632
- D*-glucono- δ -lactone 691
- D*-glucosamine 832, 851, 853
- glucose oxidase (GOx) 654, 656, 691–693, 695, 696, 1057, 1058
- glutathione (GSH)-based solvophilic macroRAFT 720, 723, 857
- glycerol methacrylate, *see* glycerol monomethacrylate
- glycerol monomethacrylate (GMMA, 2,3-dihydroxypropyl methacrylate, glycerol methacrylate) 242–244, 268, 393–394, 411, 414, 561, 572, 686–687, 697–699, 711, 718, 719, 722–724, 726, 735–737, 844, 1041–1042
- with dithiocarbamates 561, 572
- with dithioesters 242–244, 268
- with trithiocarbonates 393–394, 411, 414
- glycidyl methacrylate (GMA) 241–243, 245, 305, 377, 383, 387, 391–393, 557, 558, 615, 711, 717–719, 723–724, 727, 737, 844, 1033, 1053
- glycomonomers 238, 273, 278, 281, 284, 287, *see also* 6-*O*-methacryloyl mannose
- with dithioesters 40, 238, 273, 278, 281, 284, 287
- with trithiocarbonates 417, 418, 420, 423
- glycopolymers 832–833, 847, 848, 851, 853, 857
- poly(2-(2,3,4,6-tetra-*O*-acetyl- β -D-glucosyloxy)ethyl methacrylate) (PACGlcEMA) 857
- gradient copolymers 26, 207–208, 517, 521, 533, 948–949
- modelling of 207–208
- graft copolymers 4, 307, 323, 433, 437, 522, 523, 587, 727, 782, 956–964, 1027, 1078
- graft-block copolymers 1022
- graft density 1017, 1021, 1029–1031, 1033, 1034, 1039
- general guidelines 963–964
- grafting from 960–963
- grafting onto 959–960
- grafting through 958–959
- graft water-soluble acrylates 624

- grafting from biomolecules 699–701
graphitic carbon nitride (g-C₃N₄) 622
group transfer polymerization (GTP) 755
- h**
- halo-olefins 514–515, *see also* vinyl chloride, vinyl halides, vinylidene chloride, vinylidene fluoride
- hemoglobin (Hb) 654, 655
- hetero Diels-Alder reaction 322, 769–773, 777, 778, 945, 947, 955
- heterodienophiles
(diethoxyphosphoryl)methanedithioates 769, 770
pyridine-2-carbodithioates 769
triazolinediones 771–772
- heterogeneous polymerization, *see also* PISA
dispersion polymerization 38, 187, 194, 195, 211, 440–442, 508, 650, 686, 691, 696–699, 709–713, 718–721, 723–726, 729–731, 733–736, 898, 900–901, 996, 998–999, 1006
emulsion polymerization 34–37, 42, 187, 195, 205, 209–210, 313–314, 413–414, 507–509, 587, 710, 1078–1080, 1092, 1095–1097, 1100, 1105, 1108–1112, 1114, 1121, 1125, 1129, 1132
inverse emulsion polymerization 532
miniemulsion polymerization 195, 205–206, 507, 509, 515, 908
precipitation polymerization 710–712
suspension polymerization 195–196
- hexa-2,4-dien-1-ol 772, 773
(2*E*,4*E*)-hexa-2,4-dien-1-yl prop-2-yn-1-yl succinate 777
- 1,1,1,3,3,3-hexafluoroisopropanol (HFIP) 838
- 1-hexanethiol 779
- hexyl isocyanate 782
- hierarchical porous polymers (HPP) 873
- high throughput (HTP) polymer libraries 630, 631, 633
- high throughput and high output experimentation (HT/HO-E)
antimicrobial agents 1064–1065
bottlenecks 1052
fundamental aspects and limitations 1052–1053
kinetic investigations 1053–1055
nanomedicine and drug delivery systems 1059–1065
star polymer synthesis 1000
synthesis and accessibility 1056, 1059
versatility 1051
- Hildebrand solubility parameter 898, 900
- horseradish peroxidase (HRP) 693–695, 713
- hydrazine hydrate 527, 773, 782, 783, 790
- 2-hydroethyl acrylate (HEA) 424, 615, 685, 691–693, 700, 845, 1057
- hydrogels and 3D printing 633–634
- hydrogen peroxide 321, 447, 528, 591, 624, 691, 693, 695, 696, 713, 769
- hydroperoxide end-terminated (co)polymers 768
- 2-hydroxybutyl methacrylate (HBMA) 727, 730
- 2-hydroxyethyl acrylate (HEA) 615, 653, 685, 1057
- 2-hydroxyethylamine 722
- 2-hydroxyethyl 2-((ethoxycarbonothioyl)thio)-2-methylpropanoate (2-ethoxythiocarbonylsulfanyl-2-methyl-propionic acid 2-hydroxyethyl ester) 498, 519, 693
- 2-hydroxyethyl methacrylate (HEMA) 38, 126, 523, 615, 693, 846, 900, 1057
- hydroxy-functional dithioesters 308
- hydroxy-functional trithiocarbonates 436

- N*-(2-hydroxyethyl)acrylamide (HEAm)
394, 398, 399, 405, 440, 696, 727, 840
- N*-(2-hydroxyethyl methacrylate (HEMA)
198, 416, 424, 701, 853, 902
- 2-hydroxypropyl)methacrylamide
(HPMAm) 45, 53, 440, 686, 687, 691, 693, 696–699, 711, 718, 719, 723, 725, 737, 831, 841, 847, 853, 854, 966
- 2-hydroxypropyl methacrylate (HPMA)
45, 53, 440, 686, 687, 691, 693, 696–699, 711, 718, 719, 723, 725, 737, 841, 847, 853, 854
- 4-hydroxystyrene 400, 421, 566, 572
- hypophosphite salts 766
- i**
- imidazole
as isocyanate protecting group 841
in trithiocarbonate synthesis 413
- 1*H*-imidazole-1-carbodithioates 552, 557, 561, 576–578
- 1*H*-imidazole-4-carbodithioates 209, 210
- indene monomers 628, 820–823
single unit monomer insertion (SUMI)
628, 820–823
- industrial applications of RAFT
531–532, 1077–1136
- biomedical applications 1132
patents 1103–1106, 1108–1110, 1112–1114, 1117, 1119, 1121, 1122, 1125–1131
- electronic applications. 1132
patents 1105, 1108, 1112, 1113, 1115, 1118, 1122, 1124, 1125, 1127, 1128
- emulsion polymerization 1100
patents 1105, 1108–1112, 1114, 1120, 1121, 1125, 1128, 1129
- end-group removal
patents 1133–1136
- macromonomer RAFT polymerization
1077–1086
- RAFT agents 1087–1088, 1101–1102
- stars and microgels 1132
- thiocarbonylthio RAFT polymerization
1082, 1086–1136
key patents 1086, 1089, 1090–1099
- iniferter 312, 553, 554, 611, 617, 624, 818, 941, 1089, *see also*
photoiniferter
- inimer RAFT agents 301, 305–306, 433–435, 629, 815–816, 966, 987
- acrylates 305
- methacrylates 305, 434, 628, 629, *see also* 1-(methacryloyloxy)propan-2-yl 4-(((butylthio)carbonothioyl)thio)-4-cyanopentanoate
- methacrylamides 434, *see also* 1-methacrylamidopropan-2-yl 4-(((butylthio)carbonothioyl)thio)-4-cyanopentanoate
- in self-condensing vinyl polymerization
dithioesters 301, 305–306
enzyme responsive 966
trithiocarbonates 433, 434–435, 987
- styrenes 209–210, 305–306, 434–435, 987, *see also* 4-vinylbenzyl 1*H*-imidazole-4-carbodithioate
- in synthesis of sequence-defined
polymers 815–816
- initiators, *see also* azo-compound
initiators, peroxide initiators,
photoinitiators
- photo-redox initiators 312
- photoiniferters (RAFT agents) 109, 130–131, 311, 438–439
- γ -radiation 109
- redox initiators 4, 47, 109, 506–507, 591, 648–656, 672, 684, 695
- Fenton's reagent 648, 652–654
bio-Fenton's reagent 654
- initer 553
- intermediate activated monomers (IAMs),
see N-vinylcarbazole

- intermediate radical termination (IRT)
59, 60, 76, 77, 80, 86, 90, 91,
119–126, 171, 187, 188, 191–194,
200–201, 205, 206, 225, 226, 288,
373, 425, 443
- inverse mini-emulsion periphery RAFT
polymerization (IMEPP)
approach 851
- iodine transfer polymerization (ITP) 708
- ionization energy (IE) 176
- iridium photocatalyst, *see* tris(2-phenyl-
pyridine)iridium(III)
(*fac*-Ir(ppy)₃), PET-RAFT
- isocyanate (NCO) group
functional monomer 294, 297, 425,
428, 840–842, *see also*
2-isocyanatoethyl acrylate,
2-isocyanatoethyl methacrylate
functional RAFT agent 409
reagent 782, *see also* allyl isocyanate,
hexyl isocyanate
selenol–isocyanate reaction 786
thiol–isocyanate reaction 774, 780–782
- 2-isocyanatoethyl acrylate 840
- 2-isocyanatoethyl methacrylate 782,
841
protected monomer, *see* azole-*N*-
carboxamide methacrylates
- isoprene 242, 280, 393, 395, 396, 399,
421, 424, 434, 944, 945, 1034,
1055
high throughput synthesis 1053
mScroRAFT agents 944, 945
SI-RAFT 1034
- isopropenyl acetate (IPAc) 842–843
- j**
- junction-functionalized diblock
copolymers 807
- k**
- ketoform reaction 423, 580, 581
- ketone functionalized particles 724
- kinetic modeling, *see* mathematical
modeling
- kinetic rate coefficients, *see* rate
coefficients
- Krytox 900, 902, 903
- l**
- lauroyl peroxide (LPO, dodecanoyl
peroxide) 318, 320, 321,
443–446, 530–531, 589, 765–767,
1135
in RAFT end-group removal 318, 320,
321, 443–446, 530–531, 589,
765–767, 1135
in SUMI 1120
- lauryl methacrylate (LMA, dodecyl
methacrylate) 416, 711, 725,
774, 853, 1034, 1059
- leaving group ability 96–97, 100, 105,
109–111, 373, 443, 810
- less activated monomers (LAMs) 46, 47,
116–118, 234, 239, 287–288, 297,
301, 371, 380–382, 424, 431, 447,
551, 552, 557, 572–575, 582, 583,
585, 587, 591, 593, 693, 764, 806,
807, 809, 934, 935, 936, 938, 939,
1100, *see also* vinyl amides, vinyl
esters, vinyl imides
- Lewis acid-catalyzed mechanisms 1160
- ligands
- 2,2'-bipyridine (bpy), *see also* copper
activators
Cu^{II}(DC)₂/bpy 668
tris(2,2'-bipyridyl)ruthenium(II)
chloride (Ru(bpy)₃Cl₂) 312,
618–619, 635, 689–691
- 4,4'-dinonyl-2,2'-bipyridine (dNbpy),
see also copper activators
CuX/ dNbpy 657
- N,N,N',N',N''*-pentamethyldiethyl-
enetriamine (PMDETA), *see*
copper activators
CuX/PMDETA 313, 656–658, 669,
759, 1024
- tris[2-(dimethylamino)ethyl]amine
(Me₆TREN), *see also* copper
activators

- ligands (*contd.*)
- CuX/Me₆TREN 120, 517, 657, 658
 - tris(2-phenylpyridine)
 - tris(2-phenylpyridine)iridium(III) (*fac*-Ir(ppy)₃) 312, 612, 617–619, 623, 627, 628, 1176
 - tris(2-pyridylmethyl)amine (TMPA), *see also* copper activators
 - CuX/TMPA 657–659, 669–670
 - light-harvesting polymers 307, 807
 - limonene 381, 620, 628
 - live cell graft polymerizations 634–635
 - Saccharomyces cerevisiae* 635
 - living cationic polymerization 519, 1160–1161
 - livingness (*L*) (fraction of living chains, end-group fidelity)
 - living polymerization 15
 - PET-RAFT polymerization 618
 - RAFT polymerization 116, 128, 225, 310–311, 552, 938–942
 - living polymerization
 - RAFT mechanism transformation 307
 - terminology 15–18
 - lower critical solution temperature (LCST) 514, 519, 714, 1027, 1029
 - lysozyme 526, 528, 624, 701, 791, 991, 1027
- m**
- (macro)alkoxyamines 205, 296, 308–310, 403, 438, 429, 717, 737, 768
 - addition-fragmentation coupling 768
 - ESARA 310, 438, 768
 - NMP inimer 429
 - RAFT-*T*-NMP 308–310, 438
 - macro-chain transfer agent (macroCTA), *see* macroRAFT agent
 - macromolecular design by interchange of xanthates (MADIX) 1, 495, 532, 553, 710, 806, 1089
 - macromolecular thiol
 - catalyzed thiol-michael additions 777–780
 - disulfide reactions 787–790
 - oxidative coupling 773
 - pristine polymeric thiol 773
 - radical thiol-ene reaction 775–776
 - thiol-epoxy ring-opening 782–783
 - thiol-halo substitution 783–786
 - thiol-isocyanate modification 780–781
 - tributylphosphine (PBu₃) 773
 - tris(2-carboxyethyl)phosphine (TCEP) 773
 - macromonomer RAFT agent (macromonomer CTA) 936, 1078
 - synthesis of
 - catalytic chain transfer (CCT) 1079
 - macromonomer RAFT polymerization (sulfur-free RAFT polymerization) 936, 1077–1082
 - emulsion polymerization 936, 1078–1079
 - industrial use 1077–1082
 - mechanism 1078
 - nanogel synthesis 1023
 - macromonomers 96, 167, 297, 298, 425–430, 433, 437, 719, 727, 730, 782, 786, 793, 806, 936, 958, 959, 963, 994, 1077–1082, 1083, 1085, 1086
 - in arm-first star synthesis 994–995
 - with dithioesters 297–299
 - in grafting through 958–959
 - methacryloyl 298–299, 430, 958
 - peptide 430
 - polycaprolactone (PCL) 269
 - poly(ethylene glycol) (PEG) 297–298, 425, 958
 - poly(dimethylsiloxane) (PDMS) 290, 958
 - polyoxazoline 298
 - with trithiocarbonates 425–430
 - macroRAFT agent (macro-chain transfer agent, macroCTA) 98, 102, 105, 118, 310, 680, 681, 683,

- 685–687, 691, 696–701, 768, 933, 934, 937, 938, 939, 943–994, 948, 986, 994
- cores 986
- macro-RAFT SUMI products 810
- macroRAFT type PISA 712
- MALDI-TOF mass spectrometry 774
- maleic anhydride (MAH) 241, 242, 244, 247, 249, 253, 254, 264, 265, 280, 307, 384, 385, 395, 396, 397, 399, 400, 430–431, 433, 559, 807, 809, 810, 830, 842–843, 809, 810, 883, 891, 915
- cyclocopolymerization with divinyl ether 430–431
- single unit monomer insertion (SUMI) 307, 433, 807, 809, 810
- styrene/maleic anhydride copolymer 241, 242, 247, 249, 264, 265, 384, 385, 396, 399, 400, 842, 891, 915
- maleimide crosslinker 845, 947, 955, 956
- in star synthesis 955, 956
- maleimide end-functional polymer 633, 780, 991
- thiol-maleimide reaction 947, 991
- maleimide-functional initiator
- radical-induced coupling 991
- maleimide-functional RAFT agent 700, 1027
- furan-protected 402, 437
- in SI-RAFT 1025, 1027
- maleimide monomers 280, 420, 588, 589, 627–628, 433, 777
- furan-protected 291, 853
- N*-methylmaleimide (NMMI) 245, 265, 280, 401, 559
- 1-(2-oxotetrahydrothiophen-3-yl)-1H-pyrrole-2,5-dione (thiolactone-functional maleimide) 843
- N*-phenylmaleimide (NPMI) 256, 264, 265, 280, 391, 393, 401, 559, 589, 820, 821
- single unit monomer insertion (SUMI) 381, 401, 433, 627–628, 807, 817, 820–823
- mathematical modeling (kinetic modeling, numerical simulation)
- ab initio* kinetic modelling 165–167
- deterministic modeling 188–204
- diffusion coefficients 196–198
- full molecular weight distributions 198–204
- explicit integration methods 199–201
- Predici® software 201–204
- probability generating function 201, 202
- hybrid methods 206
- matrix-free polychloroprene grafted silica nanocomposites 1034
- method of moments 188–190
- heterogeneous systems 194–196
- homogeneous systems 190–194
- polymer networks
- kinetic random branching theory (KRBT) 915–918
- trifunctional molecule approach 907–910
- multifunctional molecule approach 910–915
- polymerization processes
- branched copolymerizations 209–210
- CSTRs/PFR 208–209
- microwave-assisted RAFT polymerization 210–211
- semibatch polymerization 206–208
- Predici® software 64, 65, 73, 103, 124, 165, 187, 201–204, 906
- stochastic modeling 204–206
- Monte Carlo 204–205
- 2-mercaptoethanol (2M) 857
- 6-mercapto-1-hexanol 779
- 3-mercaptopropionic acid (3M) 760, 857
- 3-mercaptopropyl polyhedral oligomeric silsesquioxane 779
- merocyanine 613, 629
- metal free photocatalysts 622–625
- metalloporphyrin 617, 620, 628

- meta-tetra(hydroxyphenyl)-chlorin (*m*-THPC) 822
- methacrylamides 43–46, 223, 233, 239, 240, 278, 325, 380, 413, 416, 420, 441, 449, 532, 552, 629, 683, 686, 691, 693, 724, 831, 838, 840, 841, 846, 847, 850, 851, 966, 985, 1057
- amino-functional, *see* *N*-(2-aminoethyl) methacrylamide hydrochloride, *N*-(3-aminopropyl)methacrylamide hydrochloride
- azide functional, *see* *N*-(3-azidopropyl) methacrylamide
- with dithioesters 223, 233, 234, 240, 244, 252, 278–279, 293
- experimental procedures
- N*-(3-aminopropyl)methacrylamide in aqueous solution 45
 - N*-(2-hydroxypropyl)methacrylamide in aqueous solution 45
 - poly(benzyl methacrylate)-*block*-poly(*N*-(2-hydroxypropyl) methacrylamide) in solution 53
- functional methacrylamides
- azide, *see* *N*-(3-azidopropyl) methacrylamide
 - hydroxyl, *see* *N*-(2-hydroxypropyl) methacrylamide
 - glycomonomers 278
 - protected amine 293, *see also* *N*-(2-aminoethyl)methacrylamide hydrochloride (AEMAm), *N*-(3-aminopropyl)methacrylamide hydrochloride (APMAm)
 - protected thiol 292
- RAFT agent selection 27–28, 43–45, 112
- with trithiocarbonates 380, 391–393, 397, 399, 404, 405, 417, 420
- 1-methacrylamidopropan-2-yl 4-(((butylthio)carbonothioylthio)-4-cyanopentanoate 434
- methacrylates 38, 239, 268, 416, 701, 767, 816, 834, *see also* butyl methacrylate (BMA), methyl methacrylate (MMA), lauryl methacrylate (LMA), octadecyl methacrylate (ODMA)
- crosslinking monomers 202
- with dithiocarbamates 557, 558–562
- balanced activity 561–562, 582–583
 - switchable 568
- with dithioesters 239–247, 260–263
- aliphatic dithioesters 564
 - aromatic dithioesters 241–247
 - bis-dithioesters 260–263
 - functional dithioesters 250–259
- experimental procedures
- benzyl methacrylate in solution 41
 - 6-*O*-methacryloyl mannose in solution 40
 - methyl methacrylate in solution 39, 40, 41
 - poly(*N*-(2-hydroxypropyl)methacrylamide)-*block*-poly(benzyl methacrylate) 53
 - poly(*N*-isopropylacrylamide-*co*-sodium methacrylate), in solution 50
 - poly(methyl methacrylate)-*block*-poly(butyl acrylate) 51
 - poly(methyl methacrylate-*co*-trimethylsilyl methacrylate-*co*-*t*-butyl(dimethyl)silyloxystyrene), in solution 49
 - poly(methacrylic acid-*co*-oleyl methacrylate) in solution 49
 - poly(triphenylmethyl methacrylate-*co*-methacrylic acid), in solution 49
- functional methacrylates
- alkyne 289, *see also* propargyl methacrylate
 - azide 289
 - epoxy, *see* glycidyl methacrylate (GMA)

- hydroxyl, *see* glycerol monomethacrylate (GMA) 2-hydroxyethyl methacrylate (HEMA), 2-hydroxypropyl methacrylate (HPMA)
- glycomonomers 273
- isocyanate-functional, *see* 2-isocyanatoethyl methacrylate 294
- oleate 844
- PEG, *see* poly(ethylene glycol) monomethyl ether methacrylate (PEGMA)
- protected amine, *see* (2-aminoethyl) methacrylate hydrochloride (AEMA)
- protected thiol 292
- silane 294
- tertiary amine, 2-(dimethylamino) ethyl methacrylate (DMAEMA) 292–293
- inimers
- ATRP 295
- RAFT 292, 305
- RAFT agent selection 27–28, 112
- star polymers 954
- with trithiocarbonates 383–384, 391–397, 416, 420
- bis-trithiocarbonates 388–389
- non-symmetric trithiocarbonates 391–397
- symmetric trithiocarbonates 383–384
- functional, trithiocarbonates 379, 408
- with xanthates 511–512, 532
- semi-batch emulsion polymerization 512
- methacrylic acid (MAA) 49–50, 209, 321, 323, 691, 711, 714, 782, 815, 831, 855, 1023, 1029, 1061, 1062, 1064
- NHS ester, *see* *N*-methacryloyloxysuccinimide (MAS)
- methacrylic poly(*N*-acryloylmorpholine) macromonomers 782
- methacryloyl chloride 839
- methacryloyl-functionalised poly(ethylene glycols) 958
- 6-*O*-methacryloyl mannose 40, 945
- [2-(methacryloyloxy)-ethyl] dimethyl(3-sulfo)propyl ammonium hydroxide 699
- 2-(methacryloyloxy)ethylisocyanate, *see* 2-isocyanatoethyl methacrylate
- 2-(methacryloyloxy)ethyl oleate (MAEO) 719, 844
- 1-(methacryloyloxy)propan-2-yl 4-(((butylthio)carbonothioyl)thio)-4-cyanopentanoate [2-(2-(butyltrithiocarbonate)-propionate)ethyl methacrylate] 628, 629
- N*-methacryloyloxysuccinimide (NMS, methacrylic acid, NHS ester) 242, 290, 426, 830–832, 855
- method of moments (MM) 188
- heterogeneous systems 194–196
- dispersion polymerization 195
- emulsion polymerization 195
- suspension polymerization 195–196
- homogeneous systems 190–194
- 2-methoxyethylacrylate (MEA) 650, 696
- methyl acrylate (MA) 39, 40, 51, 75, 76, 117, 130, 164, 172, 205, 234, 380, 421, 505, 507, 552, 556, 572, 615, 622, 648, 666, 696, 759, 808, 814, 844, 986, 987, 1053, 1175
- rate coefficients for radical addition (k_i , k_p) 107
- 2-methyl-acrylic acid 6-(3,4-dicyanophenoxy)-hexyl ester 850
- Methylene-bis-acrylamide (MBA) 526, 717, 907, 996
- methyl 2-bromo-2-propanoate 783
- 2-methylene-4-phenyl-1,3-dioxolane 725

- methyl 2-((ethoxycarbonothioyl)thio)propanoate (*O*-ethyl-*S*-(1-methoxycarbonyl)ethyl)dithiocarbonate, Rhodixan A1) 47, 496, 510–516, 857
- application scope 496, 510–516
 - experimental procedure
 - vinyl acetate in solution 47
 - transfer coefficients 510
- methyl 2-((isopropoxycarbonothioyl)thio)acetate (methyl (isopropoxycarbonothioyl)sulfanylacetate) 33 52
- methyl 3-mercaptopropionate (M3) 857
- methyl 2-((methyl(pyridin-4-yl)carbamothioyl)thio)propanoate 30, 47, 51, 52
- application scope 569
 - experimental procedure
 - methyl acrylate in solution 51
 - poly(methyl acrylate)-*block*-poly(*N*-vinylcarbazole) in solution 51
 - styrene in solution 52
 - polystyrene-*block*-poly(vinyl acetate) in solution 52
- methyl methacrylate (MMA) 31, 38, 88, 99, 104, 111, 117, 130, 147, 148, 154, 172, 193, 234, 380, 381, 505, 512, 551, 552, 556, 648, 666, 810, 846, 966, 1056, 1175
- experimental procedures
 - methyl methacrylate in solution 39, 40, 41
 - poly(methyl methacrylate)-*block*-poly(butyl acrylate) 51
 - poly(methyl methacrylate-*co*-trimethylsilyl methacrylate-*co*-*t*-butyl(dimethyl)silyloxystyrene), in solution 49
 - mathematical modeling 193–194, 197–198, 208
 - model system with dithiobenzoate 88–89
 - RAFT equilibrium constant 88
 - rate coefficients for radical addition (k_i, k_p) 107
- methyl methanethiolsulfonate (methyl methanesulfonothioate) 317
- methyl(pyridin-4-yl)carbamodithioates (*N*-methyl-*N*-(4-pyridinyl)dithiocarbamates) 51, 116–117, 556–557, 567–571, 583–585, *see also* cyanomethyl methyl(pyridin-4-yl)carbamodithioate, methyl 2-((methyl(pyridin-4-yl)carbamothioyl)thio)propanoate
- application scope 116–117, 568–571
 - canonical forms 583–584
 - monomers 568–571
 - polyMAM-*block*-polyLAM synthesis 116, 584
 - transfer coefficients 556
- N*-methyl-*N*-(4-pyridinyl)dithiocarbamates, *see* methyl(pyridin-4-yl)carbamodithioates
- 4-methylstyrene 241, 249, 264, 280, 411, 421
- Me₆TREN, *see* tris[2-(dimethylamino)ethyl]amine, *see* copper activators
- micellar nucleation 696, 711
- miktoarm stars 525, 950, 951, 955, 956, 986, 994, 1002, 1009
- missing-step reaction 77–86, 88–90
- Mitsunobu reaction 238, 413, 833
- molar mass distribution (MMD) 61, 187, 204, 208, 210, 211, 445, 514, 516, 905, 908
- molecular orbital calculations, *see* quantum-chemical calculations
- molecular weight distribution (MWD) 25, 31, 32, 36, 37, 48, 97, 106, 109, 111, 115, 116, 119, 122, 125, 126, 129, 187, 318, 511, 513, 515, 519, 522, 523, 620, 625, 636, 755, 760, 764, 765, 782, 823, 831, 832, 836, 838, 877, 905, 953, 962, 1162

- monomers, *see* intermediate activated monomers (IAMs), less activated monomers (LAMs), more activated monomers (MAMs)
- more activated monomers (MAMs) 2, 38, 114, 226, 371, 551, 693, 712, 764, 806, 934, 1100
- 1,1-disubstituted MAMs 239–240, 280, 416, 420, 557, 572, *see also* methacrylamides, methacrylates
- monosubstituted MAMs, *see* acrylamides, acrylates, acrylonitrile, diene monomers, styrenic monomers 240, 280, 421, 424
- multiblock copolymers 105, 521, 532, 618, 626, 652, 805, 806, 878, 942–943, 968, 1065, 1176
- from multiRAFT agents 937
- by PET-RAFT 618, 626, 632
- by sequential addition 942–943, 1057, 1065
- multicomponent reactions (MCR) 849–850
- multiolefinic monomers 301, 432, 874, *see also* divinyl monomers
- multiple unit monomer insertion (MUMI) 813, 814
- multiRAFT agents (multifunctional RAFT agents) 879, 895, 937
- multistep polymerization 877, 878
- multiwalled carbon nanotubes (MWNTs) 1007, 1008
- Mycobacterium smegmatis* 631, 632
- n**
- N-carboxyanhydrides (NCA) 436–437, 767, 790, 983, 995
- γ -benzyl-L-glutamate
- N-carboxyanhydride (NCA) 767, 790
- RAFT-*F*-ROP 436–437, 767, 790
- N-isopropylacrylamide (NIPAM) 44, 46, 50, 210, 234, 380, 505, 508, 552, 615, 616, 634, 650, 686, 711, 780, 783, 810, 843, 846, 944, 949, 965, 991, 996, 1003, 1063
- 3-nitrobenzonitrile 667
- o*-nitrobenzyl functionalities 629
- o*-nitrobenzyl methacrylate 629
- p*-nitrophenyl ester 835–836
- S*-nitrosothiol species 790
- 2-nitro-*m*-xylene 667
- nanoclay attapulgite (ATP) 649, 650
- NHS ester, *see* N-hydroxysuccinimide (NHS) ester
- N-hydroxysuccinimide (NHS) ester 815, 830–832, 1004
- of acrylic acid, *see* N-acryloyloxysuccinimide (NAS)
- in bioconjugation 857
- by copolymerization of multifunctional monomers 966–967
- hyperbranched polymers 517, 526, 964–968, 986
- of methacrylic acid, *see* N-methacryloyloxysuccinimide (NMS)
- NHS functional monomer 290, 426, 830–832
- NHS functional RAFT agent 251–252, 402–405, 500, 857, 860
- dithioester 251–252
- trithiocarbonate 402–405
- xanthate 500
- by self-condensing vinyl polymerization (SCVP) 964–966, 986
- of 4-vinylbenzoic acid, *see* 2,5-dioxopyrrolidin-1-yl 4-vinylbenzoate
- nitroxide-mediated polymerization (NMP) 118, 176–178, 277, 297, 308, 309–310, 421, 425, 436, 438, 525, 552, 554, 708, 712, 717, 726, 755, 905, 939, 962, 964, 983, 985
- RAFT mechanism transformation (RAFT-*F*-NMP) 438
- nitroxide-terminated polymers 768

- NMR spectroscopy 36, 41, 49, 52, 53, 81, 85, 513, 515, 521, 700, 760, 774, 779, 782, 783, 785, 786, 831
- non-activated alkenes 807
- non-Aldol carbonyl reaction 780
- non-homopolymerizable monomers 820
- non-symmetric trithiocarbonates 372, 373, 380, 387, 389–390, 401, 404
- norbornenyl-functional RAFT agents 760
- xanthate 759
- nucleophile (phosphine) catalyzed mechanism 777, 779
- numerical simulation, *see* mathematical modeling
- O**
- octadecyl acrylate (ODA) 42
- blocks 383, 385
- copolymers 262, 388, 397
- experimental procedure 42
- homopolymer 42, 385, 397
- with trithiocarbonates 42, 385, 397
- 1-octanethiol 834, 1166
- octanol-water partition coefficient 726
- octylamine 779
- oleic acid decorated nano-objects 719
- oligo(ethylene glycol) acrylate (OEGA) 43, 69, 240, 421, 425, 615, 690, 693, 699, 701, 727, 730, 943, 1175
- oligo(ethylene glycol) methacrylate (OEGMA) 634, 701, 1041, 1059, 1060
- oligomer separation 813
- organotellurium-mediated radical polymerization (TERP) 708
- oxygen tolerance
- PET-RAFT 624, 691
- P**
- para-fluoro substitution reaction (PFTR) 194, 723, 834, 835
- parathyroid peptide hormone, (PTH) 612, 622, 634, 761, 762
- parent hydrogels 634
- Passerini reaction-derived methacrylates 720, 835
- PEG dithiol 633
- pendent double bonds (PDB) 873, 906
- pentafluorobenzyl (PFB) ester monomers, *see* pentafluorobenzyl methacrylate (PFBMA)
- pentafluorobenzyl methacrylate (PFBMA) 723, 834, 835
- pentafluorophenyl (PFP) ester 830, 832–835
- monomers, *see* pentafluorophenyl methacrylate, pentafluorophenyl 4-vinylbenzoate
- nanoparticle shell-units 720
- RAFT agents 789, 837
- pentafluorophenyl ester-functionalized R-group dithiobenzoate 789
- pentafluorophenyl methacrylate (PFPMA) 38, 240, 720, 723, 832–835, 848, 851, 854
- decorated particles 720
- pentafluorophenyl-(4-phenylthio-carbonylthio-4-cyanovalerate) 837
- pentafluorophenyl 4-vinylbenzoate (PFP4VB) 833
- pentafluorostyrene (PFSty) 421, 786, 834, 844
- based homo- and copolymers 834
- pentamers 818, 820
- N,N,N',N''*-pentamethyldiethylene-triamine (PMDETA), *see* copper activators
- peroxide initiators 32, *see also*
- ammonium persulfate (APS), benzoyl peroxide (BPO), *t*-butyl hydroperoxide, cumene hydroperoxide (CHP), lauroyl peroxide (LPO)
- 'solvent-initiated' RAFT 654

- PET-RAFT, *see* photoinduced energy or electron transfer-RAFT
- pH 736
- sensitive DMAEMA block 698
 - zwitterionic pH-responsive copolymers 843
- phenazine organophotocatalysts 617
- 2-((phenylcarbonothioyl)thio)acetic acid industrial use 1102
- phenothiazine 612, 617, 622
- 1-phenylethyl benzodithioate 109, 111, 120, 122, 124, 205, 228, 229, 247, 307, 888
- application scope 247
 - chain transfer coefficient 228, 229
 - crosslinking polymerization 888
- N*-phenyl maleimide (PMI) 820, 821
- 2-phenylpropan-2-yl benzodithioate (cumyl dithiobenzoate, CDB) 33, 34, 40, 41, 49, 67, 108, 126, 149, 164, 228, 233, 241, 307, 314, 658, 832, 836, 913
- application scope 27, 241
 - experimental procedure
 - benzyl methacrylate in solution 41–42
 - 6-*O*-methacryloyl mannose in solution 49
 - styrene in solution 33–35
 - chain transfer coefficient 228
 - industrial use 30
 - side reactions 126
 - stability 126
- 2-phenylpropan-2-yl 2-phenylethane-dithioate (cumyl phenyldithioacetate, CPDA) 105–106, 264
- application scope 264
- 3-phenylpropyl methacrylate 720
- phosphate buffered saline (PBS) 691, 692, 693, 695
- photoinduced energy or electron transfer-RAFT (PET-RAFT) 311–312, 438–439, 506, 507, 617–625, 628, 629, 635, 688–690, 701, 817, 991, 1055, 1064–1065
- with dithiobenzoates 311–312
 - high throughput experiments 629–633, 1955, 1064–1065
 - oxygen tolerance 442, 624, 691
 - photocatalysts
 - eosin Y (EY) 312, 612, 623–625, 630, 635, 689–691, 701, 1027
 - pheophorbide A 617, 628
 - rose bengal 624, 719
 - tris(2,2'-bipyridyl)ruthenium(II) chloride (Ru(bpy)₃Cl₂) 312, 618–619, 635, 689–691
 - tris(2-phenylpyridine)iridium(III) (fac-Ir(ppy)₃) 312, 612, 617–619, 623, 627, 628, 1176
 - zinc(II) (2,3,7,8,12,13,17,18-octaethyl-5,10,15,20-tetraphenylporphyrin) (ZnOETPP) 621–622
 - zinc-iron oxide semiconductor (Zn_{0.64}Fe_{2.36}O₄) 616
 - zinc(II) meso-tetra(*N*-methyl-4-pyridyl) porphine tetrachloride (ZnTPMPyP) 629, 691
 - zinc(II) meso-tetra(4-sulfonatophenyl)porphyrin) (ZnTPPS⁴⁻) 621, 689, 691
 - zinc(II) 5,10,15,20-tetraphenyl-21H,23H-porphine (zinc tetraphenyl porphyrin, ZnTPP) 312, 617, 620, 628–631, 991, 1055
 - star polymer synthesis 991–994
 - single unit monomer insertion (SUMI) 307, 439, 625–628, 817–820
 - with trithiocarbonates 438–439
 - photoiniferter process 109, 130–131, 311, 553, 438–439, 613–625, 817
- photoiniters 553
- photoinitiation 130, 307, 311–312, 382, 438–439, 575, 585, 612–613, 685

- photoinitiators, *see also* photoinduced energy or electron transfer (PET)-RAFT, photoiniferter process
 in pulsed laser photolysis (PLP) 60–65
 type I 612–613
 acetophenones 612
 acyl-phosphine oxides 612
 benzoin ethers 612
 benzoin ketals 612
 benzophenone 612
 2,2-dimethoxy-2-phenylacetophenone 760
 2-methyl-1-[4-(methylthio)phenyl]-2-morpholin-4-ylpropan-1-one 63, 70, 73–75, 85, 88
 type II 612
 photoRAFT polymerization, *see also* photoinduced energy or electron transfer (PET)-RAFT
 high throughput (HTP) polymer libraries 630
 hydrogels and 3D printing 633–634
 live cell graft polymerization 634–635
 photoiniferter polymerization 613–625
 photoinitiation 612–613
 photoredox catalysis 617–625
 single unit monomer insertion (SUMI) 625–628
 wavelength orthogonal polymerization 628–629
 photoRAFT-SUMI, *see* photoinduced energy or electron transfer (PET)-RAFT SUMI
 photoredox initiation, *see* photoinduced energy or electron transfer (PET)-RAFT
 photoresponsive monomer
 7-[4-(trifluoromethyl)coumarin] methacrylamide 691
 physisorption 903, 905, 1018
 Pickering emulsions 722, 739
 4-pinacoloborylstyrene (pBSt) 849
 PISA, *see* polymerization-induced self-assembly
 polyacrylamide gel electrophoresis (PAGE) 700
 polydentate amine ligand 656
 poly(ethylene glycol) (PEG) 38, 39, 240, 297, 557, 649, 650, 685, 693, 713, 721, 727, 737, 843, 847, 857, 944, 995, 996, 1009, 1057, 1064, 1175
 (poly(ethylene glycol) monomethyl ether)acrylamide 635
 poly(ethylene glycol) monomethyl ether acrylate (PEGA) 298, 685, 996
 blocks 249, 253
 copolymers 244
 with dithioesters 247, 248
 homopolymer 247, 248
 poly(ethylene glycol) monomethyl ether methacrylate (PEGMA) 416, 425, 440, 557, 650, 659, 693, 699, 721, 726, 788, 790, 847, 857, 858, 861, 958, 992, 995, 996, 1057, 1059, 1064
 thermo-responsive polymer 857
 poly(ethylene glycol) methyl ether thiol (PEG-SH) 857
 poly(ethylene oxide) (PEO) 37, 518, 683, 715, 1037–1038, *see also* poly(ethylene glycol) (PEG)
 polyhedral oligomeric silsesquioxane (POSS) 5, 777, 779, 1009
 poly(lactide) 944
 polymer, *see* under respective monomer
 polymer grafted nanoparticles 1022–1027, 1039–1042
 in composites 1030–1039
 synthesis
 assemble-crosslink-disperse 1009, 1339–1040
 PISA 1040–1042
 SI-RAFT 1022–1027
 polymer grafted surfaces, *see* surface-initiated RAFT (SI-RAFT)
 polymer monoliths 913
 polymer networks

- background 906–907
- mathematical modeling
 - kinetic random branching theory (KRBT) 915–918
 - multifunctional molecule approach 910–915
 - trifunctional molecule approach 907–910
- model co-networks 301, 304, 895
- RAFT crosslinking polymerization 301, 432, 876–898
 - crosslinking monomers (divinyl monomers, multivinyl monomers) 301, 303, 432, 879–898
 - RAFT agents (controllers) 879–898
 - synthesis pathways 877–879
 - structure and characteristics 875–876
 - supercritical carbon dioxide 898–904
- polymer–protein binding affinity 631
- polymeric CTA, *see* macroRAFT agent
- polymeric disulfides 773–774, 781, 790
- polymeric ionic complexes (PIC) 734
- polymeric Janus nanosheets (PJS) 1023, 1024
- polymeric *S*-nitrosothiol 790–791
- polymerization-induced self-assembly (PISA) 509, 629, 696, 679, 708–710, 712, 725, 710, 942, 1040, 1041
 - applications 738–740
 - cooperative assembly (PICA) 712–714, 733
 - dispersion polymerization 711
 - electrostatic self-assembly (PIESA) 713–715, 734
 - emulsion polymerization 710–711
 - from spherical to anisotropic block copolymer particles 725–727
 - hierarchical self-assembly (PIHSA) 713–715, 733
 - history/origins of 709–710
 - initiation in RAFT-PISA 712–713
 - macroRAFT type 712
 - parameters that impact the particle morphology 727–733
 - poly(methacrylic acid) (PMAA) 714
 - polymerization-induced cooperative assembly (PICA) 714
 - polymerization-induced electrostatic self-assembly (PIESA) 714
 - polymerization-induced hierarchical self-assembly (PIHSA) 714–715
 - polymerization-induced thermal self-assembly (PITSA) 713–714
 - post-polymerization morphological transitions/chain reorganization 735–738
 - precipitation polymerization 711–712
 - reactive/functional nano-objects 716–717
 - solvophilic macroRAFT 713
 - strategies to stir specific morphologies 733–735
 - surfactant-free latexes 715–716
- polymerization-induced thermal self-assembly (PITSA) 508, 713–714
- population balance equations (PBEs) 188–189, 199
- precipitation polymerization 710–712
- Predici® software, *see* mathematical modeling
- pristine polymeric thiol 773
- propacrylic acid (PAA, 2-methylenebutanoic acid) 392, 393, 395, 396, 440, 442
- propagation rate coefficient, *see* rate coefficients
- pro-TAD-functional trithiocarbonate 772
- propargyl acrylate 39, 421, 649, 779–780, 991
- propargyl methacrylate (PgMA) 523, 847–842
- 2-(propylthiocarbonothioylthio)-2-methylpropionoic acid (PMP) 691

- prop-2-yn-phosphocholine (PPhCh) 847
- protecting group
for isocyanate, *see* imidazole, azole-*N*-carboxamide methacrylates
for primary amine, *see*
t-butyloxycarbonyl (BOC) 129
for thiol, *see* benzothioic *S*-acid, *S*-(2-((*t*-butoxycarbonyl)amino)ethyl) methanesulfonothioate, 2,2'-dipyridyl disulfide
- protein-polymer conjugates 679, 699, 790–791
- protein-thermoreponsive polymer conjugates 761
- pseudo-kinetic rate constants method 907
- pulsed-laser polymerization (PLP) 60, 61, 65, 151, 201
SP-PLP-EPR method 60–61, 64, 71–72, 88, 90, 379
- 1*H*-pyrazole-1-carbodithioates 28, 561, 582–583, *see also* 2-cyanobutan-2-yl 3,5-dimethyl-1*H*-pyrazole-1-carbodithioate, cyanomethyl 3,5-dimethyl-1*H*-pyrazole-1-carbodithioate
application scope 28, 561–564, 582–583
- 1-pyrenemethyl methacrylate 738
- pyren-1-ylmethyl acrylate 859
- pyridine-2,6-dicarboxaldehyde 724
- 2-pyridyl disulfide end-group 406, 761–762, 788–789, 815–816
- 2-pyridyl disulfide functional monomer 292, 427, 842, 843, 855–857
acrylamide 395, 427
acrylate 292, 392, 427
copolymer 243, 244, 392, 395, 855–857
with vinyl lactam 855–857
methacrylamide 244, 292, 427
methacrylate 39, 243, 244, 292, 395, 427, 843, 856–857
- 2-pyridyl disulfide functional RAFT agent 406
- 1*H*-pyrrole-1-carbodithioates 552, 577, 578, 581, 582, 588, 593, 894, *see also* benzyl 1*H*-pyrrole-1-carbodithioate, 2-cyanopropan-2-yl 1*H*-pyrrole-1-carbodithioate
application scope 558–559
crosslinking polymerization 894
synthesis of 577–579
- q**
- quantum-chemical calculations 60, 65–66, 72, 79, 87–88, 90, 139–180
ab initio kinetic modelling 152–167
methodology 65–66, 140–152
radical stabilization energy (RSE) 176
RAFT agent design 176–179
RAFT equilibrium coefficient 65–66, 87–88, 152–156, 159–164
retardation mechanisms 164, 179
slow fragmentation model 167–171
side reactions in RAFT 156–159
structure-reactivity predictions 167–179
transition state theory
quasi-block copolymers 118, 1054, 1056–1060, 1064
standard 140–147
variational 140, 146
- quinone 612
- r**
- radical stabilization energy (RSE) 176
- radical thiol-ene reaction 760, 775–776, 779
- radical thiol-yne reaction 776–777, 780, 779
- RAFT agent (chain-transfer agent, CTA, ZC=SSR), *see also*
dithiocarbamates, dithioesters, macromonomer RAFT agent, trithiocarbonates, xanthates
chain transfer coefficient (C_{tr}) 96–103, 226–230, 374
dithiocarbamates 554–556, 580
dithioesters 101, 228–229

- trithiocarbonates 375–377
- xanthates 509–510
- R-substituent (homolytic leaving group) 109, 111, 379–382
 - affect on chain transfer coefficient 96–97
 - affect on re-initiation and initialization 105, 107–108
 - for block copolymer synthesis 103–106, 381
 - 2-carboxyprop-2-yl radicals 381
 - for 1,1-disubstituted MAMs 380
 - for IAMs and LAMs 381
 - for monosubstituted MAMs 381
 - stability 109
 - with trithiocarbonates 379–382
- side reactions 109, 114–116, 126–130, 680–681
 - hydrolysis 109, 680–681
- Z-substituent (activating group) 109, 111
 - affect on chain transfer coefficient 113
 - affect on reactivity 116–118
 - affect on reaction kinetics 118–119
 - intermediate radical termination 119–123
 - side reactions, stability 114–116, 126–130
 - slow fragmentation 123–126
- RAFT agent synthesis
 - alkylating agent with carbodithioate 237, 413, 502, 575–577
 - 1,3-dipolar cycloaddition 413, 502
 - of dithiocarbamates 575–580
 - of dithioesters 237–239
 - esterification or amidation 238, 413
 - Markovnikov addition of dithioacid to olefin 237
 - ketoform reaction 413, 580, 581
 - nucleophile with dithiochloroformate or thiocarbonyl-bis-imidazole 413, 577–578
 - radical-induced ester exchange 238, 413
 - radicals with a bis(thioacyl)disulfide 238, 239, 413, 423, 502, 578–580
 - single unit monomer insertion (SUMI) 238, 306–307, 413, 433
 - thionation 238
 - thiol exchange 238, 413, 506
 - of trithiocarbonates 412–413, 433
 - of xanthates 495–502, 506
- RAFT crosslinking polymerization, *see* polymer networks
- RAFT (end-)group removal/transformation
 - dithiocarbamates 588–591
 - cross-coupling with aryl boronic acids 591
 - radical-induced coupling 558, 589
 - radical-induced disproportionation 558, 589
 - radical-induced reduction 589, 590
 - reaction with nucleophiles 589, 590
 - oxidation 591
 - thermolysis 590, 591
 - dithioesters 314–325
 - cross-coupling with aryl boronic acids 324, 325
 - electrocyclic reaction 322, 323, 326
 - oxidation 321, 322
 - radical-induced coupling 318–320
 - radical-induced disproportionation 318–320
 - radical-induced reduction 318, 321
 - reaction with nucleophiles 315–317
 - thermolysis 322
 - hetero Diels Alder 769–772
 - radical-induced coupling 318–320, 442–445, 558, 589, 766–768
 - radical-induced oxidation 768–769
 - radical-induced reduction 318, 321, 445–446, 530–531, 589, 590, 765–766
 - reaction with nucleophiles (reactions of thiol-ends) 315–317, 446–447, 527–528, 589, 681–684, 590, 772–790
 - aminolysis 681–684, 772–790

- RAFT (end-)group removal/
transformation (*contd.*)
- hydrolysis 680–681
 - thermolysis 322, 446, 448–449,
529–530, 590, 591, 763–765
 - trithiocarbonates 442–447
 - oxidation 447, 449
 - radical-induced coupling 442–445
 - radical-induced disproportionation
445
 - radical-induced reduction 445–446
 - reaction with nucleophiles 446–447
 - thermolysis 446, 448–449
 - xanthates 527–531
 - reaction with nucleophiles 527–528
 - oxidation 528–529
 - radical-induced reduction 530–531
 - thermolysis 529–530
- RAFT mechanism transformation
- 307–310, 433, 436–438, 587–588
 - with dithiocarbamates 587–588
 - RAFT-*T*-ATRP 588
 - RAFT-*T*-ROMP 587–588
 - ROP-*T*-RAFT 587
 - ROMP-*T*-RAFT 587–588
 - with dithioesters 307–310
 - ESARA process 309–310
 - RAFT-*T*-ROMP 256, 308
 - ROMP-*T*-RAFT 308
 - RAFT-*T*-ROP 256, 308
 - ROP-*T*-RAFT 308
 - RAFT-*T*-ATRP 308–309
 - RAFT-*T*-NMP 309–310
 - with trithiocarbonates 433, 436–438
 - cationic-*T*-,RAFT 438, 1175–1178
 - RAFT-*T*-anionic 438
 - RAFT-*T*-ATRP 438
 - RAFT-*T*-NMP 438
 - RAFT-*T*-ROAMP 437
 - RAFT-*T*-ROMP 437
 - RAFT-*T*-ROP 436–437
 - ROP-*T*-RAFT 436–437
 - ROMP-*T*-RAFT 437
 - with xanthates 519–521
 - cationic-*T*-,RAFT 519–520,
1175–1178
 - ROP-*T*-RAFT 520
- RAFT PISA, *see* polymerization-induced
self-assemblyRAFT
- polymerization, *see* reversible
addition-fragmentation chain
transfer polymerization
- RAFT-single-unit monomer-insertion
(RAFT-SUMI) 4, 178, 235, 280,
306–307, 325, 373, 379, 413, 420,
425, 433, 436, 625–628, 806–823
- into dithioesters 306–307
 - sequential SUMI
 - styrene-NIPAm dimer 810
 - of styrene 810
 - into trithiocarbonates 433
- rate coefficients
- addition to RAFT agent (k_{add}) 95, 101,
103, 125
 - addition to monomer (k_i) 105, 107
 - initiator decomposition (k_d) 100
 - fragmentation ($k_{\text{-add}}$, k_{add} , k_{frag}) 95,
103, 125
 - propagation (k_p) 96–101, 105, 107, 125
- rate constants, *see* rate coefficients
- γ -ray irradiation 506–507
- reactive groups 716, 722, 736–737, 788,
829–830, 833, 853, 963, 1039, *see*
also alkyne group, azide group,
isocyanate group
- active ester 251–252, 290–291,
404–405, 426, 830–838, *see also*
acyl chloride, acetone oxime
(AO) group, *p*-dialkylsulfonium
phenoxy ester (DASPE),
anhydride group,
N-hydroxysuccinimide (NHS)
ester, nitrophenyl ester,
pentafluorobenzyl (PFB) ester,
pentafluorophenyl (PFP) ester,
salicylic acid ester
- monomer 290–291, 426, 830–838
 - RAFT agent 251–252, 404–405
 - acyl chloride 830, 839

- acryloyl chloride 780–781, 839
- methacryloyl chloride 839
- 4-vinylbenzoyl chloride 839
- alkyl halide 830, 839–840
- alkyne, *see* alkyne group
- aryl boronic acid 325, 392, 591, 791, 830, 848–849, 853
 - cross coupling with aryl bromide group 848
 - cross-coupling with RAFT agent 325, 591, 791
 - (4-vinylphenyl)boronic acid 392, 849
- azlactone group 291, 830, 842, 859, 863
 - functional monomer, *see* 2-vinyl-4,4-dimethylazlactone
 - functional RAFT agent 406
 - polymers 859, 863
- azide group, *see* azide group
- aziridine group 830
- disulfide group 830, 843, *see also* 2-pyridyl disulfide
 - thiol–disulfide exchange 843
- epoxide group 843–844
- isocyanate group, *see* isocyanate group
- redox initiation 4, 47, 109, 312–313, 506–508, 591, 648–656, 672, 684, 695, *see also* eRAFT, photoredox initiation
- ammonium persulfate–sodium
 - formaldehyde sulfoxylate 508, 650, 787
- benzoyl peroxide–amine 507, 648–649
- benzoyl peroxide–*N,N*-dimethylaniline 648–649
- t*-butyl hydroperoxide–ascorbic acid 648, 652
- cumene hydroperoxide–ascorbic acid 648, 652
- iron(II)–hydrogen peroxide (Fenton’s reaction) 648, 652–654
 - bio-Fenton’s reaction 648, 654
- potassium persulfate–sodium
 - thiosulfate 650
- potassium persulfate–sodium ascorbate 650, 652
- redox-RAFT 179, 312–313, 656–660
 - copper activator 656–659
 - dissociative electron transfer (DET)-RAFT 179, 660
 - sodium dithionite activator 660
- redox responsive hydrogels 787–788
- retardation in RAFT polymerization 32, 37–38, 59–60, 66–69, 76–77, 86, 88–91, 107, 116–120, 122–124, 126, 164, 179, 187, 190–191, 193–194, 205, 210, 223, 225–226, 230, 233, 235, 237, 280, 287–288, 313–314, 325, 371, 379–381, 420, 425, 506, 525, 532, 551, 581–584, 935
- thiolactone group reactive polymers, *see* reactive group
- reversible addition-fragmentation chain transfer (RAFT) polymerization
 - overview 1–5
 - terminology 18–24
- reversible complexation-mediated polymerization (RCMP) 708
- reversible crosslinking copolymerization (RCC) 526
- reversible deactivation polymerization (RDP, controlled polymerization)
 - RAFT mechanism transformation 307
- reversible deactivation radical polymerization (RDRP, controlled radical polymerization)
 - RAFT mechanism transformation 307
 - terminology 15–18
- R-group, *see* RAFT agent
- Rhodixan A1, *see* methyl 2-((ethoxy-carbonothioyl)thio)propanoate
- ring closure reactions 522, 949

- ring-opening alkyne metathesis
 - polymerization (ROAMP) 437
 - RAFT mechanism transformation (ROAMP-*T*-RAFT) 437
 - ring-opening metathesis polymerization (ROMP) 308, 437, 587–588, 708, 958–959, 962, 983
 - RAFT mechanism transformation
 - RAFT-*T*-ROMP 256, 308, 437
 - ROMP-*T*-RAFT 308, 437
 - ring-opening polymerization (ROP) 5, 297, 301, 308, 431, 436–438, 515, 525–526, 519, 587, 629, 767, 790, 844, 958–959, 962, 983, 985, 995, 1009
 - RAFT mechanism transformation
 - RAFT-*T*-ROP 436–437
 - ROP-*T*-RAFT 308, 436–437
 - ring-opening copolymerization-induced self-assembly (rROPISA) 725
 - room temperature RAFT polymerization 648, 845
 - rose bengal 624, 719
 - ruthenium photocatalyst, *see* tris(2,2'-bipyridyl)ruthenium(II) chloride ($\text{Ru}(\text{bpy})_3\text{Cl}_2$)
- S**
- salicylic acid (SA) 837
 - Se*-benzyl *O*-(4-methoxyphenyl) carbonodiselenoate 786
 - selective initialization 108, 306–307, 808–809
 - self-assembly of block copolymers 709, 851
 - self-condensing vinyl polymerization (SCVP) 4, 526, 964–966, 986, 1132
 - RAFT inimers
 - dithioesters 301, 305–306
 - enzyme responsive 966
 - trithiocarbonates 433, 434–435
 - semi-batch RAFT polymerization
 - 187, 188, 193–195, 206–208, 313, 440–441, 512, 907, 1077–1078
 - copolymerization 187, 188, 193–195, 206–208, 907
 - mathematical modeling 206–208, 907
 - starved feed emulsion polymerization
 - with dithioesters 313
 - with macromonomer RAFT 1077–1078
 - with trithiocarbonates 440–441
 - with xanthates 441, 512
 - sequence controlled (SC) polymers 179, 626, 628, 805, 816
 - sequence-defined (SD) polymers
 - discrete polymers 806
 - oligo(acrylate)s 810
 - oligomers 816
 - shell-crosslinked (SCL) micelles 846
 - shell-crosslinked nanoparticles 1009
 - shell-functional nanoparticles 712–723, 950, 956
 - silanes 445, 765–766, 863, 1021
 - silica nanoparticles 722, 739, 1021, 1023, 1029, 1033–1034
 - single unit monomer insertion (SUMI), *see* RAFT-single unit monomer insertion (RAFT-SUMI)
 - single-stranded DNA (ssDNA) 701
 - size exclusion chromatography (SEC) 98, 101–102, 119, 201, 205, 513, 516, 590, 693, 695, 700, 726, 774, 781–782, 786–787, 812–813, 937–938, 1055, 1057, *see also* gel permeation chromatography (GPC)
 - slow fragmentation 59, 76–77, 85–86, 119–120, 123–126, 164, 167–171, 187, 203, 649, 669, 817, 935
 - sodium borohydride (NaBH_4) 773, 783, 791, 947
 - sodium styrene-4-sulfonate, *see* 4-vinylbenzenesulfonate sodium salt

- sodium dodecyl sulfate (SDS) 36, 908
- sodium phenyl-2,4,6-trimethylbenzoyl phosphinate (SPTP) 685–686, 695–696
- sodium 4-vinylbenzenesulfonate, *see* 4-vinylbenzenesulfonate
- sodium salt
- solvents (for polymerization)
- acetic acid 615
 - acetone 615
 - acetonitrile (MeCN) 40–41, 46, 51, 52, 615, 996–997
 - anisole 41
 - benzene 39, 52
 - chlorobenzene 52
 - N,N*-dimethylacetamide 40, 46
 - dimethyl carbonate (DMC) 114, 116, 723
 - N,N*-dimethylformamide (DMF) 48, 53, 413, 615, 447, 577, 589, 650, 661–672, 720, 831, 834–835, 859, 913, 992, 996–997
 - dimethyl sulfoxide (DMSO) 46, 413, 442, 577, 619, 620, 622, 624, 630, 631, 659
 - 1,4-dioxane 36, 43, 46, 47, 49, 50, 210, 307, 615, 652, 654, 686, 691, 731, 768, 837
 - ethanol 43, 126, 577, 615
 - ethyl acetate (EtOAc) 578
 - heptane (*n*-heptane) 719, 1059
 - methanol 50, 126, 615, 618
 - N*-methyl-2-pyrrolidone (NMP) 621
 - 2-propanol 445, 765
 - supercritical carbon dioxide (scCO₂) 32, 195
 - tetrahydrofuran (THF) 42, 43, 307, 447, 649
 - toluene 41, 120, 445, 446, 524, 578, 615, 618, 996–997
 - water 32, 36, 37, 42, 48, 126, 513, *see also* aqueous RAFT
 - acetate buffer 45
 - phosphate buffered saline (PBS) 691, 692, 693, 695
- solvophilic macroRAFT agent 711, 712–713, 720, 726, 731
- spherical micelles 518, 697–698, 726, 730, 735–736, 1041
- squaric ester 3,4-diethoxycyclobut-3-ene-1,2-dione
 - in bioconjugation 791
- stable radical-mediated polymerization (SRMP) 755, *see also* nitroxide-mediated radical polymerization (NMP)
- stannanes 765–766
 - by-products 765
- star polymers 523, 950–956
 - applications 1002–1010
 - advanced materials 1007–1010
 - biomedical 1003–1004
 - emulsion stabilization 1006–1007
 - classification of 955–956
 - compact nature 983
 - controlled radical polymerization 985
 - core-crosslinked, *see* arm-first (convergent) synthesis
 - high throughput synthesis 630–631, 1000
 - in-out stars 956
 - mikto-arm stars 951, 953, 956
 - polymer libraries 630–631
 - ring-opening metathesis polymerization (ROMP) 983
 - ring-opening polymerization (ROP) 983
 - synthesis 950–956, 985–1002
 - arm-first (convergent) synthesis 994–1002
 - core-first (divergent) synthesis 630–631, 985–987
 - grafting-to approach 1002
 - R-connected core 988–991
 - Z-connected core 987–988
- star-star coupling 80, 953–954, 987–988, 990–991, 994–995
- starved feed polymerization, *see* semi-batch polymerization

- stimuli-responsive polymer brushes 1027–1029
- stochastic modeling techniques (SMTs) 204–206
 - Monte Carlo 204–206
- strain-promoted alkyne cycloadditions
 - alkyne-functional RAFT agents 407
 - alkyne-nitrile oxide cycloaddition (SPANOC) 848
 - alkyne-nitrone cycloaddition (SPANC) 848
 - azide-alkyne cycloaddition (SPAAC) 407, 522, 634, 847, 848
- styrene (St) 33, 112, 648, 709, 759, 808
 - experimental procedures
 - emulsion polymerization 36–37
 - solution polymerization 33–35
 - RAFT agent selection 112, 113
 - rate coefficients for radical addition (k_i, k_p) 107
 - selective initialization 108, 149–151, 307, 808, 809
 - single unit monomer insertion (SUMI) 108, 808–810
 - with dithiobenzoates
 - kinetics 76, 78, 81, 82, 91, 118–126
 - RAFT agent stability 126
 - with switchable RAFT agent 116–118
 - polystyrene-*block*-poly(vinyl acetate) 117
- styrenic monomers (styrenes, styrenics) 34, 237, 239, 280, 286–287, 307, 413, 416, 421, 422, 557, 572, 575, 627–628, 807, 842, 843, 1171–1172, *see also*
 - 4-chloromethylstyrene,
 - 4-hydroxystyrene, styrene,
 - 4-methylstyrene,
 - 2-vinylpyridine, 4-vinylpyridine
- with dithiocarbamates 552, 558–566, 572, 575
- with dithioesters 234, 280, 286–287
 - aromatic dithioesters 241–245
 - aliphatic dithioesters 264–266
- RAFT agent selection 27–28, 112
 - with trithiocarbonates 380, 381, 421–423
 - symmetric trithiocarbonates 383–386
 - non-symmetric trithiocarbonates 391–401
 - with xanthates 496, 498, 510, 532
- styrene-4-sulfonate sodium salt, *see* 4-vinylbenzenesulfonate sodium salt
- N*-substituted maleimides 817
- N*-substituted oligo(acrylamide)s 822
- sugar moieties and amino acids 719–720
- sulfur-fluoride exchange (SuFEx) chemistry 945
- sulfur-free RAFT polymerization, *see* macromonomer RAFT polymerization
- sulfur-free RAFT agents, *see* macromonomer RAFT agents
- SUMI, *see* single unit monomer insertion
- supercritical carbon dioxide (scCO₂) 32, 187, 195, 211, 721, 898–904
- supramolecular block copolymers 947–948
- supramolecular interactions 734, 737, 933, 947
- surface-initiated photoiniferter-mediated polymerization (SI-PIMP) 554
- surface-initiated RAFT (SI-RAFT) 235, 412, 554, 649, 859, 861, 1019–1021, 1023, 1025, 1029–1031, 1033–1034, 1037, 1039, 1042, *see also* brush architecture
- bioconjugation 1027–1030
- brush architecture 1020–1025
 - bimodal molar mass 1020, 1021
 - block copolymer 1020, 1023
 - Janus 1020, 1022–1024
 - looped 1020, 1022
 - mixed composition (mikto-arm) 1020
 - patchy 1020, 1022
- composites 1030–1039

- stimuli responsive brush 1027–1030
 surface/particle modification 859–864
 surfactant-free latexes 715–716, 740
 switchable RAFT agents 2, 30, 47, 51–52, 116–118, 132, 557, 567–571, 583–585, 587, 591, 593, 764, 809, 935, 1053–1054, 1100, *see also* aryl(pyridin-4-yl)carbamodithioates, cyanomethyl methyl(pyridin-4-yl)carbamodithioate, methyl(pyridin-4-yl)carbamodithioates, methyl 2-((methyl(pyridin-4-yl)carbamothioyl)thio)propanoate, methyl(pyridin-4-yl)carbamodithioates
 symmetrical trithiocarbonates 390, 413, 442, 879
 synthetic metalloporphyrins 620
- t**
- t*-butyl-, *see* under ‘B’
 telechelic polymers 522, 526, 768, 1165, 1172
 discrete decamers 813–814
 hydroxyl copolymers 767
 temperature-induced morphological transitions (TIMT) 735–736
 templating agents 739
 terephthalaldehyde (TDA) 846
tert-butyl-, *see t*-butyl – indexed under ‘b’
 tetrafunctional trithiocarbonate (4-TTC) 993
 5,10,15,20-tetraphenyl-21*H*,23*H*-porphine zinc (ZnTPP) 312, 617, 620, 628–631, 991, 1055
 tetraphenylporphyrin (TPP) 620–621, 667
 thermal initiators 32, 194, 310–311, 439, 506, 647, 684–685, 693, 809–810, 991
 thermolysis 322, 446, 448, 527, 529–530, 590–591, 763–764, 953
 thiazine 613
S-thiocarbamate 781
 2,2'-(thiocarbonylbis(sulfanediy))bis(2-methylpropanoic acid)
 (2-(1-carboxyl-1-methylethyl-sulfanylthiocarbonylsulfanyl)-2-methylpropionic acid,
S,S'-bis(α,α' -dimethyl- α'' -acetic acid)-trithiocarbonate) 36–37, 46, 50, 383, 681, 684–685, 701
 aqueous RAFT polymerization 681, 684–685, 701
 experimental procedures
 acrylamide in solution 46
 N,N-dimethylacrylamide in solution 36
 poly(*N,N*-dimethylacrylamide)-*block*-polystyrene by surfactant-free emulsion polymerization 36–37
 poly[*N*-isopropylacrylamide-*co*-(sodium methacrylate)-*co-N*-cyclohexylacrylamide] in solution 50
 functional derivatives 402–404
 active ester 402
 protected maleimide 402
 yne 403
 polymerizations 383
 thiol
 thiol-disulfide exchange 843
 thiol-ene reaction 719, 775, 841, 844
 thiol-epoxy ring-opening 782–783
 thiol-halo substitution 783–786
 thiol-isocyanate reaction 780–782
 thiol-methanethiosulfonate reaction 317, 774, 789
 thiol-Michael reaction 779, 780
 thiol-*para*-fluorophenyl reaction 786
 thiol-terminated polystyrene 775
 thiol-yne click copolymerization 776
 thiolactones 843
 functional species 761
 -functionalized acrylamide monomers (TLAm) 843
 -functionalized maleimide (TLAMI) 843

- thiolactones (*contd.*)
 -functionalized styrenic monomer 843
 R-group functionalized RAFT agent 761
- thiolates 321, 661, 663, 763, 773, 777, 791
- thiophenol 834, 1165
- thiourethane 781, 840–841
- thioxanthene 613
- thiuram disulfides 578–580, 590, 768, 1089, 1100
- toluene 41, 49, 63–65, 69–76, 81–82, 85, 87–88, 90, 120, 201, 203–204, 445, 615, 618, 765, 777, 859, 906, 908–909, 996–997, 1006
- trans,trans*-2,4-hexadien-1-ol 770
- transfer coefficient, *see* chain transfer coefficient
- transition state theory
 standard 140–147
 variational 140, 146
- triazole-functional trithiocarbonates 759
- 1,2,4-triazoline-3,5-dione (TAD) 771, 772, 845
 triazolinedione Alder-ene reaction 772
 triazolinedione-diene Diels Alder reaction 772
- triblock copolymers, *see* ABA triblock copolymers, ABC triblock copolymers, BAB triblock copolymers
- tributylphosphine (PBu₃) 773, 908
- tributylstannane (Bu₃SnH) 445–446, 531, 590, 765–766
- 2,4,6-trichloro-1,3,5-triazine (TCT) 830, 840
- triethylamine (Et₃N) 238, 621, 783, 834
- triethylene glycol dimethacrylate (TEGDMA) 383, 432, 435, 908
- trimethylaluminium (TMA) 859
- (2,4,6-trimethylbenzoyl) diphenylphosphine oxide (TPO) 685
- trimethylsilyl propargyl methacrylate (TMS-PgMA) 848–849
- triphenylphosphine 739, 768, 850
- tris(2,2'-bipyridyl)ruthenium(II) chloride (Ru(bpy)₃Cl₂) 312, 618–619, 635, 689–691
- tris(2-carboxyethyl)phosphine (TCEP) 773, 781
- tris[2-(dimethylamino)ethyl]amine (Me₆TREN), *see* copper activators
- tris(2-phenylpyridine)iridium(III) (*fac*-Ir(ppy)₃) 312, 506, 612, 617–619, 623, 627, 628, 1176
- tris(2-pyridylmethyl)amine (TMPA), *see* copper activators
- tris(trimethylsilyl)silanes 445, 765
- trithiocarbonates (carbonotrithioates) 72, 74, 75, 77, 86, 130, 194, 226, 237, 371–494, 614–615, 620, 624, 628, 661, 662, 753, 759, 775, 809, 817–820, 823, 935, 936, 1034, 1089, 1101, 1132, *see also* butyl 2-(((dodecylthio)carbonothioyl)thio)-2-methylpropanoate, 4-(((2-carboxyethyl)thio)carbonothioyl)thio)-4-cyanopentanoic acid, 3-(((1-carboxyethyl)thio)carbonothioyl)thio)propanoic acid, 2-(((ethylthio)carbonothioyl)thio)-2-methylpropanoic acid, 2-(((butylthio)carbonothioyl)thio)propanoic acid (BTPA), 4-cyano-4-((phenylcarbonothioyl)thio)pentanoic acid, 2-cyanopropan-2-yl dodecyl carbonotrithioate, dibenzyl carbonotrithioate (DBTTC), 2-(dodecylthiocarbonothioylthio)-2-methylpropionic acid (DDMAT), 2-(((dodecylthio)carbonothioyl)thio)propanoic acid (DoPAT), 2,2'-(thiocarbonylbis(sulfanediyl))bis(2-methylpropanoic acid)
 activating group (Z) selection 380, 382
 ATRP initiator 410, 425, 438, 657

- dibromotrithiocarbonate 657
- bis-trithiocarbonates 382, 387–389
 - R-connected 387–389
 - R" (or Z) -connected 382, 387
- chain transfer coefficients 374–379
- conjugates 390, 416
 - lysozyme 624
- crosslinking polymerization 431–433
 - multiolefinic monomers 432
 - networks 432
 - stars, microgels 432–433
- cyclopolymerization 429–431
- emulsion/miniemulsion/dispersion
 - polymerization 440–441
- functional trithiocarbonates 390, 402–412, 416
 - description 390, 412
 - non-symmetric 404–412
 - symmetric 402–403
- group removal/transformation
 - 442–447
 - by oxidation 447
 - by radical-induced
 - disproportionation 445
 - by radical-induced processes 442–445
 - by radical-induced reduction 445–446
 - by reaction with nucleophiles 446, 447
 - by thermolysis 446
- homolytic leaving group (R) selection
 - 379–386, 391–401
 - for block copolymer synthesis 381
 - for 1,1-disubstituted MAMs 380
 - for IAMs and LAMs 381
 - for monosubstituted MAMs 381
- industrial use 1101–1102, 1132
- monomers, *see also* polymerizations
 - acrylamides 419–421
 - acrylates 418, 421
 - diene monomers 421, 424
 - 1,1-disubstituted monomers 417
 - methacrylamides 417, 420
 - methacrylates 414–416
 - monosubstituted monomers 424–425
 - macromonomers 425, 430
 - with reactive functionality 425, 426–429
 - styrenics 421–423
 - vinyl monomers 424–425
- photoinitiation 438–439
- polymerizations
 - bis-trithiocarbonates
 - R-connected 388–389
 - Z-connected 387
 - non-symmetric trithiocarbonates 391–401
 - functional 404–412
 - symmetric trithiocarbonates 383–386
 - functional 402–403
- publication statistics 372, 390
- ring-opening polymerization 431
- RAFT equilibrium coefficients 72–76, 378
- RAFT mechanism transformation
 - cationic-*T*-,RAFT 438, 1175–1178
 - RAFT-*T*-anionic 438
 - RAFT-*T*-ATRP 438
 - RAFT-*T*-NMP 438
 - RAFT-*T*-ROAMP 437
 - RAFT-*T*-ROMP 437
 - RAFT-*T*-ROP 436–437
 - ROP-*T*-RAFT 436–437
 - ROMP-*T*-RAFT 437
- RAFT single-unit monomer insertion (RAFT SUMI) 433
- RAFT mechanism 72–76, 371–374
 - symmetric trithiocarbonates 373
 - non-symmetric trithiocarbonates 374
- reaction conditions and side-reactions 441–442
- redox-initiated RAFT with 439
- self-condensing vinyl polymerization 433–435
- structural features of 372

- trithiocarbonates (carbonotrithioates) (*contd.*)
 synthesis of 412–413
 triblock copolymer synthesis 382, 387, 389
- U**
- ultra-high molecular weight (UHMW) polymers 511, 654, 992–994
- undecanoic acid 775
- urazole derivative 772
- urazole functional trithiocarbonate (URZ-TTC) 772
- UV or visible light 506–507, 514, 685
- V**
- vinyl acetate (VAc) 47, 105, 107, 114, 117, 156–157, 164, 172, 175–176, 179, 515, 551, 628, 712, 759, 760, 764, 808, 817, 842, 936, 945, 948, 1053
 experimental procedure 47
 rate coefficients for radical addition (k_i , k_p) 107
 with dithioesters 234
- vinyl amides 2, 27–28, 46–48, 371, 572, 753, 806, 855, *see also* *N*-vinylcaprolactam, *N*-vinyl monomers, *N*-vinyl-2-piperidone, *N*-vinylpyrrolidone
 with dithiocarbamates 572
 RAFT agent selection 27–28, 112
 with xanthates 504, 513–514
- vinyl benzoate 48, 242, 498, 512
 experimental procedure 48
- 4-vinylbenzaldehyde 244, 848, 1041
- 4-vinylbenzenesulfonate sodium salt, (styrene-4-sulfonate sodium salt) 244, 247, 248, 261, 262, 280, 679, 882
- 4-vinylbenzoic acid 398, 400, 884
- NHS ester, *see* 2,5-dioxypyrrolidin-1-yl 4-vinylbenzoate
- 4-vinylbenzyl chloride (VBC), *see* 4-chloromethylstyrene
- 4-vinylbenzyl 1*H*-imidazole-4-carbodithioate 209–210, 306
 mathematical modeling 209–210
- N*-vinylcaprolactam 47, 508, 562, 565, 572, 712, 855
 with dithiocarbamates 562, 565, 572
 with xanthates 496, 504, 508
- N*-vinylcarbazole 28, 47, 51, 116, 234, 287, 371, 380, 381, 425, 496, 497, 500, 501, 504, 513, 514, 518, 519, 525, 551, 552, 556, 567, 569, 571, 572, 660, 1053–1054
 with dithiocarbamates 551, 552, 556, 567, 569, 571, 572
 with dithioesters 234, 249, 287
 experimental procedure 47
 with xanthates 496, 497, 500, 501, 504, 513, 514, 518, 519, 525
- vinyl chloride (VC) 151, 179, 498, 505, 515, 518
 with dithiocarbamates 179, 565, 572
 with xanthates 498, 505, 515, 518
- 2-vinyl-4,4-dimethylazlactone 46, 244, 291, 824, 842
 copolymer 244, 842
- vinyl esters 2, 27–28, 46–48, 371, 503, 506, 510, 512, 533, 551, 572, 585, 753, 806, 950, *see also* vinyl acetate, vinyl benzoate
 RAFT agent selection 27–28, 112
 with dithiocarbamates 572
 with xanthates 503, 510, 512
- vinyl ethers 936, 1160–1179
 2-chloroethyl vinyl ether (CEVE) 1170, 1171
 dodecyl vinyl ether 775
 ethyl vinyl ether 1170, 1171, 1174
 2-ethoxyethyl vinyl ether (EOEVE) 1170, 1171, 1175
 isobutyl vinyl ether 1160, 1164, 1170, 1171, 1174
 2-methoxyethyl vinyl ether 1169
- vinyl halides, *see* halo-olefins, vinyl chloride

- vinylidene chloride (VCl₂) 245, 248, 249
- vinylidene fluoride (VF₂) 496, 501, 505, 509, 514, 518, 520, 521, 533, 566
- with dithiocarbamates 566, 572
- with xanthates 496, 501, 505, 509, 514, 518, 520, 521, 533
- vinyl imides, *see* *N*-vinylphthalimide 47
- vinyl monomers
- N*-vinyl monomers, *see also* vinyl amides, *N*-vinylcarbazole, *N*-vinylcaprolactam, vinyl imides, *N*-vinylphthalimide, *N*-vinyl-2-piperidone, *N*-vinylpyrrolidone
- with dithiocarbamates 572, 576
- with xanthates 504–505, 513–514
- O*-vinyl monomers, *see* vinyl esters, vinyl ethers
- P*-vinyl monomers, *see* vinyl phosphonic acid
- S*-vinyl monomers 504, 512–513
- Si*-vinyl monomers, *see* vinyl triethoxysilane
- vinyl phosphonic acid (VPA) 510, 513, 518, 533
- N*-vinylphthalimide 48, 248, 400, 505, 514, 559, 572, 809
- experimental procedure 48
- with dithiocarbamates 559, 572
- N*-vinyl-2-piperidone 496, 504, 505, 514, 855
- N*-vinylpyrrolidone 28, 46–48, 116, 234, 241, 244, 248, 287, 288, 380, 506, 508–510, 513–515, 517, 519, 522, 525, 527, 532, 552, 559, 565, 567, 568, 569, 571, 572, 575, 583, 587, 591, 652, 693, 1053
- experimental procedure 48
- with dithiocarbamates 552, 559, 565, 567–569, 571, 572, 575, 583, 587, 591
- with dithioesters 234, 241, 244, 248, 287
- with xanthates 506, 508–510, 513–515, 517, 519, 522, 525, 527, 532
- vinyl triethoxysilane (VTES) 722
- vinylbenzoyl chloride (VBzC) 839
- 2-vinylpyridine 241, 247, 253, 258, 261, 306, 316, 538, 572, 583, 615, 649
- 4-vinylpyridine 241, 245, 247, 249, 253, 306, 572, 722
- Vitamin B12 release profiles 905
- vitamin B2 (riboflavin)- riboflavin 5'-mononucleotide (FMN) 690
- vitamin C, *see* ascorbic acid 695
- W**
- water soluble macroRAFT agent 680, 686, 691, 696
- water soluble RAFT agent 680
- water-soluble methacrylamide polymers 838
- water-soluble trithiocarbonate 650, 652, 654
- wavelength orthogonal polymerization 628–629
- wormlike micelles 697
- X**
- xanthates (dithiocarbonates) 48, 69, 72, 112–114, 194, 198, 495, 509–510, 519, 522, 526, 650, 662, 753, 759, 936, 948, 1164–1165, 1175–1176, *see also* 2-((ethoxycarbonothioyl)thio)acetic acid, ethyl 2-((phenoxycarbonothioyl)thio)propanoate, 2-hydroxyethyl 2-((ethoxycarbonothioyl)thio)-2-methylpropanoate, methyl 2-((ethoxycarbonothioyl)thio)propanoate
- canonical forms 114
- cationic RAFT polymerization 1164–1165, 1175–1176
- end-group removal 527–531
- nucleophilic reaction 527–528
- oxidation 528–529
- radical-induced reduction 530–531

- xanthates (dithiocarbonates) (*contd.*)
- thermolysis 529–530
 - industrial applications 531–532
 - initiation 506–508
 - kinetics 509–510
 - macromolecular architectures 516–526
 - block copolymers 518–521
 - cyclic polymers 521–522
 - end-functional polymers 517–518
 - gradient copolymers 521
 - graft copolymers 522–523
 - hyperbranched polymers 526
 - star polymers 523–526
 - mathematical modeling 194, 198
 - monomers 510–516
 - acrylates and acrylamides 510–511
 - cyclic ketene acetals 515–516
 - diallyl monomers 516
 - ethylene 515
 - methacrylates 511–512
 - styrenics 510
 - vinyl esters 512
 - N*-vinyl monomers 513–514
 - S*-vinyl monomers 512–513
 - vinyl phosphonic acid 513
 - polymerization conditions 508–509
 - heterogeneous polymerization 508–509
 - high pressure polymerization 508
 - RAFT mechanism transformation
 - cationic-*T*-,RAFT 1175–1178
 - side reactions 114–116
 - synthesis of 495, 496, 502
 - xanthene dyes 623–625, 635
 - xanthogen disulfides 502
 - xanthone 613
 - xyloglucan (XG) 719–720
- y**
- yne, *see also* alkyne
- ω -yne end-functional polymer 779
 - poly(*N*-isopropylacrylamide) 779
 - yne-functional monomer, *see*
 - alkyne-functional monomer, propargyl acrylate, propargyl methacrylate 288–289, 426
 - yne-functional trithiocarbonate 759
 - 2-cyano-5-oxo-5-(prop-2-yn-1-ylamino)pentan-2-yl-dodecyl-trithiocarbonate 760
 - yne-hydroxyl-functional trithiocarbonate 780
- yttrium(III) trifluoromethanesulfonate (Y(OTf)₃) 618
- z**
- Z group, *see* RAFT agent
- Z-connected bis-RAFT agents 879
 - zinc-iron oxide semiconductor (Zn_{0.64}Fe_{2.36}O₄) 616
 - zinc(II) *meso*-tetra(*N*-methyl-4-pyridyl) porphine tetrachloride (ZnTPMPyP) 629, 691
 - zinc(II) tetraphenyl porphyrin (ZnTPP) 312, 617, 620, 628–630, 991, 1055
 - zinc(II) *meso*-tetra(4-sulfonatophenyl) porphyrin (ZnTPPS⁴⁻) 621, 689, 691
 - zinc(II) (2,3,7,8,12,13,17,18-octaethyl-5,10,15,20-tetraphenylporphyrin) (ZnOETPP) 621–622
 - zwitterionic pH-responsive copolymers 843