

Index

a

- accessible intraparticle porosity 77
- acrylate-based resin, pore size
 - distributions of 128
- adsorbed phase diffusion 184, 193, 218, 222
- adsorption
 - azeotropes 256–257
 - definition 159
 - equilibrium 69, 78, 89, 159–180, 219, 231, 237, 244, 251, 255, 256, 258–259, 265, 275, 289, 296, 304, 342, 371, 381–382
 - isotherm 161–165, 167, 174, 177–179, 204, 206, 212, 216–219, 221, 233–234, 236, 241–243, 246, 254–255, 259, 278, 294, 301, 313, 320, 338, 366, 381
- adsorptive membranes 119, 121, 133, 268, 358
- adventitious agents 18, 44, 49, 51, 355, 358
- affinity purification systems 111
- agarose polymers 122–123
- aggregates 6, 37, 40–41, 48, 51, 95–96, 169, 355, 358, 364
- air sensors 156
- AKTA Pure systems
 - dynamic mixer used in 145
 - pump head in 143
 - tandem piston pumps 144
- alcohol oxidase promoter (AOX)
 - expression system 45
- amino acids 3, 5
 - naturally found 3, 4
 - side chains 5
 - structure 3
- amino residues 107
- ammonium sulfate 10, 32–33, 66, 99, 100, 163, 179, 333–337, 358
- amphipathic α -helices 7, 11
- ampholytes 345–346
- anion exchangers 15, 28, 103–104, 106, 118, 172, 250–251, 261, 312, 340–341, 345–346, 349, 351
- antibodies 1–2, 7, 11–12, 26–27, 29–30, 41–43, 46–47, 49, 52–53, 55, 56, 59–60, 66, 72, 74, 96, 100–101, 103, 109–114, 118, 124, 166–167, 173, 177–179, 183, 190, 195, 202, 214, 248–251, 336–340, 343, 345, 359–361, 370, 405–407
- arginine 118
- artificial chaperones 58
- asymmetry factor 158, 270, 271, 273, 275
- attenuated total reflectance Fourier transform infrared spectroscopy (ATR-FTIR) 180
 - conformational changes of protein 3D-structure 9
- axial diffusion 78–79, 82–83, 228, 286
- axial dispersion 82, 201, 227–233, 237, 243, 250, 265, 276, 278, 283–285, 288, 290, 291, 303, 314, 338, 386, 388
- axial dispersion effects 78–79
- axially-dispersed plug flow model 227–228, 278

b

- bacterial immunoglobulin-binding
 - proteins 110, 113
- band broadening factors 78, 265, 381, 386
- batch adsorption
 - advantage 202
 - binding kinetics control 210
 - in an infinite bath 211
 - laboratory studies 203
 - LDF rate equation 210–211
 - of 1 mg/ml α -chymotrypsinogen on SP-Sepharose-FF 212
 - rate equations 204
 - with external mass transfer control 207
 - with pore diffusion control 208–210
 - with solid diffusion control 207–208
- beads 54–55, 95, 119, 121–123, 126, 129–131, 157
- BET isotherm 168
- bi-Langmuir isotherm 166–167
- bind-wash-elute sequence 312
- biological products 1, 47–48, 54
- biopharmaceutical manufacturing 42, 54, 273, 357, 359
 - environmental impact 59–60
- biopharmaceutical process development
 - purity requirements 51–52
 - quality by design (QbD) 50
 - validation of 50–51
- biopharmaceuticals 1
 - downstream processes 355
 - market share 1, 2
 - product quality requirements 46
- bioprocessing operations
 - shear rates in 35
- biosimilars 52–53
- biospecific interaction chromatography (BIC) 66, 93, 94
 - advantage 109
 - fusion protein concept 110
 - steps in 108
- biospecific ligands classification 110
- Biot number 210, 213
- blood plasma fractionation 1
- boundary layer mass transfer coefficient 185
- bubble traps 78, 149, 156–157
- buffers 18, 58, 60, 101, 115, 117–120, 140–141, 145–146, 250, 253, 255–256, 341–346, 349, 356, 372–374

c

- capacity utilization (CU) 73, 356, 365–369
- capture downstream process 357
 - adsorbed contaminants 361
 - CIP step 364
 - compressible media 368, 369
 - continuous capture 370–375
 - effluent profiles 362
 - elution step 364
 - feed load 361
 - load steps 362–363
 - productivity and capacity utilization 365–369
 - re-equilibration step 365
 - rigid particles 365, 366
 - wash step 363–364
- carboxyhemoglobin, solubility of 32
- Carman–Kozeny equation 133, 197–198, 366, 369
- cation exchangers 27–28, 72, 103–104, 106, 117, 129, 169, 171–173, 176–178, 191–192, 195, 221, 250–251, 259, 262, 312, 340–341, 346–347, 351
- cell disruption 359
- ceramic hydroxyapatite
 - amino acids, peptides and proteins, retention of 117
 - SEM and TEM of 116
- ceramic hydroxyapatite (CHT) 115–117
- chaotropic salts 34, 359
- chemical stability, of proteins 33
- chromatofocusing 345, 349
- chromatographic capture 71, 367
- chromatographic columns 53
 - efficiency 81–84
 - flow non-uniformity 78, 79

- fluid flow 77
- laminar flow conditions 77
- length requirements 86
- packed 77
- scaling 88–90
- separation performance 78, 79
- velocities in 139
- chromatographic methods 68, 161
- chromatographic process
 - capture 357
 - volume reduction 357
- chromatographic resolution 84
 - adsorption equilibrium 382
 - band-broadening effects 381
 - feed volume 376
 - isocratic elution 378
 - linear isotherm and isocratic elution 379–382
 - pulse injections 376
- chromatographic separation 1, 14, 50, 64, 73–74, 78–79, 84, 89, 204, 270, 339, 350, 358
- chromatographic systems 140
 - specially designed workstations 139
 - superficial mobile phase velocities and column lengths 140
- chromatographic velocity 232, 337, 344
- chromatography 63, 67
 - defined 63
 - displacement 71–72
 - elution 69–70
 - frontal analysis 70–71
 - modern liquid chromatography, branches of 66, 67
 - performance factors 76–80
 - scaling relationships 87–90
- chromatography columns
 - axially-dispersed plug flow model 227
 - displacement chromatography 255–263
 - disposable 150, 151
 - local equilibrium theory
 - chromatographic velocity 232
 - coupling of flow and adsorption equilibrium 231
 - frontal loading of column 234–236, 241–243
 - isocratic elution, preparative injection 239–241
 - linear isocratic elution 233–234
 - mobile and stationary phases 231
 - protein breakthrough curves 243
 - saturated elution of column, Langmuir isotherm 236–239
- modes of operation, entrance conditions
 - dimensionless equations 230–231
 - displacement chromatography 229
 - elution chromatography 229
 - frontal analysis 229
- radial flow 153–155
- multicomponent systems
 - breakthrough profiles 248–250
 - characteristic velocities 247
 - characteristic velocity 245
 - coherent paths 245, 246, 248
 - column dynamic behavior 252
 - frontal analysis behavior, for binary mixture 246–248
 - Na⁺ adsorption isotherms and outlet profiles 254
 - outlet concentration profiles 248
 - pH and sodium concentration profiles 256
 - pH transitions 250–252
 - protein ion exchange chromatography 250
 - quaternary ammonium ion (Q) functional groups 250
 - theory of coherence 244
- chromatography media 94
 - base matrices 119, 121
 - buffers and mobile phases 117–118
 - categories 118, 121
 - electrostatic interaction 103–106
 - hydrophobic interaction 96–103
 - mixed mode interaction 113–117
 - particle size and particle size distribution 131

- chromatography media (*contd.*)
- porosity, pore size and surface area 125
 - steric interaction 94–96
- clean-in-place (CIP) protocols 122–123, 125, 361–362, 364–365, 372
- coherence, theory of 244
- colloidal models 174
- column chromatography 64, 65, 67, 139, 140
- column efficiency 265–275
- plate model 275–278
 - competitive adsorption systems 303–308
 - extracolumn contributions 268
 - flow and thermodynamics 266
 - HETP 265–267, 272–275
 - linear isotherm 266
 - rate models
 - with linear isotherms 278–287
 - with non-linear isotherms 287–303
- combined mass transfer resistances, analytical solutions for 211
- commercial simulators 338
- competitive adsorption systems 178, 219, 287, 303–308
- compression factor 157, 318, 321, 329, 332–333
- conductivity monitors 147, 149, 267
- confocal scanning laser microscopy (CLSM) 215, 217, 222
- conformational stability, of a protein 33
- constant-separation factor isotherm 166, 244
- contaminants 2–40, 48, 49, 53, 101, 238, 248, 355, 357, 361–363, 365–366
- continuous annular chromatography (CAC) 57, 64, 65
- continuous bed structures 76
- continuous stationary phases 119
- controlled porous glass (CPG) 124, 125
- cosmeceuticals 2
- countercurrent adsorption process, for binary separations 75
- critical impurities 48
- critical operational parameters 50
- critical velocity 134–135, 368
- culture broth, components of 45
- culture media 18, 44, 45
- cysteine 5, 6, 108
- d**
- deamidation 13, 15, 16, 26, 355
- Debye length 104, 160, 174, 191
- Debye–Hückel theory 32, 98, 342
- design space 50, 51
- diaphragm dampeners 143
- differential migration process 64
- diffusional hindrance coefficient 188–190, 192, 195, 301
- diffusivity
 - of non-globular proteins 39–40
 - of proteins 38
 - dispersion and rate process, on column performanceband broadening factors 265
 - empirical characterization, of column efficiency 270, 271
 - extracolumn contributions 268
 - flow and thermodynamics 266
 - HETP 265–267, 272–275
 - linear isotherm 266
 - symmetrical peaks 268
 - plate model 275–278
 - rate models
 - competitive adsorption systems 303–308
 - with linear isotherms 278–287
 - with non-linear isotherms 287–303
- dispersion-repulsion forces 159
- dispersive factors 69, 78–79, 81–83, 87, 243, 275, 289–290, 338
- displacement chromatography 71
- adsorption azeotropes 256
 - feed components 255
 - isotachic train 255, 257–262
 - transient profiles 262–263

- disulfide bond formation 6
- DNA 16, 17
 purity 22
 specific mass extinction coefficient 22
- downstream processing 1, 355, 360
 chromatography in 53–59
 flow sheet for proteins purification 55
 post-translational modifications 12–16
 solubility 31–33
- dynamic binding capacity (DBC) 86–89, 287–288, 291–296, 301–303, 361–369
- dynamic light scattering (DLS) 24, 40
- e**
- E. coli*, protein expression in 41
- economic viability 355
- effective pore diffusivities 188, 190–194, 212–213, 219, 222, 285–287, 296, 298, 300–302, 306, 320, 325, 363, 365–366, 373
- electron microscopy methods 129
- electrostatic forces 159–160
- Elelyso (taliglucerasealfa) 42
- elution chromatography 63, 67–71, 76, 84, 100, 229, 235, 279, 311–351, 391
- endotoxin units (EU) 18, 46
- endotoxin-free water 18
- endotoxins 18
 chemical structure 18, 19
 coagulation test 20
 concentrations 20
 detection 18
- engineered Protein A 110
- environmental scanning microscopy (ESM) technique 129
- equilibrium binding capacity (EBC) 87, 89, 287–288, 291, 293–296, 302–303, 363, 365–369
- equilibrium factors 78, 81
- exponentially modified Gaussian (EMG) function 271–274
- expression systems, for recombinant proteins 40
E. coli BL21 41
 mammalian expression system 42
 plants and plant cells 42
 transgenic mammals 42
- external mass transfer coefficient 187
- external mass transfer resistance 185, 325
- extraction systems
 liquid–liquid 53
- extraparticle porosities 76–77, 125, 127, 133, 228, 243, 365, 368–369
- f**
- favorable adsorption isotherm 234
- film model 184
- flow behavior index 36
- flow consistency index 36
- flow-through process 86
- fluorescence spectroscopy
 of amino acids 11
- free surface model 197–198
- Freundlich isotherm 167
- frontal analysis 68, 70–72, 74, 161, 206, 229, 244–246, 248–251, 275, 304, 307–308
- fusion protein concept 110
- g**
- gas chromatography (GC) 83
- Gaussian distribution 277
- gel filtration chromatography 93
- gel permeation chromatography 93
- generalized van Deemter curves 81, 82
- generalized van Deemter equation 83
- Gibbsian dividing surface 159
- globular proteins
 diffusivity of 39
- globular proteins, size of 22
- glycosylation 13–15
- gradient elution chromatography 69, 70
 with pH Gradients 340–351
 “bind-wash-elute” sequence 312
 at high protein loads 337–340
 LGE relationships

- gradient elution chromatography
(*contd.*)
- conditions, for minimize 331–332
 - experimental data 325–327
 - for RPC and HIC 332–337
 - HETP experiments 323–325
 - mobile phase modulator 320
 - protein retention factor 320
 - retention factor and mass action law parameters 322–323
 - salt concentration at elution, determination of 327–329
 - linear gradients 312
 - mobile phase composition 312, 313
 - mobile phase modulator 311
 - pH gradients 312
 - reversed phase chromatography (RPC) 311
 - with linear isotherms
 - axial dispersion and flow heterogeneities 314
 - characteristic velocities 314
 - equilibrium parameters 317
 - isocratic elution experiments 317
 - linear driving force (LDF) approximation 319
 - linear gradient elution 313
 - peak compression 314
 - protein characteristic velocity 317
 - retention factors 317
 - gradient mixers 145
 - gradient SMB 76
- h**
- Hamaker constant 174
 - height equivalent to a theoretical plate (HETP) 81, 265–267, 272, 273
 - contributions 82
 - estimation 83–84
 - preparative/process protein chromatography 83
 - reduced 81, 83
 - α -helices 6–7, 11
 - hemoglobin 3, 11, 21, 243
 - Henry's law 69, 85, 241
 - high-pressure mixing 145, 146
 - His residues 107, 108
 - His-tags 108–109
 - histones 17
 - Hofmeister series 33, 99
 - homogeneous diffusion 193
 - host cells composition 43
 - human serum albumin 27
 - positive and negative charge distribution 29
 - human von Willebrand factor 2
 - hydrodynamic dispersion 78–79, 82–84, 214, 228, 286
 - hydrogen-deuterium exchange mass spectrometry (HDMS) 335
 - hydrophilic α -helices 7
 - hydrophilic interaction liquid chromatography (HILIC) 93
 - hydrophobic α -helices 7
 - hydrophobic chromatography 66, 96
 - hydrophobic interaction chromatography (HIC) 29, 58, 93, 94, 96, 311
 - ammonium sulfate 99
 - characteristics of 97
 - elution 101
 - isocratic elution 96, 98
 - labile proteins and virus purification 102
 - ligands used in 96, 97
 - limitations 100
 - polishing of antibodies 103
 - protein retention vs. ionic strength 100
 - protein retention vs. salt concentration 96, 98
 - retention behavior 98
 - vs. RPC 100, 101
 - hydrophobicity of proteins 28
 - hydroxyapatite 115, 117
 - column, pH transients in 117, 118
 - retention characteristics of biomolecules 116
 - hydroxyapatite crystal functional groups 116
- i**
- ideal adsorbed solution theory (IAS) 338

- iminodiacetic acid (IDA) 106–109
 immobilized folding catalysts 58
 immobilized metal affinity
 chromatography (IMAC) 58,
 106, 108
 performance of 109
 immunoglobulin-binding bacterial
 proteins 113
 impingers 146
 induced pH gradients 346–351
 inorganic materials 123, 124
 insulin 1, 2, 41
 interstitial velocity 77, 81, 86, 139, 140,
 231
 intraparticle convection 77–78,
 196–201, 287, 293
 intraparticle Peclet number 199–201
 intraparticle porosities 76, 77,
 125–126, 188, 323, 332, 363, 366,
 379
 intrinsic viscosity
 defined 36
 of biologically important
 macromolecules 37, 38
 inverse size exclusion chromatography
 (iSEC) 126–127, 129–130
 ion exchange chromatography (IEC)
 93, 94, 103, 311
 advantages 104
 ligands used 103
 separation power 104
 stoichiometric displacement model
 104
 ion exchange column 106, 251–255,
 296, 358
 ion-exchange chromatography (IEC)
 66, 72, 93, 94, 103, 104, 113, 118,
 162, 180, 311–312, 314, 320–333,
 340–341, 358
 iso-resolution Curve 320–332
 isocratic elution 85
 isocratic elution 69–70, 72–73, 76,
 78–79, 85, 87, 96, 100, 140, 161,
 231–235, 239–241, 283, 314,
 317–319, 325, 336, 338,
 377–382
 isoelectric focusing (IEF) gel separation,
 of recombinant human
 erythropoietin 14
 isoelectric point 14, 15, 26–28, 31, 104
 isopropyl- β -D-thiogalactopyranoside
 (IPTG) 44
 isotachic pattern 71
 isotachic train 71
- j**
- Jones–Dole B-coefficient 33
- k**
- kinetic stability, of proteins 34
 kinetics of protein binding 201
 Kohlrausch law 147
 kosmotropic salts 34, 94, 99, 163
- l**
- laboratory scale chromatographic
 systems
 components 140, 141
 corrections for extracolumn effects
 149, 150
 mixers 145–146
 monitors 146–149
 pump types 141–144
 α -lactalbumin 12, 35, 57, 100
 CD-spectrum of 9
 β -lactoglobulin, solubility of 31
 Lambert–Beer law 21, 146
 Langmuir isotherm 163, 164, 166–168,
 174–179, 202, 207, 209–210,
 234–240, 246–248, 258–261,
 263, 288, 291, 303–304, 306, 338,
 371–373, 381
 Langmuir model 165, 167, 172, 176,
 206, 298, 338
 Langmuir parameters 165
 Langmuir–Freundlich isotherm
 167–168, 179, 241243
 Laplace transform 284
 limulus amoebocyte lysate (LAL) test
 18
 linear driving force (LDF) 278
 approximation 205, 319
 rate coefficient 205

- linear gradient elution (LGE) 161, 313, 315, 317, 320–337
 - linear isotherms 85, 170, 172–174, 177, 206, 208, 211, 231–233, 235, 256, 269–270, 277, 278–287, 289, 312–320, 333, 339–340, 359, 363, 376–377, 379–382, 388–391
 - liquid–liquid extraction systems 53
 - low-pressure mixing 145
 - lyotropic series 33
 - lysozyme
 - net charge of 26, 27
 - positive and negative charge distribution 29
 - surface hydrophobicity 30
- m**
- machine learning methods 178
 - macropore diffusion 188
 - MagneVist 80
 - mammalian cell system 41–46, 49, 359
 - Mark–Houwink equation 37
 - mass spectrometry 25–26, 146, 180, 335
 - mass transfer effects 78, 82, 161, 262, 266, 386
 - mass transfer enhancement 200
 - matrix-assisted refolding 57–58
 - mechanical stability, of proteins 35
 - melting temperature 34
 - mercury intrusion 126–128
 - metal chelate affinity chromatography 106
 - metal chelate chromatography 93
 - metal interaction chromatography (MIC) 66, 93, 94, 106, 108
 - metal ion interaction 106–108
 - methanol 45, 96
 - microheterogeneity 355
 - micropore diffusion 193
 - mixed mode chromatography (MMC) 66, 93–94, 113, 115
 - mobile phase 12, 63, 93, 139, 161, 183, 227, 266, 311, 358
 - mobile phase modifier 69, 101, 107
 - mobile phase modulator 66, 68, 70, 311, 316, 320, 358, 361
 - modifier 66, 69, 96, 100–101, 107, 169, 332–333, 382
 - modulator 66, 68–70, 229, 279, 311–316, 318, 320, 332–333, 338, 358, 361
 - molal surface tension increments 98, 99
 - molecular diffusion coefficient 38
 - monoliths 77, 119, 121, 126, 129–131, 133, 140, 153, 155, 183, 201–202, 214, 268
 - moving bed system 65
 - multicolumn countercurrent solvent gradient purification process (MCSGP) 76
 - multicomponent adsorption kinetics 218, 223
 - multimodal chromatography (MMC) 66, 93, 94, 115
 - multimodal interaction 113, 115
- n**
- natural carbohydrate polymers 121–122
 - nitrilotriacetic acid (NTA) 107–109
 - nitrogen adsorption 126–127
 - non-critical impurity 48
 - non-proteinaceous biomolecules 1
 - normal phase chromatography (NPC) 93
 - nucleic acids 21–22, 43–44, 359
 - UV absorbance 21
 - nucleotides 16, 45
 - in DNA 17
- o**
- oxygen transport proteins 45
- p**
- packed chromatographic columns 77
 - particle conservation equations 204, 206
 - particle size distribution (PSD) 131–132, 204, 206, 213, 299
 - PCC system 371, 373–375
 - peak compression 314, 317–318, 320–322, 325, 329, 332–333

- PEGylated proteins 40
- peptide bond formation 3
- percolation processes 63
- perfusion 42, 78, 140, 196, 198, 370
- perfusion chromatography 140
- periodic countercurrent process 73
- periodic countercurrent strategy 371
- peristaltic pumps 143, 156
- pH 5, 66, 93, 141, 162, 212, 250, 272, 312, 359
- pH monitors 141, 149
- plasmid DNA 16, 17, 36
- plasmids 1, 17, 22, 38, 40, 116, 130
- plate model 275–278, 321, 326–327
- plate number 81, 86–88, 266–267, 279, 321–322, 358
- Poisson distribution 276
- polishing, downstream process 358
- polynucleotides 16–17
- pore diffusion 81–82, 183–184, 188–193, 195, 199–200, 202, 208–217, 219, 222, 266–267, 269–270, 279, 283–285, 294, 297, 299–303, 307, 319, 363–364, 366, 371, 373
- pore size
 - for SP-Sepharose-FF 129
 - of Protein A adsorbents 129, 130
- porous glass 124
- post-translational modifications, of proteins 12–15
- pressurized continuous annular chromatography (P-CAC) 57
- primary recovery 54
- Process Analytical Technologies (PAT) 51
- Process Mass Intensity (PMI) 59, 60
- process robustness 355
- process scale chromatography system
 - flow diagram 155
 - gas bubbles 156
 - peristaltic pumps 156
- rotary lobe pumps 156
- process scale columns 151
 - design 150
 - large-scale 150
 - packing 157, 158
- process-related impurities 48
- process-scale columns
 - pre-packed 152
 - pre-packed, disposable 153
 - single use, pre-packed 152
- product-related impurities 48, 357–359
- proline 5
- Protein A affinity chromatography 103, 117–118
- Protein A media 110, 111, 114
 - protein adsorption 195
 - rate mechanisms 183–202
 - in spherical adsorbent particles 205
 - transport and kinetic resistances 184
- protein adsorption equilibrium
 - attractive/repulsive interactions 160
 - charged and hydrophobic groups 159
 - chromatographic methods 161
 - diffusional resistances 160
 - empirical correlation of data 178
 - empirical/semi-empirical approaches 160
 - frontal analysis experiments 161
 - ionic strength 160
 - multicomponent systems
 - Langmuir isotherm 174
 - native and deamidated antibody isoforms 178
 - multiple binding sites/ligands 159
- protein conformational changes 180
- robotic operated equipment 160
- single component systems
 - bi-Langmuir isotherm 166
 - bi-Langmuir model 167
 - colloidal models 174
 - constant-separation factor isotherm 166
 - deamidated isoform 173
 - empirical isotherm models 167
 - IEC, HIC, and BIC 162
 - ion-exchange and hydrophobic adsorption 162
 - isotherm data 171, 172

- protein adsorption equilibrium (*contd.*)
- Langmuir isotherm model 163, 167
 - low-loading behavior 172
 - mobile phase composition 164
 - pH values 163
 - protein concentration and free protein 161
 - protein retention and salt concentration 170
 - protein-counterion exchange 169
 - steric mass action model (SMA) 170
 - with Langmuir parameters 165
- protein mass transfer data examples 194
- protein refolding, reaction scheme for 56
- proteins
- attenuated total reflectance Fourier transform infrared spectroscopy 9
 - biochemical and biophysical properties 20–40
 - CD-spectroscopy 7, 9
 - charges 26–28
 - chemical stability 33–34
 - in cosmetics 1
 - description 2–3
 - diffusivity of 38–40
 - folding process 12
 - α -helices in 6
 - hydrophobicity of 28–31
 - ionization methods 25
 - isoelectric points 26, 28
 - loops in 7
 - mechanical stability of 34–35
 - oligonucleotides and polynucleotides 16–17
 - post-translational modifications of 12–16
 - primary structure 3–11
 - quaternary structure 3, 11–12
 - secondary structure 3
 - shape of 40
 - β -sheets in 6–8
 - size 22–26
 - solubility 31–33
 - specific absorbance of 22
 - tertiary structure 3, 11
 - UV absorbance 20–22
 - viscosity of 35–38
 - with quaternary structure 11
 - secondary structure 6
- pulsation dampeners 143
- pyrogens 18
- q**
- quality by design (QbD) 50
- r**
- radial flow chromatography columns 153–155
- rate factors 78, 231
- recombinant antibody
- purification of 55, 56
 - surface hydrophobicity 30
- recombinant human erythropoietin (rhEPO), isoelectric focusing of 14
- recombinant Protein A 41, 110
- recombinant proteins
- host cells composition 43
 - single cells composition 43
- reduced HETP 81–83, 158, 266, 325
- refolding yield of protein 58
- reproducibility 101, 273, 345, 355
- retention factor 85–86, 96, 99–101, 230, 279, 281, 314, 317–320, 322–323, 326–327, 329, 332–333, 336–337, 363, 376, 379
- reversed phase chromatography (RPC) 93, 311, 332
- characteristics of 97
 - elution 101
 - vs. HIC 100, 101
 - ligands used in 96, 97
 - mobile phase modifier, qualitative effect of 101
 - retention 93
 - retention factor 101
- retention mechanisms 100
- Reynolds number 77, 188, 286
- rhizo-secretion 42

- RNA 16, 17, 21, 22, 44
 molecules, classes of 17
- S**
- Saccharomyces cerevisiae* 41
- salt tolerance 115
- scanning electron microscopy (SEM)
 115, 129, 130
- SDS-polyacrylamide gel electrophoresis
 (SDS-PAGE) 5, 25, 41,
 102–103, 105, 109
- sensitivity coefficient 99, 101, 332–333
- separation factor 166, 206, 244, 263,
 288, 291–292
- separation performance metrics
 81–90
- septic shock 18
- shallow bed adsorption systems 203,
 204
- β -sheets 6–8, 11
- Sherwood number 185–186
- shock layer thickness 303, 306–308
- short-range repulsive forces 159
- shrinking core model 210, 217,
 221–222, 294, 307
- Simpson rule 243
- simulated moving bed (SMB) process
 57, 65, 66, 73
 advantages 76
 design 382–391
 principle of operation 74
- single piston syringe pump 142
- size exclusion chromatography (SEC)
 23–25, 93, 94, 358
 advantage 96
 porous chromatography particles
 94, 95
 promoted refolding 57
 purposes of 95
 separation in 93
- solid core particles 125
- solid diffusion 192–196, 205, 207–208,
 210–211, 214–215, 218–219,
 293, 299
- solubility
 of ammonium sulfate solutions 33
 of carboxyhemoglobin 32
 of β -lactoglobulin 31
 of proteins 31
- solvophobic theory 96, 98, 100, 333,
 337
- Sorbex process 74
- SP-Sepharose-FF
 effective pore diffusivities for
 lysozyme in 191
 particle size distribution 132
 pore size of 129
- specific hydraulic permeabilities 88,
 133
- spiking experiments 48, 49
- Spodoptera frugiperda* (Sf9) 42
- standard operating protocol (SOP) 50
- Staphylococcal Protein A* (SpA) 55,
 109, 110–111, 125
- static capacity 87
- static light scattering 23
- static mixers 146
- stationary phase 54, 63, 93, 139, 160,
 183, 227, 265, 312, 356
- steady-state chromatographic
 separation 64
- stepwise elution 70, 140
- steric exclusion, of molecules 94
- steric mass action (SMA) model 104,
 170–173, 176–178, 259–261,
 296–297
- stirred batch adsorption systems 203
- stoichiometric displacement (SD) model
 104, 169–171, 320, 323
- strong ion exchangers 104
- superficial velocity 77, 88–89, 139,
 140, 155, 187, 197, 298, 300, 306,
 369
- surface diffusion 193
- synthetic polymers 122–124, 134
- syringe pumps 142
- t**
- Temkin isotherm 167
- thermodynamic stabilities, of proteins
 34
- Thomas solution 288–290
- tortuosity factor 188, 192, 286
- transition analysis 266

- transmission electron microscopy (TEM) 115, 123, 129–130
- transport mechanisms, in protein chromatography 214
- Trp residues 30, 107
- true countercurrent moving bed (TMB) system 74–75, 382–386, 388–390
- turnaround time 370–373

- U**
- UV absorbance, in protein solution 20–21

- V**
- validation, of biopharmaceutical process development 50
- van der Waals forces 28, 66, 96, 159, 188, 196

- viral contamination 51
- virus clearance values of retroviridae virus 49
- virus filtration 55, 359
- viscous fingering effects 79, 358, 364
- volume overloaded 283, 326, 376

- W**
- weak cation exchangers 104, 106, 117, 251, 346–347
- weak ion exchangers 104
- wetting angle of agarose beads 122
- wetting angle of polymethacrylate 122
- width of mass transfer zone 282
- wild-type Protein A 110–111

- Z**
- zirconium salt treatment 124