

# 1

## Techniques for Enzyme Purification

Adrie H. Westphal<sup>1</sup> and Willem J. H. van Berkel<sup>1,2</sup>

<sup>1</sup> Wageningen University & Research, Laboratory of Biochemistry, Stippeneng 4, 6708WE Wageningen, The Netherlands

<sup>2</sup> Wageningen University & Research, Laboratory of Food Chemistry, Bornse Weiland 9, 6708WG, Wageningen, The Netherlands

### 1.1 Introduction

Biocatalysis is the chemical process through which enzymes or other biological catalysts perform reactions between organic components. Biocatalysis gives an added dimension to synthetic chemistry and offers great opportunities to prepare industrial useful chiral compounds [1, 2]. Depending on the goal of the chemical conversion and the costs involved, biocatalyst-driven reactions are performed using whole cell systems or isolated enzymes, either in free or immobilized form [3–5].

Initially, industrial applications utilizing isolated enzymes were mainly developed with amylases, lipases, and proteases [6–8]. These hydrolytic enzymes were usually applied in a partially purified form, also because crude enzyme preparations are often more stable than the purified ones. However, for obtaining highly pure products, especially in the pharmaceutical industry, the purity of the enzyme preparation can be a critical factor.

Many enzyme purification methods have been developed over the years. Traditional purification procedures make use of the physicochemical properties of the enzyme of interest. These procedures were developed during the twentieth century for elucidating enzyme mechanisms and solving protein three-dimensional structures but also appeared to be valuable for the preparation of highly pure biocatalysts. Yet, progress in the preparation of biocatalysts has been given the biggest boost by the amazing developments in recombinant DNA technology and the accompanying revolutionary changes in enzyme production, enzyme purification, and enzyme engineering [9].

Here, we describe our experiences with the contemporary techniques for enzyme purification. For more information about the practical issues of enzyme purification, the reader is referred to the “Guide to Protein Purification” in *Methods in Enzymology* 463 [10].

## 1.2 Traditional Enzyme Purification

Before summarizing the traditional enzyme purification methods, it is important to note that the purification of enzymes is made easier by the fact that they are such specific catalysts. This enables the determination of the amount of a given enzyme in units (where 1 unit [U] of enzyme activity is defined as the amount of enzyme that catalyzes the conversion of 1  $\mu\text{mol}$  substrate per minute) and its specific activity (in  $\text{U mg}^{-1}$ ) in crude extracts and after each purification step. The specific activity is a good indication of the purity and quality of the enzyme preparation, especially if the specific activity of the pure enzyme under defined conditions is known. During enzyme purification, the improvement in specific activity and the yield of the enzyme after each purification step can be summarized in a purification scheme. The purification factor (specific activity obtained after a purification step divided by that of the starting material) provides an insight into the “efficiency” of each step. If a pure enzyme is obtained, it also indicates the relative amount of that enzyme present in the starting material. A theoretical example of a purification scheme, comprising three purification steps, is shown in Table 1.1.

Enzymes that are used for biocatalysis are typically purified from microbial cells or from culture media after or during growth of microorganisms (in case of excreted proteins). The enzyme purification generally starts with a cleared cell extract in which the enzyme is present in a soluble form. If the enzyme to be purified is excreted into the culture medium, it is usually sufficient to remove the cells from the medium by centrifugation (for small-scale purifications) or by filtration (for large-scale industrial purifications). In the case of an intracellular enzyme, cells should be broken first to release the protein into solution. Depending on the type of cells, different techniques are employed. The microbial cells are first harvested from the culture medium by centrifugation and resuspended in a small amount of buffer. The cells can be broken using a variety of techniques, e.g. by treatment with enzymes that digest cell walls (e.g. lysozyme), followed by osmotic shock, by using lysis buffers containing detergents, by exposure to ultrasound using sonicators, by pushing cells under high pressure through a small orifice using a pressure cell system, or by grinding frozen cells in liquid nitrogen. Extracts thus obtained are cleared from unbroken cells and large, insoluble particles by centrifugation or filtration. To prevent enzyme inactivation during these treatments, and also in the following purification steps, the temperature

**Table 1.1** Imaginary traditional enzyme purification scheme.

Step	Volume (ml)	Activity (U)	Protein (mg)	Specific activity ( $\text{U mg}^{-1}$ )	Yield (%)	Purification factor
CE	500	3000	15 000	0.2	100	1.0
AS	100	2400	4000	0.6	80	3.0
IEC	45	1440	500	2.9	48	14.5
GF	50	1000	125	8.0	33	40.0


Steps: CE, cell extract; AS, ammonium sulfate fractionation; IEC, ion exchange chromatography; GF, gel filtration.

of the enzyme solution is usually kept around 4 °C. Proteolytic degradation of the enzyme to be purified can be precluded by adding a protease inhibitor cocktail during breaking of the cells.


Once a cell-free extract has been obtained, several methods can be employed for further purification of the desired enzyme. These separation methods can be roughly divided into the following categories: (i) selective precipitation, (ii) separation based on charge, (iii) separation based on molecular size, (iv) separation based on bio-affinity, and (v) separation based on adsorption principles. Except for the first category, all these methods generally make use of column chromatography, with column sizes depending on the scale of the sample volumes and protein concentrations.

The strategy applied during enzyme purification is such that separation methods belonging to different categories are carried out in a logical order until the goal is reached. A good purification results in the recovery of most of the enzyme activity (i.e. a high yield) and in removal of many “contaminating” proteins and other types of (bio)molecules (i.e. a strong increase in specific activity). An often-experienced phenomenon during purification is the inactivation and/or aggregation of the enzyme (Figure 1.1). Because of increased enzyme concentration in the final steps of purification, aggregation can occur. If proteases are still present, the enzyme becomes more and more the only target for the protease, which can lead to proteolysis. In addition, wrong physical conditions (pH, temperature, and ionic strength) can lead to (partly) unfolding, followed by aggregation and/or proteolysis. Changing the type of buffer, pH, and/or ionic strength and the addition of protecting agents may alleviate these processes.

The purity of the final enzyme preparation can be tested in several ways. The most common methods used are sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) (Figure 1.2), analytical gel filtration, and mass spectrometry [11].



**Figure 1.1** Enzyme aggregation and proteolytic degradation processes.




**Figure 1.2** Example of an SDS-PAGE gel. (1) Molecular mass markers, (2) cell extract, (3) sample after ammonium sulfate fractionation, (4) sample after ion exchange chromatography, and (5) sample after gel filtration. Mr., relative molecular mass (kDa).

Traditional enzyme purification procedures many times start with an ammonium sulfate fractionation. This type of fractionation makes use of the fact that individual proteins precipitate at different concentration ranges of ammonium sulfate [12].

To make an estimation of the fractionation range, a small-scale pilot experiment can be performed. For such an experiment, different amounts of ammonium sulfate (from 0% to 90% saturation) are added to small samples of cell extract (usually, 1 ml). After dissolving the ammonium sulfate and removal of the formed protein precipitates by centrifugation, enzyme activity of the supernatants is measured (Figure 1.3). Such an analytical pilot experiment tells us at which saturation value the enzyme starts to precipitate (in our pilot, around 30%) and at which degree of saturation precipitation of the enzyme is more or less complete (in our pilot, around 65%). Once these values have been determined, the bulk of the cell extract is fractionated using these percentages and the precipitate obtained after the second addition of ammonium sulfate is used for further purification. If desired, removal of ammonium sulfate can be accomplished by dialysis, ultrafiltration, or gel filtration (e.g. with desalting columns).

Ammonium sulfate fractionation has been used in our group for the purification of several oxidoreductases. For the purification of vanillyl alcohol oxidase from *Penicillium simplicissimum* [13], the ammonium sulfate fractionation of the cell extract from 30% to 60% saturation gave a yield of 85%. Although few protein impurities were removed (as judged from SDS-PAGE and from the rather low purification factor of 1.2), this step appeared to be advantageous for the subsequent purification using a Phenyl Sepharose column, especially because ammonium sulfate removal could be omitted before this hydrophobic interaction chromatography (HIC) step. A similar experience was made with the purification of catalase peroxidase from *P. simplicissimum* [14] and 4-hydroxybenzoate 3-hydroxylase from



**Figure 1.3** Ammonium sulfate fractionation. In this example, a pilot experiment is performed on small cell extract samples. Next, 30% saturation is used on the total extract sample and the precipitates formed are removed by centrifugation. Then, the supernatant is brought to 65% saturation and the precipitate, which contains most of the enzyme activity, is collected by centrifugation.

*Rhodococcus opacus* 557 [15]. With hydroquinone dioxygenase from *Pseudomonas fluorescens* ACB [16], the cell-free extract was adjusted to 25% ammonium sulfate saturation before loading onto a Phenyl Sepharose column. Ammonium sulfate can also be used to concentrate the solution during enzyme purification. In the case of 4-hydroxybenzoate 1-hydroxylase from *Candida parapsilosis* [17], the enzyme fraction obtained after Q-Sepharose ion exchange chromatography (IEC) was adjusted to 70% saturation with pulverized ammonium sulfate and the resulting precipitate, collected by centrifugation, was dissolved in a small amount of buffer.

### 1.2.1 Ion Exchange Chromatography

IEC is one of the most widely used methods for enzyme purification. It separates protein molecules according to their differences in charge [18]. The stationary phase (matrix) in IEC carries charged functional groups fixed by chemical bonds. The fixed groups are associated with exchangeable counterions. In anion exchange chromatography, the fixed groups have positive charges and in cation exchange chromatography, these groups are negatively charged. As a rule of thumb, proteins bind to an anion exchanger at pH values above their isoelectric point (pI) and to a cation exchanger at pH values below the pI. Protein IEC usually involves the following steps:

- (1) *Equilibration*: The ion exchange resin is equilibrated with a low-salt buffer that allows binding of the enzyme of interest.
- (2) *Sample application and adsorption*: Protein molecules with a proper charge displace counterions and bind reversibly to the matrix. The ionic strength of the buffer in which

the protein sample is loaded should be low, as a high concentration of salt usually prevents binding. Any volume of sample can be applied as long as the total amount of protein does not exceed the binding capacity of the matrix. Yet, a large sample volume having a low concentration of precipitates may eventually clog the column. Proteins in the sample not bound by the matrix can be washed from the column using a loading buffer. To follow elution of nonbinding proteins, the absorbance of the column effluent can be monitored at 214 or 280 nm in the case of a low amount of protein in the sample or at 305 nm in the case of high protein concentrations.

- (3) *Desorption of bound proteins:* A stepwise increase of salt concentration or, in most cases, a gradual increase of the salt concentration (gradient) of the elution buffer is used. Again, the elution of proteins can be monitored by measuring the absorbance of the column effluent at 214 or 280 nm and, in addition, at a visible wavelength in the case of colored proteins. Elution with a shallow continuous gradient has the advantage that proteins with small differences in pI values are better separated and elute from the column in sharp, symmetrical peaks. For some enzymes, activity may be lost at high salt concentrations (e.g. because of dissociation of subunits). In that case, an elution can be attempted using a pH change step or a pH change gradient.
- (4) *Cleaning of the column:* Proteins and other substances that are bound very strongly to the column are removed. This is usually done by “cleaning-in-place,” using 2 M NaCl or 0.5 M NaOH solutions, followed by washing with water/buffer and 20% ethanol for storage.

IEC is a very powerful (preparative) purification method because (i) the high binding capacity of ion exchange columns allows elution of proteins in a very concentrated form and (ii) a proper choice of elution conditions results in separation of the bound proteins at high resolution.

For many years, IEC was included in almost every enzyme purification procedure, both on lab scale and at industrial level. Although this picture has changed after the introduction of the recombinant DNA technology providing the use of affinity tags, IEC remains a superior technology for enzyme purification because of its large resolving power and high recovery of enzyme activity.

In our experience, the IEC technique appeared to be crucial for the purification of a wide range of oxidoreductases, including monooxygenases, oxidases, dioxygenases, peroxidases, reductases, and dehydrogenases (Table 1.2).

A specific application of IEC involved the separation of native and oxidized forms of the flavoenzyme 4-hydroxybenzoate 3-hydroxylase from *P. fluorescens* [21]. The sensitivity of this dimeric enzyme to air oxidation resulted in different isoforms, which could be separated on a preparative scale with a DEAE-Sepharose column (Figure 1.4). Further analysis with an analytical Mono-Q column and isoelectric focusing experiments revealed the 10 different isoforms possible, assigned to combinations of the sulfhydryl, sulfenic acid, sulfinic acid, and sulfonic acid state of the surface-accessible Cys116 of each subunit. Mixing a native enzyme and a fully oxidized enzyme resulted in extremely slow formation of hybrid dimers with one native and one fully oxidized subunit, pointing to the high stability of the enzyme dimer.

**Table 1.2** Ion exchange chromatography of oxidoreductases.


Enzyme family	Enzyme	References
Flavoprotein hydroxylases	4-Hydroxybenzoate 3-hydroxylase	[14, 19–22]
	4-Hydroxybenzoate 1-hydroxylase	[16]
	3-Hydroxyphenylacetate 6-hydroxylase	[23]
	Hydroquinone hydroxylase	[24]
	Phenol hydroxylase (PheA1)	[25]
	3-Hydroxybenzoate 6-hydroxylase	[26]
Baeyer–Villiger monooxygenases	4-Hydroxyacetophenone monooxygenase	[27]
Copper-dependent monooxygenases	Polyphenol oxidase (tyrosinase)	[28]
	Lytic polysaccharide monooxygenase	[29]
Flavoprotein oxidases	Vanillyl alcohol oxidase	[13]
	Eugenol oxidase	[30]
Multicopper oxidases	Laccase-like multicopper oxidase	[31, 32]
Non-heme iron dioxygenases	Hydroquinone dioxygenase	[33]
Heme-dependent peroxidases	Catalase peroxidase	[12]
	Cationic peroxidase	[34]
Reductases	NADH reductase	[25, 35]
	Flavin reductase (PheA2)	
Nicotinamide-dependent dehydrogenases	Alcohol dehydrogenase	[36]
	Carveol dehydrogenase	[37]
Flavin-dependent dehydrogenases	Galactonolactone dehydrogenase	[38, 39]
	Proline dehydrogenase	[40]

Most of the listed enzymes were purified with several traditional separation methods described in this review. See references for details.

### 1.2.2 Gel Filtration

In gel filtration, also referred to as molecular sieve or size exclusion chromatography (SEC), sample molecules do not bind to the column but are fractionated based on their relative size and shape [41]. The liquid phase in such a column (total volume,  $V_t$ ) has two measurable volumes: external or “void” volume, consisting of the liquid between the beads ( $V_0$ ), and the internal volume ( $V_i$ ), constituted by the liquid within the pores of the beads. Molecules being too large to enter the pores cannot equilibrate with  $V_i$  and therefore emerge first from the column, while small molecules can equilibrate with  $V_i$  and therefore elute later.

The most important parameters in SEC are (i) the diameter of the pores allowing access to the internal volume of the beads, (ii) the total internal volume of the beads, (iii) the hydrodynamic diameter of the sample molecules, (iv) the flow rate of the liquid phase, and (v) the operation temperature and viscosity of the buffer used.



**Figure 1.4** IEC of isoforms of highly pure 4-hydroxybenzoate 3-hydroxylase (PHBH) from *Pseudomonas fluorescens*. Preparative separation on DEAE-Sepharose CL-6B (650 mg protein) using a gradient elution. The small peaks after the main peak contain differently oxidized forms of PHBH. Source: Modified from van Berkel and Müller [21].


Elution volumes of fractionated molecules should be intermediate between  $V_0$  and  $V_t$ . The elution volume ( $V_e$ , Figure 1.5a) relates to the accessibility of the molecule to the pores of the beads:  $V_e = V_0 + K_{AV} \cdot V_i$  (where the partition coefficient  $K_{AV} = (V_e - V_0)/(V_t - V_0)$ ). A semi-logarithmic plot illustrating the relation between  $K_{AV}$  and protein molecular weight ( $M_r$ ) is given in Figure 1.5b. The separation of proteins according to  $M_r$  is greatest in the central, linear region of the sigmoidal curve, spanning  $K_{AV}$  values between 0.2 and 0.8. This span is described as the fractionation range of a size exclusion matrix. A steep slope of the sigmoidal curve indicates a large resolving power of a matrix for a certain molecular weight range.

Next to being a suitable purification step [28, 29, 35, 40, 42], SEC is extremely useful to get information about the molecular weight of the native protein and its possible subunit composition [43]. By using this technique, we established that 4-hydroxybenzoate 3-hydroxylase from *P. fluorescens* is a homodimer, both in its holo and apo form [19, 44]. For lipoamide dehydrogenase from *P. fluorescens*, we experienced that nicotinamide adenine dinucleotide reduced (NADH) binding strongly stimulates flavin adenine dinucleotide (FAD)-induced dimerization [45].

For vanillyl alcohol oxidase from *P. simplicissimum*, we found that the holoenzyme favors the octameric state [13, 46], whereas the apoenzyme [47] mainly exists as a dimeric species. The octamer–dimer equilibrium of the holoenzyme varied with the ionic strength of the buffer solution, with kosmotropic salts stimulating the octameric state [48, 49]. More recently, it was established that a single loop at the protein surface is essential for the octamerization of vanillyl alcohol oxidase [30].

For hydroquinone dioxygenase from *P. fluorescens* ACB, we obtained strong indications from gel filtration that this non-heme, iron-dependent enzyme is an  $\alpha_2\beta_2$





**Figure 1.5** Gel filtration. (a) Elution profile and (b)  $K_{AV}$  vs  $\log M_r$  plot.

heterotetramer [33]. With proline dehydrogenase from *Thermus thermophilus*, we showed that the native enzyme is a homotetramer [40, 50] and that dimerization of the protein subunits strongly increases the enzyme thermostability [51]. For 3-hydroxybenzoate 6-hydroxylase from *Rhodococcus jostii* RHA1, we obtained evidence that the monomer-monomer contact of the dimer is stabilized by the binding of a phosphatidylinositol ligand [20].

### 1.2.3 Bio-affinity Chromatography

Bio-affinity chromatography is one of the most powerful procedures in protein purification. This method has a very high selectivity as it utilizes the specific, reversible

interactions between biomolecules [52]. Classic enzyme affinity chromatography mainly focused on methods that made use of the specific interactions of enzymes with ligands, such as substrates, coenzymes, inhibitors, and activators. Through immobilization of such ligands on suitable matrices, enzymes can be selectively bound to these resins (see example of old yellow enzyme given below). Preferably, the dissociation constant ( $K_d$ ) of an enzyme-immobilized ligand complex should not change substantially compared to that of the enzyme–ligand complex free in solution. The dissociation constant range of the complex may vary from micromolar (enzyme–coenzyme complexes) to nanomolar (enzyme inhibitor complexes).

In bio-affinity chromatography, proteins to be purified are brought onto a column containing the immobilized ligand. After application of sample to the column, nonbinding proteins are washed out. Protein(s) that are retained on the column by their specific interaction with the ligand are removed by changing the elution conditions. The most specific way is by using a soluble ligand, which is competitive for the matrix-bound ligand with which the enzyme is associated. Bio-specific elution is not always possible, in which case elution can be stimulated by, for example, using a gradient of increasing salt concentration or by changing the pH-value of the elution buffer.

Because bio-affinity chromatography makes use of specific interactions, it may result in a high degree of purification. In some cases, using this technique, an enzyme can be obtained from a crude extract almost completely pure in a single chromatographic step (see example of Old Yellow Enzyme given below). However, commercially available bio-affinity resins are often very costly and have a limited choice in coupled ligands. In addition, these resins are also not easily prepared “at home,” especially when expensive biomolecules must be used as immobilized ligands. Bio-affinity columns containing such ligands are usually more difficult to clean than, e.g., ion exchangers; therefore, the lifetime of a bio-affinity column is often limited.


We applied traditional bio-affinity chromatography for the purification of a number of enzymes, ranging from oxidoreductases to transferases. For 4-hydroxybenzoate 3-hydroxylase, we developed a Cibacron Blue dye affinity matrix, which appeared to be very useful for increasing the specific activity and yield of the enzyme, as isolated from different microbial sources [14, 15, 19–21]. Glutathione *S*-transferase isoenzymes from rat liver were purified using *S*-hexylglutathione affinity chromatography, followed by chromatofocusing on a Mono-P column [53]. A novel branched-chain alcohol dehydrogenase was purified from *Saccharomyces cerevisiae* using a Procion Red dye affinity column, which was selected based on its capacity to bind to a wide range of nicotinamide adenine dinucleotide phosphate (NADP)-dependent enzymes [36].

Old Yellow Enzyme from *Saccharomyces carlsbergensis* is the canonical member of a large family of ene reductases [54]. These flavoenzymes catalyze the asymmetric *trans*-hydrogenation of alkenes, resulting in industrially relevant chiral products [55]. Because Old Yellow Enzyme strongly interacts with phenolic compounds that act as competitive inhibitors, the enzyme was purified originally in high yield from brewer's bottom yeast by affinity chromatography using *N*-(4-hydroxybenzoyl)aminoethyl agarose [56–58]. This affinity matrix was prepared from agarose in four steps [56]:

- (1) Agarose beads were equipped with an aminohexyl spacer arm by activating the beads with cyanogen bromide in the presence of 1,6-diaminohexane.
- (2) The resulting aminohexyl agarose was reacted with 4-acetoxybenzoic acid to give *N*-(4-acetoxybenzoyl)aminohexyl agarose.
- (3) Remaining free amino groups were acetylated with acetic anhydride.
- (4) The protecting acetoxy group was removed from the ligand by incubation with imidazole, yielding the *N*-(4-hydroxybenzoyl)aminohexyl agarose affinity matrix (Figure 1.6).

The purification of Old Yellow Enzyme from brewer's bottom yeast then went as follows [56]:

- (1) 350 g dried yeast was suspended in 1 l of demineralized water, containing 10  $\mu$ M phenylmethylsulfonylfluoride to inactivate serine proteases.
- (2) The suspension was homogenized for 30 seconds at the high-speed setting of a Waring Blender.
- (3) *Autolysis*: the mixture was transferred to a glass beaker and mechanically stirred for 4 hours at 37 °C. All subsequent operations were performed at 0–4 °C.
- (4) The extract was clarified by centrifugation and precipitated with solid ammonium sulfate to 78% saturation.
- (5) *Enzyme reduction to remove phenolic ligands*: the precipitate was collected by centrifugation and dialyzed overnight against 6 l of 0.1 M Tris-HCl pH 8.0, containing 0.1 M ammonium sulfate, 10  $\mu$ M phenylmethylsulfonylfluoride, and 10 mM sodium dithionite.
- (6) *Enzyme reoxidation*: dialysis with the same buffer, omitting sodium dithionite, continued for another 6 hours with one additional buffer change.
- (7) Centrifugation to remove a white precipitate and stirring the clarified yeast extract for another 30 minutes to ensure reoxidation.
- (8) A column of *N*-(4-hydroxybenzoyl)aminohexyl agarose (Figure 1.6, bed volume 20 ml) was washed with 0.1 M Tris-HCl pH 8.0, containing 0.1 M ammonium sulfate and 10  $\mu$ M phenylmethylsulfonylfluoride.
- (9) The clarified yeast extract was applied on the column and the column was extensively washed with buffer (about 2 l) until the absorbance at 280 nm is lower than 0.2.
- (10) The Old Yellow Enzyme was eluted with 400 ml of washing buffer, which was degassed, flushed with oxygen-free nitrogen, and supplemented with 3 mM sodium dithionite.



**Figure 1.6** *N*-(4-Hydroxybenzoyl)aminohexyl agarose affinity matrix.

- (11) The enzyme eluted directly upon flavin reduction and turned bright yellow after reoxidation by air.
- (12) The collected enzyme (about 100 ml) was concentrated by ultrafiltration and stored frozen in 1 ml aliquots.
- (13) Regeneration of the affinity matrix was accomplished by washing the agarose beads with 0.2 M acetate buffer pH 5.0, containing 6 M GuHCl.
- (14) Storage of the gel in 10% ethanol with 1 mM sodium azide to prevent microbial damage.

SDS-PAGE showed that the Old Yellow Enzyme from *S. carlsbergensis* (relative subunit molecular mass = 49 kDa) was obtained in pure form. Absorption spectral analysis confirmed that the dimeric enzyme contained one tightly bound molecule of flavin mononucleotide (FMN) per subunit. The yield of enzyme was about 85% (130 mg) and its specific activity (turnover number) was slightly higher than the value obtained for the enzyme purified by conventional procedures.

Besides IEC, there are other chromatographic separation methods that make use of the properties of a protein's surface for adsorption to a specific chromatographic resin. The most commonly applied methods are described below.

#### 1.2.4 Hydrophobic Interaction Chromatography

HIC is a very useful technique for the fractionation of proteins [59]. In proteins, some hydrophobic groups, or clusters of hydrophobic groups, can occur at the surface of a protein and thus contribute to the surface hydrophobicity. The surface hydrophobicity allows a protein to undergo hydrophobic interactions not only with other proteins but also with column materials carrying hydrophobic groups.

Hydrophobic interactions between nonpolar compounds are enhanced by a polar environment and are energetically favorable because of a gain in entropy on forming. It is the liberation of ordered water molecules in contact with hydrophobic surfaces that drives clustering of hydrophobic groups. It follows that hydrophobic interactions will be affected if the structure of water is changed by dissolved salts or organic solvents. Kosmotropic salts (e.g. ammonium sulfate) tend to favor the strength of hydrophobic interactions, whereas chaotropic salts (e.g. sodium thiocyanate) disrupt the structure of water and thus tend to decrease the strength of hydrophobic interactions. Organic solvents are also commonly used to alter the polarity of water.

Although the mechanisms of hydrophobic interactions are complicated, chromatographic techniques based on hydrophobic interactions are easy to use. The most common resins for HIC are substituted with *n*-butyl, *n*-octyl, or phenyl groups. For an uncharacterized protein, as a start, phenyl-substituted resin is usually the best choice because strongly hydrophobic proteins are not easily eluted from the highly hydrophobic octyl-substituted resins. The phenyl ligand is intermediate in hydrophobicity between *n*-butyl and *n*-pentyl and will bind to aromatic amino acids through  $\pi$ - $\pi$  interactions.

A salt concentration just below that used for salting out of a protein is normally used for binding the protein to a hydrophobic matrix. A common procedure is to start purification of a protein from a crude extract with an ammonium sulfate precipitation at a

concentration of ammonium sulfate that leaves the protein of interest just in solution, followed by removal of proteins that precipitate at this salt concentration by centrifugation, and loading the clarified extract onto an HIC column.

Proteins bound to an HIC column are eluted by reducing the concentration of kosmotropic salt (e.g. ammonium sulfate) in the buffer using a negative gradient. This successively releases proteins from the column in order of hydrophobicity. Proteins that are tightly bound such that they do not elute at zero salt concentration can be eluted using a positive gradient of a polarity-reducing organic solvent (usually up to 50% ethylene glycol).

We applied the HIC technology in quite some traditional enzyme purifications, usually after ammonium sulfate fractionation, and often in combination with IEC and an additional chromatographic step. This enabled us to characterize the catalytic properties of several oxidoreductases [12–14, 23, 24, 27, 33, 37, 39, 60, 61] (see also Table 1.2).


A special case of HIC concerns the purification of the recombinant forms of lipoamide dehydrogenase from *A. vinelandii* [62] and *P. fluorescens* [63]. Both these enzymes, overproduced in *Escherichia coli*, could be purified in a single chromatographic step by binding them to a Sepharose 6B gel filtration column, equilibrated in 0.1 M potassium phosphate buffer pH 7.0, containing ammonium sulfate at 50% saturation. After washing with equilibration buffer, the lipoamide dehydrogenases were obtained in pure form by eluting with the same buffer at 25% ammonium sulfate saturation.

We also applied the HIC technology for the reversible removal of the flavin cofactor in a number of flavoproteins, including lipoamide dehydrogenase, glutathione reductase, mercuric reductase, and butyryl-CoA dehydrogenase [64, 65]. Figure 1.7 presents a schematic overview of the procedures. Each flavoprotein is bound to Phenyl Sepharose in a high-salt buffer. After changing to low pH, the FAD cofactor is released from the bound protein. Reconstitution of holoenzyme with natural FAD, chemically modified FAD, or isotopically enriched FAD (FAD\*) is performed on-column at neutral pH. Next, the reconstituted protein is released from the column with 50% ethylene glycol. Alternatively, the apoprotein is released from the column with 50% ethylene glycol and the holoenzyme is reconstituted in solution. The HIC technology appeared to be superior to classical methods for the reversible dissociation of the FAD cofactor of these enzymes, especially because the reversible immobilization procedure gave excellent yields and could be applied at large scale.

### 1.2.5 Hydroxyapatite Chromatography

Another adsorption chromatography method for enzyme purification concerns hydroxyapatite (HAP) chromatography [66]. HAP is a chromatographic support consisting of calcium phosphate crystals.

The amino groups of proteins adsorb to HAP primarily as a result of nonspecific electrostatic interactions between their positive charges and negative charges on the HAP column when the column is equilibrated with phosphate buffer. The carboxyl groups in proteins bind specifically by complexation to the calcium sites on the column. It was also found that basic proteins are eluted either as a result of normal Debye–Hückel charge screening or by a specific displacement with  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  ions, which



**Figure 1.7** Preparation and reconstitution of apo-flavoproteins on Phenyl Sepharose. FAD\* is isotopically enriched ( $^{13}\text{C}$ ,  $^{15}\text{N}$ )-FAD. Source: Modified from van Berket et al. [64].

form a complex with column phosphates and neutralize their negative charges. Acidic proteins are eluted by displacement of their carboxyl groups from calcium sites by ions that form stronger complexes with calcium than do carboxylate groups, e.g. fluoride or phosphate. The ineffectiveness of chloride as an eluent of acidic proteins is due to the fact that it does not form a complex with  $\text{Ca}^{2+}$  and, thus, cannot compete with the calcium carboxylate complexes. The ability of  $\text{CaCl}_2$  and  $\text{MgCl}_2$  to strengthen the interaction of acidic proteins with HAP is due to the formation of additional bridges between protein carboxyl groups and column phosphate sites. In practice, proteins are often applied in 5–10 mM phosphate buffer at pH 7.0 and eluted with an increasing gradient of phosphate buffer.

We successfully applied HAP chromatography for the purification of vanillyl alcohol oxidase [13, 60], NADH reductase [35], catalase peroxidase [12], 4-hydroxybenzoate 3-hydroxylase [20], 4-hydroxybenzoate 1-hydroxylase [61], carveol dehydrogenase [37], flavin reductase PheA2 [25], and galactonolactone oxidoreductase [39] (see also Table 1.2). In most cases, HAP chromatography was introduced at the final stage of purification, after IEC or HIC, and before SEC.

### 1.3 Example of a Traditional Enzyme Purification Protocol

We already discussed the theoretical aspects and some specific applications of the traditional enzyme purification methods. In this paragraph, we describe the purification of 3-hydroxyphenylacetate 6-hydroxylase from *Flavobacterium* JS-7 as an example of a traditional enzyme purification protocol [23].

- (1) 50 g of *Flavobacterium* JS-7 cells, grown with 0.1% phenylacetic acid as a sole source of carbon and energy, was suspended in 10 mM potassium phosphate at pH 7.0, containing 0.5 mM ethylenediaminetetraacetic acid (EDTA) (for complexation of heavy metal ions), 0.5 mM dithiothreitol (DTT) (to prevent cysteine oxidation), 1.5 mM 3-hydroxyphenylacetic acid (binding of this aromatic substrate in the active site might protect the enzyme from inactivation), and 10  $\mu\text{M}$  FAD (to prevent apoenzyme formation). Next, 1 mg of DNase was added and cells were broken in a precooled French Press. The resulting cell extract was clarified by centrifugation at 4 °C. All further operations were performed between 0 and 4 °C using the abovementioned starting buffer omitting the aromatic substrate. After each chromatographic step, the enzyme solution was concentrated by ultrafiltration and desalted through either dialysis or gel filtration before performing the next step.
- (2) Cell extract, containing about 3 g of protein, was loaded onto a DEAE-Sepharose CL-6B column (17  $\times$  2.4 cm; bed volume 77 ml). After washing with five volumes of starting buffer, the enzyme was eluted with a 0–0.5 M KCl gradient in 1000 ml of starting buffer. Pooled active fractions eluting between 0.10 and 0.15 M KCl were loaded onto a Phenyl Sepharose CL-4B column (2.4  $\times$  9 cm; bed volume 41 ml), equilibrated with 15% ammonium sulfate. After washing with five volumes of equilibration buffer, the enzyme was eluted with a decreasing gradient of 15 to 0% ammonium sulfate in 500 ml starting buffer. Pooled active fractions eluting between 8.4 and 0% ammonium sulfate were

**Table 1.3** Purification of 3-hydroxyphenylacetate 6-hydroxylase from *Flavobacterium JS-7*.

Step	Activity (U)	Protein (mg)	Spec. activity (U mg <sup>-1</sup> )	Yield (%)
CE	1050	3080	0.3	100
IEC	799	250	3.2	76
HIC	712	60	11.9	68
BAC	510	28	18.2	49

Source: Adapted from Van Berkel and van den Tweel [23].

Steps: CE, cell extract; IEC, ion exchange chromatography; HIC, hydrophobic interaction chromatography; BAC, bio-affinity chromatography.

loaded onto a Reactive Red 120 agarose column (2.4×9 cm; bed volume 41 ml). After washing with five volumes of starting buffer, the enzyme was eluted with a linear gradient of 0–1 M KCl in 500 ml starting buffer. Pooled active fractions eluting between 0.16 and 0.46 M KCl were concentrated in 50 mM starting buffer, and 1 ml aliquots with a protein concentration of 2 mg ml<sup>-1</sup> was distributed over Eppendorf tubes, flash frozen in liquid nitrogen, and stored at –70 °C. The results of this purification procedure are summarized in Table 1.3.

## 1.4 Purification of Recombinant Enzymes

Developments in recombinant DNA technology and genome sequencing at the turn of the millennium have led to new procedures for the discovery and production of biocatalysts (see also Chapter 2). Nowadays, most biocatalysts are produced in heterologous hosts through the expression of synthetic genes. Many strategies for the expression of these recombinant proteins exist [67–69]. Many new biocatalysts are produced as fusion proteins, either to allow their rapid and efficient purification, enhance their proper folding and solubility, or facilitate cofactor regeneration [70].

One of the most widely applied methods for the purification of recombinant proteins concerns immobilized metal affinity chromatography (IMAC). This versatile method is described in the following section.

### 1.4.1 Immobilized Metal Affinity Chromatography

IMAC is a specialized adsorption chromatography technique that has turned into a powerful tool for single-step purification of recombinant proteins into which a metal binding site (“His-tag”) has been manufactured by genetic modification [71]. In IMAC, transition metal ions such as cobalt, nickel, or zinc act as electron acceptors (Lewis acids) for groups with electron-donating atoms (Lewis bases). The imidazole nitrogen atoms of His-tags introduced at the N- or C-terminal ends of proteins appear to be perfect electron donors for these metal ions.



In order to utilize this interaction for chromatographic purposes, the metal ion must be immobilized onto an insoluble support. The most widely applied chelating group for that is nitrile-triacetic acid (NTA), which has four bases for binding a nickel ion.

Binding of proteins to IMAC columns is usually best between pH 6.0 and pH 8.0 where the imidazole groups of the histidine residues are deprotonated. Chelating agents such as EDTA or citrate, but also Tris buffer, can reduce the binding strength. In order to prevent a metal affinity column to also function as an ion exchanger, buffers for loading, washing, and eluting proteins usually contain a relatively high ionic strength.

His-tags can be inserted at the N- or C-terminal ends of proteins or within exposed loops. The tag is added by modification of the gene encoding the protein to include codons for 6 to 10 consecutive histidine residues. These codons can be added by site-directed mutagenesis or by cloning the gene directly, in the correct reading frame, into a vector that contains these codons, resulting in a fusion protein. It is of major importance that the inserted histidine residues do not prohibit proper folding of the protein nor interfere with its enzyme activity. Site-directed mutagenesis also facilitates removal of His-tags after purification of the protein. This is commonly done by insertion of a recognition site for a highly specific endopeptidase between His-tag and native protein.

Ni-NTA metal affinity columns have a high binding capacity (5–10 mg protein per ml packed gel). His-tagged proteins bound to Ni-NTA are eluted from the column by inclusion of a high-concentration (250–500 mM) imidazole in the elution buffer. Before eluting His-tagged proteins, it is worth washing the column with a low concentration of imidazole (10–50 mM) in order to remove weakly interacting proteins.

We used the His-tag technology for the purification of several oxidoreductases including galactonolactone dehydrogenase from *Arabidopsis thaliana* [72], styrene monooxygenase from *R. opacus* 1CP [73, 74], 3-hydroxybenzoate 6-hydroxylase from *Rhodococcus jostii* RHA1 and *Pseudomonas alcaligenes* [26, 75], ene reductase from *R. opacus* CP1 [76], pyranose 2-oxidase from *Arthrobacter siccitolerans* [77], styrene monooxygenase reductase from *R. opacus* 1CP [78], vanillyl alcohol oxidase from *P. simplicissimum* [79], eugenol oxidase from *Rhodococcus jostii* RHA1 [80], 5-(hydroxymethyl)furfural oxidase from *Methylovorus* sp. strain MP688 [81, 82], and 4-hydroxybenzoate hydroxylase from *Cupriavidus necator* [83].

The hyperthermostable laminarinase from *Pyrococcus furiosus* (LamA) was also purified using the His-tag technique. The cell-free extract, obtained after breaking overexpressing *E. coli* cells and subsequent centrifugation, was subjected to Ni-NTA affinity chromatography, in which the LamA was eluted with a linear gradient of 0–0.5 M imidazole [42]. Next to highly active monomers, a minor part of the soluble enzyme constituted less active native-like oligomers and non-native monomers.

It is important to mention here that almost 50% of the overproduced LamA protein ended up in the pellet fraction that was obtained after initial centrifugation of the lysed cells. Notably, a large fraction of the soluble and insoluble LamA aggregates could be recovered in fully active form by incubating them for several hours at 80 °C in the presence of 3 M guanidine hydrochloride (GuHCl). Similar chaotropic heat treatment protocols might be useful for increasing the yield of other hyperthermostable enzymes that tend to aggregate during production in *E. coli*.

Obviously, this method is not applicable to less thermotolerant enzymes that initially are produced in inclusion bodies. For such His-tagged proteins, a soluble enzyme exhibiting the desired activity might be obtained by solubilizing them with a proper unfolding agent and subsequent purifying and refolding using a Ni-NTA affinity matrix [84].

Care must be taken regarding the correct unfolding and refolding conditions. For instance, for aryl-alcohol oxidase from *Pleurotus eryngii* overproduced in *E. coli* (without His-tag), the washed inclusion bodies were first solubilized in 20 mM Tris-HCl buffer at pH 8.0, containing 2 mM EDTA, 30 mM DTT, and 8 M urea. Next, optimization of the refolding conditions yielded a decently active enzyme after incubating the recombinant protein at 0.4 mg ml<sup>-1</sup> for 80 hours at 16 °C and pH 9.0 in the presence of 35% glycerol, 80 μM FAD, 0.6 M urea, 1 mM DTT, 2.5 mM oxidized glutathione, and 1.25 mM reduced glutathione [85].

Recently, Fraaije and coworkers used the IMAC technology for the purification of a self-sufficient cytochrome P450 monooxygenase from *Thermothelomyces thermophila* [86]. They also developed experimental protocols for screening of thermostable Baeyer-Villiger monooxygenases by purifying the His-tagged proteins with Ni-NTA beads in a 96-well plate format [87]. Robust cyclohexanone monooxygenases, stabilized by applying mutations found by computational and experimental library design, were characterized after purification by metal affinity chromatography [88].

A special application of Ni-NTA affinity chromatography concerns the deflavinylation and reconstitution of flavoproteins [89]. This generic method, originally developed for the NifL PAS domain from *A. vinelandii* [89], also turned out to be very useful for the reversible resolution of flavin and pterin cofactors of His-tagged *E. coli* DNA photolyase [90].

#### 1.4.2 Affinity Chromatography with Protein Tags


Next to IMAC, affinity chromatography procedures are available for enzyme purification that make use of a specific interaction of protein tags with accompanying resins. Well-known examples of such fusion tags are thioredoxin (Trx), glutathione transferase (GST), maltose binding protein (MBP), and small ubiquitin modifier (SUMO) [70]. These so-called solubility tags fold rapidly into a stable and highly soluble protein upon translation and fusion to the N-terminus of the enzyme of interest might stimulate the expression and solubility of the target enzyme. Again, removal of the tag from the purified protein can be facilitated by insertion of a recognition site for a highly specific endopeptidase between the tag and the native protein.

Dijkman and Fraaije used the SUMO technology for the production of 5-(hydroxymethyl)furfural oxidase (HFMO) from *Methylovorus* sp. strain MP688 [81]. To that end, they designed a His<sub>6</sub>-SUMO-HFMO gene construct, which, after ligation in a pET-SUMO vector, could be expressed in *E. coli* BL21(DE3) cells. After production, the His<sub>6</sub>-SUMO-HMFO fusion was cleaved with His<sub>6</sub>-SUMO protease. Subsequently, a Ni-NTA column was used to capture both the His<sub>6</sub>-SUMO tag and His<sub>6</sub>-SUMO protease, yielding purified HMFO in the flow through. A similar strategy was applied for the production of *Thermocrispum municipale* cyclohexanone monooxygenase (*Tm*CHMO) variants, with the aim of selecting for variants of *Tm*CHMO with changed stereo- and regioselectivity [91, 92] and substrate preference [93].

The gene coding for polycyclic ketone monooxygenase (PockeMO) from *Thermothelomyces thermophila* was fused into a His<sub>6</sub>-SUMO-containing pET vector (yielding pET-His<sub>6</sub>-SUMO-PockeMO) and also into a His<sub>6</sub> cofactor-recycling phosphite dehydrogenase (PTDH) containing the pBAD vector (yielding pBAD-His<sub>6</sub>-PTDH-linker-PockeMO) by Fraaije and coworkers [94]. For the expression of the SUMO fusion, *E. coli* BL21(DE3) was used and for expression of the PTDH fusion, *E. coli* NEB10 $\beta$  cells were used. Both constructs yielded high expression levels and the fusion proteins were easily purified with Ni-NTA affinity chromatography by exploiting the N-terminal His<sub>6</sub>-tags. PockeMO is active with bulky ketones and can perform enantioselective oxidations on steroids. The PTDH-PockeMO protein showed an increased activity compared to the native enzyme. After removal of the SUMO tag, applying the method described above for *Tm*CHMO, the protein could be crystallized and its structure solved. PockeMO exhibits the typical Bayer–Villiger monooxygenase organization with an FAD domain, an NADP domain, and a helical domain [94].

Fraaije and coworkers also showed that the flavoprotein alditol oxidase (AldO) from *Streptomyces coelicolor* is expressed at extremely high levels in *E. coli* when the enzyme is fused to MBP [95]. The extreme overproduction of the fused protein allowed for a single-step purification procedure using a Q-Sepharose ion exchanger. To find out if fused MBP-AldO protein behaves similarly to native AldO, the MBP tag was cleaved off with trypsin. Using amylose affinity chromatography, the MBP tag was removed. Electrospray ionization mass spectrometry analysis of free AldO established that cleavage had occurred at the expected site and that a homogeneous AldO preparation was obtained. AldO turned out to be active with the same range of substrates as found for fused MBP-AldO protein. AldO accepts various polyols, with xylitol and sorbitol being the preferred substrates. Intriguingly, AldO also oxidizes thiols such as DTT to the corresponding thiocarbonyls [82].

Incorporation of an N-terminal MBP solubility tag strongly increased the yield of recombinant form of the membrane-associated proline dehydrogenase from *T. thermophilus* (*Tt*ProDH) [37]. Expressed MBP-*Tt*ProDH comprised about 50% of the total protein content of expressing *E. coli* cells, and by applying amylose affinity chromatography, more than 250 mg of active enzyme was obtained per liter of culture. Native *Tt*ProDH was obtained by cleaving the fusion protein with trypsin in the presence of the detergent *n*-octyl  $\beta$ -D-glucopyranoside. Analytical gel filtration suggested that both native and MBP-fused *Tt*ProDH form tetramers that are prone to aggregation through non-native self-association. Site-directed mutagenesis (F10E/L12E variant) then showed that the hydrophobic N-terminal helix of ProDH is responsible for the self-association process [96]. Truncation of the N-terminal arm of *Tt*ProDH ( $\Delta$ A and  $\Delta$ AB variants) resulted in highly active tetramers [50], while selective disruption of two ion pairs in the dimerization interface of the enzyme (D205K/E207K variants) resulted in monomer formation (Figure 1.8) [51]. The newly created *Tt*ProDH monomer showed excellent catalytic properties but a significant lower thermal stability than the tetramer. Finally, using a riboflavin auxotrophic *E. coli* strain and MBP as a solubility tag, we also succeeded in producing the apoprotein of *Tt*ProDH. Reconstitution experiments together with structural studies and flavin content analysis led to the surprising conclusion that *Tt*ProDH does not discriminate between FAD and FMN as a cofactor [97].



**Figure 1.8** Oligomeric state of *TtProDH* and its site-directed mutants. EE: F10E/L12E variant.  $\Delta$ A: *TtProDH* without the N-terminal helix  $\alpha$ A.  $\Delta$ AB: *TtProDH* without the N-terminal helices  $\alpha$ A and  $\alpha$ B. KK: D205K/E207K variants of EE,  $\Delta$ A and  $\Delta$ AB. The flavin cofactor is shown in yellow, helix  $\alpha$ A in green, helix  $\alpha$ B in red, and helix  $\alpha$ C in blue. Source: Modified from Huijbers et al. [51].

## 1.5 Column Materials

A small selection of commercially available column materials is shown in Table 1.4. These bulk resins can be used to construct columns of various sizes in accordance with the type and volume of samples applied. In addition, a large number of prepacked columns for direct use are also available from these companies.

**Table 1.4** Selection of various commercially available chromatography media.

Ion exchange	Matrix	Functional group	Capacity (mg ml <sup>-1</sup> )	Brand
Capto Q	Highly cross-linked agarose with a dextran surface extender	Quaternary amine	High	1
Capto S	Highly cross-linked agarose with a dextran surface extender	Sulfonate group	High	1
Source 15Q	Rigid polystyrene – a divinyl benzene polymer	Quaternary amine	High	1
Source 15S	Rigid polystyrene – a divinyl benzene polymer	Sulfonate group	High	1
Macro-Prep High Q	Methacrylate copolymer	Quaternary amine	37 BSA	2

Table 1.4 (Continued)

Ion exchange	Matrix	Functional group	Capacity (mg ml <sup>-1</sup> )	Brand
Macro-Prep High S	Methacrylate copolymer	Sulfonate group	49 IgG	2
Macro-Prep DEAE	Methacrylate copolymer	Diethylaminoethyl	40 BSA	2
Macro-Prep CM	Methacrylate copolymer	Carboxy methyl	40 BSA	2
<b>Gel filtration</b>	<b>Matrix</b>	<b>Separation range (kDa)</b>		<b>Brand</b>
Sephacryl S-100 HR	Copolymer of allyl dextran and <i>N,N'</i> -methylene bisacrylamide	1–100		1
Sephacryl S-200 HR	Copolymer of allyl dextran and <i>N,N'</i> -methylene bisacrylamide	5–250		1
Sephacryl S-500 HR	Copolymer of allyl dextran and <i>N,N'</i> -methylene bisacrylamide	40–20000		1
Superdex 75 Prep Grade	Composite of cross-linked agarose and dextran	3–70		1
Superdex 200 Prep Grade	Composite of cross-linked agarose and dextran	10–600		1
Bio-Gel P-30	Copolymerization of acrylamide and <i>N,N'</i> -methylenebisacrylamide gel	2.4–40		2
Bio-Gel P-100	Copolymerization of acrylamide and <i>N,N'</i> -methylenebisacrylamide gel	5–100		2
Bio-Gel A 1.5m	Agarose	10–1500		2
<b>(Bio)-affinity</b>	<b>Matrix</b>	<b>Functional group</b>	<b>Capacity (mg ml<sup>-1</sup>)</b>	<b>Brand</b>
HiPrep Heparin FF	Cross-linked 6% agarose	Heparin	High	1
Ni Sepharose HP	Highly cross-linked agarose	Ni-charged	40	1
Glutathione Sepharose 4 FF	Highly cross-linked 4% agarose	Glutathione	10	1
Strep-Tactin <sup>®</sup> XT Sepharose	Rigid cross-linked agarose	Strep-Tactin	10	1
Blue Sepharose 6 Fast Flow	Cross-linked 6% agarose	Cibacron blue 3G	High	1
Nuvia <sup>™</sup> IMAC	Inert hydrophilic beads	Nitrilotriacetic acid	>40	2

(Continued)

**Table 1.4** (Continued)

<b>Ion exchange</b>	<b>Matrix</b>	<b>Functional group</b>	<b>Capacity (mg ml<sup>-1</sup>)</b>	<b>Brand</b>
Affi-Gel <sup>®</sup> Blue Gel	Cross-linked agarose	Cibacron blue F3GA	>11	2
<b>Hydrophobic interaction</b>	<b>Matrix</b>	<b>Functional group</b>	<b>Capacity (mg ml<sup>-1</sup>)</b>	<b>Brand</b>
Phenyl Sepharose 6 FF	Cross-linked 6% agarose	Phenyl	High	1
Butyl Sepharose 4 FF	Cross-linked 4% agarose	Butyl	High	1
Octyl Sepharose 4 FF	Cross-linked 4% agarose	Octyl	High	1
Macro-Prep <sup>®</sup> Methyl HIC	Methacrylate	Methyl	12	2
Macro-Prep <i>t</i> -butyl HIC	Methacrylate	<i>t</i> -Butyl	12	2
<b>Hydroxyapatite interaction</b>	<b>Matrix</b>	<b>Functional group</b>	<b>Capacity (mg g<sup>-1</sup>)</b>	<b>Brand</b>
Bio-Gel <sup>®</sup> HT Hydroxyapatite	Calcium phosphate	Calcium phosphate	10 BSA	2

1. GE Healthcare Life Sciences/Cytiva, VWR International B.V., Amsterdam, The Netherlands; 2. Bio-Rad Laboratories B.V., Veenendaal, The Netherlands.

## 1.6 Conclusions

Traditional enzyme purification procedures, whereby different chromatographic steps are required in order to obtain a pure protein, have played an eminent role in the development of the biocatalysis field. Nowadays, these methods have been replaced for the most part by affinity chromatographic techniques in which recombinant proteins fused with a specific tag can be effectively purified in a single step. Regarding the latter, it should be kept in mind that in many cases, an additional polishing step is required to remove critical impurities.

In this review, we have summarized our longstanding experience with the purification and characterization of redox enzymes. Purification procedures for these proteins do not significantly differ from the purification methods used for, e.g., hydrolases or transferases. However, with redox enzymes, and, in general, with proteins containing cofactors, potential cofactor dissociation during purification steps should be taken into account. Thus, addition of excess cofactor during purification can be essential for keeping the enzyme stable and active. We show that conventional chromatographic purification techniques can also be used for the large-scale preparation of apoproteins and for gaining insights into the molecular and hydrodynamic properties of the enzymes. Combining the specific

opportunities of the different chromatographic techniques with the still growing DNA-recombinant toolbox will make it possible in the future to identify and purify new exciting biocatalysts.

## References

- 1 Kohls, H., Steffen-Munsberg, F., and Höhne, M. (2014). Recent achievements in developing the biocatalytic toolbox for chiral amine synthesis. *Curr. Opin. Chem. Biol.* 19: 180–192.
- 2 Zmijewski, M. (2000). *Stereoselective biocatalysis*. New York: American Chemical Society.
- 3 Faber, K. (2018). *Biotransformations in organic chemistry*, 1–434. Heidelberg: Springer-Verlag.
- 4 Sheldon, R.A. and Brady, D. (2018). The limits to biocatalysis: pushing the envelope. *Chem. Commun.* 54: 6088–6104.
- 5 Woodley, J.M. (2019). Reaction engineering for the industrial implementation of biocatalysis. *Top. Catal.* 62: 1202–1207.
- 6 de Souza, P.M. and de Oliveira Magalhães, P. (2010). Application of microbial  $\alpha$ -amylase in industry: a review. *Braz. J. Microbiol.* 41 (4): 850–861.
- 7 Bornscheuer, U.T. and Kazlauskas, R.J. (2006). *Hydrolases in organic synthesis: regio- and stereoselective biotransformations*, 1–355. Weinheim: Wiley-VCH Verlag GmbH and Co. KGaA.
- 8 Filho, D.G., Silva, A.G., and Guidini, C.Z. (2019). Lipases: sources, immobilization methods, and industrial applications. *Appl. Microbiol. Biotechnol.* 103 (18): 7399–7423.
- 9 Bornscheuer, U.T., Huisman, G.W., Kazlauskas, R.J. et al. (2012). Engineering the third wave of biocatalysis. *Nature* 485 (7397): 185–194.
- 10 Burgess, R.R. and Deutscher, M.P. (eds.) (2009). *Guide to Protein Purification*. Methods in Enzymology 463 San Diego: Academic Press.
- 11 Rhodes, D.G. and Laue, T.M. (2009). Determination of protein purity. In: *Methods in Enzymology* (eds. R.R. Burgess and M.P. Deutscher), 463: 677–689. San Diego: Academic Press.
- 12 Burgess, R.R. (2009). Protein precipitation techniques. In: *Methods in Enzymology* (eds. R.R. Burgess and M.P. Deutscher), 463: 331–342. San Diego: Academic Press.
- 13 de Jong, E., van Berkel, W.J.H., van der Zwan, R.P. et al. (1992). Purification and characterization of vanillyl-alcohol oxidase from *Penicillium simplicissimum*. *Eur. J. Biochem.* 208 (3): 651–657.
- 14 Fraaije, M.W., Roubroeks, H.P., Hagen, W.R. et al. (1996). Purification and characterization of an intracellular catalase-peroxidase from *Penicillium simplicissimum*. *Eur. J. Biochem.* 235: 192–198.
- 15 Jadan, A.P., van Berkel, W.J.H., Golovleva, L.A. et al. (2001). Purification and properties of *p*-hydroxybenzoate hydroxylases from *Rhodococcus* strains. *Biochem. Mosc.* 66 (8): 898–903.
- 16 Moonen, M.J.H., Kamerbeek, N.M., Westphal, A.H. et al. (2008). Elucidation of the 4-hydroxyacetophenone catabolic pathway in *Pseudomonas fluorescens* ACB. *J. Bacteriol.* 190 (15): 5190–5198.

- 17 Eppink, M.H.M., Schreuder, H.A., and van Berkel, W.J.H. (1998). Interdomain binding of NADPH in *p*-hydroxybenzoate hydroxylase as suggested by kinetic, crystallographic and modeling studies of histidine 162 and arginine 269 variants. *J. Biol. Chem.* 273 (33): 21031–21039.
- 18 Jungbauer, A. and Hahn, R. (2009). Ion-exchange chromatography. In: *Methods in Enzymology* (eds. R.R. Burgess and M.P. Deutscher), 463: 349–371. San Diego: Academic Press.
- 19 Müller, F., Voordouw, G., van Berkel, W.J.H. et al. (1979). A study of *p*-hydroxybenzoate hydroxylase from *Pseudomonas fluorescens*. *Eur. J. Biochem.* 101 (1): 235–244.
- 20 Seibold, B., Matthes, M., Eppink, M.H.M. et al. (1996). 4-Hydroxybenzoate hydroxylase from *Pseudomonas* sp. CBS3. Purification, characterization, gene cloning, sequence analysis and assignment of structural features determining the coenzyme specificity. *Eur. J. Biochem.* 239 (2): 469–478.
- 21 van Berkel, W.J.H. and Müller, F. (1987). The elucidation of the microheterogeneity of highly purified *p*-hydroxybenzoate hydroxylase from *Pseudomonas fluorescens* by various biochemical techniques. *Eur. J. Biochem.* 167 (1): 35–46.
- 22 van Berkel, W.J.H., Westphal, A.H., Eschrich, K. et al. (1992). Substitution of Arg214 at the substrate binding site of *p*-hydroxybenzoate hydroxylase from *Pseudomonas fluorescens*. *Eur. J. Biochem.* 210 (2): 411–419.
- 23 van Berkel, W.J.H. and van den Tweel, W.J.J. (1991). Purification and characteriation of 3-hydroxyphenylacetate 6-hydroxylase: a novel FAD-dependent monooxygenase from a *Flavobacterium* species. *Eur. J. Biochem.* 201 (3): 585–592.
- 24 Eppink, M.H.M., Cammaart, E., van Wassenaar, D. et al. (2000). Purification and properties of hydroquinone hydroxylase, a FAD-dependent monooxygenase involved in the catabolism of 4-hydroxybenzoate in *Candida parapsilosis* CBS604. *Eur. J. Biochem.* 267 (23): 6832–6840.
- 25 Kirchner, U., Westphal, A.H., Müller, R. et al. (2003). Phenol hydroxylase from *Bacillus thermoglucosidasius* A7, a two-protein component monooxygenase with a dual role for FAD. *J. Biol. Chem.* 278 (48): 47545–47553.
- 26 Montersino, S. and van Berkel, W.J.H. (2012). Functional annotation and characterization of 3-hydroxybenzoate 6-hydroxylase from *Rhodococcus jostii* RHA1. *Biochim. Biophys. Acta* 1824 (3): 433–442.
- 27 Kamerbeek, N.M., Moonen, M.J.H., van der Ven, J.G.M. et al. (2001). 4-Hydroxyacetophenone monooxygenase from *Pseudomonas fluorescens* ACB. *Eur. J. Biochem.* 268 (9): 2547–2557.
- 28 Kuijpers, T.F.M., van Herk, T., Vincken, J.-P. et al. (2014). Potato and mushroom polyphenol oxidase activities are differently modulated by natural plant extracts. *J. Agric. Food Chem.* 62 (1): 214–221.
- 29 Frommhagen, M., Sforza, S., Westphal, A.H. et al. (2015). Discovery of the combined oxidative cleavage of plant xylan and cellulose by a new fungal polysaccharide monooxygenase. *Biotechnol. Biofuels* 8 (101): 12.
- 30 Ewing, T.A., Gygli, G., and van Berkel, W.J.H. (2016). A single loop is essential for the octamerization of vanillyl-alcohol oxidase. *FEBS J.* 283 (13): 2546–2559.
- 31 Tamayo-Ramos, J.A., van Berkel, W.J.H., and de Graaff, L.H. (2012). Biocatalytic potential of laccase-like multicopper oxidases from *Aspergillus niger*. *Microb. Cell Factories* 11: 165.



- 32 Ferraroni, M., Westphal, A.H., Borsari, M. et al. (2017). Structure and function of *Aspergillus niger* laccase McoG. *Biocatalysis* 3 (1): 10.
- 33 Moonen, M.J.H., Synowsky, S.A., van den Berg, W.A.M. et al. (2008). Hydroquinone dioxygenase from *Pseudomonas fluorescens* ACB: a novel member of the family of nonheme-iron(II)-dependent dioxygenases. *J. Bacteriol.* 190 (15): 5199–5209.
- 34 Dicko, M.H., Gruppen, H., Hilhorst, R. et al. (2006). Biochemical characterization of the major sorghum grain peroxidase. *FEBS J.* 273 (10): 2293–2307.
- 35 Weber, F.J., van Berkel, W.J.H., Hartmans, S. et al. (1992). Purification and properties of the NADH reductase component of alkene monooxygenase from *Mycobacterium* strain E3. *J. Bacteriol.* 174 (10): 3275–3281.
- 36 van Iersel, M.F., Eppink, M.H.M., van Berkel, W.J.H. et al. (1997). Purification and characterization of a novel NADP-dependent branched-chain alcohol dehydrogenase from *Saccharomyces cerevisiae*. *Appl. Environ. Microbiol.* 63 (10): 4079–4082.
- 37 van der Werf, M.J., van der Ven, C., Barbirato, F. et al. (1999). Stereoselective carveol dehydrogenase from *Rhodococcus erythropolis* DCL14: A novel nicotinoprotein belonging to the short chain dehydrogenase/reductase superfamily. *J. Biol. Chem.* 274 (37): 26296–26304.
- 38 Leferink, N.G.H., Heuts, D.P.H.M., Fraaije, M.W. et al. (2008). The growing VAO flavoprotein family. *Arch. Biochem. Biophys.* 474 (2): 292–301.
- 39 Kudryashova, E.V., Leferink, N.G.H., Slot, I.G.M. et al. (2011). Galactonolactone oxidoreductase from *Trypanosoma cruzi* employs a FAD cofactor for the synthesis of vitamin C. *Biochim. Biophys. Acta* 1814 (5): 545–552.
- 40 Huijbers, M.M.E. and van Berkel, W.J.H. (2015). High yields of active *Thermus thermophilus* proline dehydrogenase are obtained using maltose-binding protein as a solubility tag. *Biotechnol. J.* 10 (3): 395–403.
- 41 Stellwagen, E. (2009). Gel filtration. In: *Methods in Enzymology* (eds. R.R. Burgess and M.P. Deutscher), 463: 373–385. San Diego: Academic Press.
- 42 Westphal, A.H., Geerke-Volmer, A.A., van Mierlo, C.P.M. et al. (2017). Chaotropic heat treatment resolves native-like aggregation of a heterologously produced hyperthermostable laminarinase. *Biotechnol. J.* 12 (6): 11.
- 43 Rhodes, D.G., Bossio, R.E., and Laue, T.M. (2009). Determination of size, molecular weight, and presence of subunits. In: *Methods in Enzymology* (eds. R.R. Burgess and M.P. Deutscher), 463: 691–723. San Diego: Academic Press.
- 44 Müller, F. and van Berkel, W.J.H. (1982). A study on *p*-hydroxybenzoate hydroxylase from *Pseudomonas fluorescens*. *Eur. J. Biochem.* 128 (1): 21–27.
- 45 van Berkel, W.J.H., Benen, J.A.E., and Snoek, M.C. (1991). On the FAD-induced dimerization of apo-lipoamide dehydrogenase from *Azotobacter vinelandii* and *Pseudomonas fluorescens*. *Eur. J. Biochem.* 197 (3): 769–779.
- 46 Fraaije, M.W., van den Heuvel, R.H.H., van Berkel, W.J.H. et al. (1999). Covalent flavinylation is essential for efficient redox catalysis in vanillyl-alcohol oxidase. *J. Biol. Chem.* 274 (50): 35514–35520.
- 47 Tahallah, N., van den Heuvel, R.H.H., van den Berg, W.A.M. et al. (2002). Cofactor-dependent assembly of the flavoenzyme vanillyl-alcohol oxidase. *J. Biol. Chem.* 277 (39): 36425–36432.
- 48 Fraaije, M.W., Mattevi, A., and van Berkel, W.J.H. (1997). Mercuriation of vanillyl-alcohol oxidase from *Penicillium simplicissimum* generates inactive dimers. *FEBS Lett.* 402: 33–35.

- 49 van Berkel, W.J.H., van den Heuvel, R.H.H., Versluis, C. et al. (2000). Detection of intact megaDalton protein assemblies of vanillyl-alcohol oxidase by mass spectrometry. *Protein Sci.* 9 (3): 435–439.
- 50 Huijbers, M.M.E., van Alen, I., Wu, J.W. et al. (2018). Functional impact of the N-terminal arm of proline dehydrogenase from *Thermus thermophilus*. *Molecules* 23 (1): 184.
- 51 Huijbers, M.M.E., Wu, J.W., Westphal, A.H. et al. (2019). Dimerization of proline dehydrogenase from *Thermus thermophilus* is crucial for its thermostability. *Biotechnol. J.* 14 (5): 1800540.
- 52 Urh, M., Simpson, D., and Zhao, K. (2009). Affinity chromatography: general methods. In: *Methods in Enzymology* (eds. R.R. Burgess and M.P. Deutscher), 463: 417–438. San Diego: Academic Press.
- 53 Vos, R.M.E., Snoek, M.C., van Berkel, W.J.H. et al. (1988). Differential induction of rat hepatic glutathione S-transferase isoenzymes by hexachlorobenzene and benzyl isothiocyanate: comparison with induction by phenobarbital and 3-methylcholanthrene. *Biochem. Pharmacol.* 37 (6): 1077–1082.
- 54 Williams, R.E. and Bruce, N.C. (2002). New uses for an old enzyme: the old yellow enzyme family of flavoenzymes. *Microbiology* 148 (6): 1607–1614.
- 55 Scholtissek, A., Tischler, D., Westphal, A.H. et al. (2017). Old yellow enzyme-catalysed asymmetric hydrogenation: linking family roots with improved catalysis. *Catalysts* 7 (5): 130.
- 56 Abramovitz, A.S. and Massey, V. (1976). Purification of intact old yellow enzyme using an affinity matrix for the sole chromatographic step. *J. Biol. Chem.* 251 (17): 5321–5326.
- 57 Eweg, J.K., Müller, F., and van Berkel, W.J.H. (1982). On the enigma of old yellow enzyme's spectral properties. *Eur. J. Biochem.* 129 (2): 303–316.
- 58 Stott, K., Saito, K., Thiele, D.J. et al. (1993). Old yellow enzyme. The discovery of multiple isozymes and a family of related proteins. *J. Biol. Chem.* 268 (9): 6097–6106.
- 59 McCue, J.T. (2009). Theory and use of hydrophobic interaction chromatography in protein purification applications. In: *Methods in Enzymology* (eds. R.R. Burgess and M.P. Deutscher), 463: 405–414. San Diego: Academic Press.
- 60 Benen, J.A.E., Sánchez-Torres, P., Wagemaker, M.J.M. et al. (1998). Molecular cloning, sequencing, and heterologous expression of the *vaoA* gene from *Penicillium simplicissimum* CBS 170.90 encoding vanillyl-alcohol oxidase. *J. Biol. Chem.* 273 (14): 7865–7872.
- 61 Eppink, M.H.M., Boeren, S.A., Vervoort, J.M. et al. (1997). Purification and properties of 4-hydroxybenzoate 1-hydroxylase (decarboxylating), a novel flavin adenine dinucleotide-dependent monooxygenase from *Candida parapsilosis* CBS604. *J. Bacteriol.* 179 (21): 6680–6687.
- 62 Westphal, A.H. and de Kok, A. (1988). Lipoamide dehydrogenase from *Azotobacter vinelandii*. Molecular cloning, organization and sequence analysis of the gene. *Eur. J. Biochem.* 172 (2): 299–305.
- 63 Benen, J.A.E., van Berkel, W.J.H., van Dongen, W.M.A.M. et al. (1989). Molecular cloning and sequence determination of the *lpd* gene encoding lipoamide dehydrogenase from *Pseudomonas fluorescens*. *Microbiology* 135 (7): 1787–1797.
- 64 van Berkel, W.J.H., van den Berg, W.A.M., and Müller, F. (1988). Large-scale preparation and reconstitution of apo-flavoproteins with special reference to butyryl-CoA dehydrogenase from *Megasphaera elsdenii*. *Eur. J. Biochem.* 178 (1): 197–207.

- 65 Hefti, M.H., Vervoort, J.M., and van Berkel, W.J.H. (2003). Deflavination and reconstitution of flavoproteins: Tackling fold and function. *Eur. J. Biochem.* 270 (21): 4227–4242.
- 66 Cummings, L.J., Snyder, M.A., and Brisack, K. (2009). Protein chromatography on hydroxyapatite columns. In: *Methods in Enzymology* (eds. R.R. Burgess and M.P. Deutscher), 463: 387–404. San Diego: Academic Press.
- 67 Brondyk, W.H. (2009). Selecting an appropriate method for expressing a recombinant protein. In: *Methods in Enzymology* (eds. R.R. Burgess and M.P. Deutscher), 463: 131–147. San Diego: Academic Press.
- 68 Zerbs, S., Frank, A.M., and Collart, F.R. (2009). Bacterial systems for production of heterologous proteins. In: *Methods in Enzymology* (eds. R.R. Burgess and M.P. Deutscher), 463: 149–168. San Diego: Academic Press.
- 69 Cregg, J.M., Tolstorukov, I., Kusari, A. et al. (2009). Expression in the yeast *Pichia pastoris*. In: *Methods in Enzymology* (eds. R.R. Burgess and M.P. Deutscher), 463: 169–189. San Diego: Academic Press.
- 70 Malhotra, A. (2009). Tagging for protein expression. In: *Methods in Enzymology* (eds. R.R. Burgess and M.P. Deutscher), 463: 239–258. San Diego: Academic Press.
- 71 Block, H., Maertens, B., Spriestersbach, A. et al. (2009). Immobilized-metal affinity chromatography (IMAC): a review. In: *Methods in Enzymology* (eds. R.R. Burgess and M.P. Deutscher), 463: 439–473. San Diego: Academic Press.
- 72 Leferink, N.G.H., van den Berg, W.A.M., and van Berkel, W.J.H. (2008). L-Galactonoyl-lactone dehydrogenase from *Arabidopsis thaliana*, a flavoprotein involved in vitamin C biosynthesis. *FEBS J.* 275 (4): 713–726.
- 73 Tischler, D., Eulberg, D., Lakner, S. et al. (2009). Identification of a novel self-sufficient styrene monooxygenase from *Rhodococcus opacus* 1CP. *J. Bacteriol.* 191 (15): 4996.
- 74 Riedel, A., Heine, T., Westphal, A.H. et al. (2015). Catalytic and hydrodynamic properties of styrene monooxygenases from *Rhodococcus opacus* 1CP are modulated by cofactor binding. *AMB Express* 5 (1): 112.
- 75 Montersino, S., te Poele, E., Orru, R. et al. (2017). 3-Hydroxybenzoate 6-hydroxylase from *Rhodococcus jostii* RHA1 contains a phosphatidylinositol cofactor. *Front. Microbiol.* 8 (1110): 11.
- 76 Riedel, A., Mehnert, M., Paul, C.E. et al. (2015). Functional characterization and stability improvement of a 'thermophilic-like' ene-reductase from *Rhodococcus opacus* 1CP. *Front. Microbiol.* 6: 1073.
- 77 Mendes, S., Banha, C., Madeira, J. et al. (2016). Characterization of a bacterial pyranose 2-oxidase from *Arthrobacter siccitolerans*. *J. Mol. Catal. B* 133: S34–S43.
- 78 Heine, T., Scholtissek, A., Westphal, A.H. et al. (2017). N-terminus determines activity and specificity of styrene monooxygenase reductases. *Biochim. Biophys. Acta* 1865 (12): 1770–1780.
- 79 Ewing, T.A., van Noord, A., Paul, C.E. et al. (2018). A xylenol orange-based screening assay for the substrate specificity of flavin-dependent *para*-phenol oxidases. *Molecules* 23 (164): 18.
- 80 Nguyen, Q.T., de Gonzalo, G., Binda, C. et al. (2016). Biocatalytic properties and structural analysis of eugenol oxidase from *Rhodococcus jostii* RHA1: a versatile oxidative biocatalyst. *ChemBioChem* 17 (14): 1359–1366.

- 81 Dijkman, W.P. and Fraaije, M.W. (2014). Discovery and characterization of a 5-hydroxymethylfurfural oxidase from *Methylovorus* sp. strain MP688. *Appl. Environ. Microbiol.* 80 (3): 1082.
- 82 Ewing, T.A., Dijkman, W.P., Vervoort, J.M. et al. (2014). The oxidation of thiols by flavoprotein oxidases: a biocatalytic route to reactive thiocarbonyls. *Angew. Chem. Int. Ed.* 53 (48): 13206–13209.
- 83 Westphal, A.H., Tischler, D., Heinke, F. et al. (2018). Pyridine nucleotide coenzyme specificity of *p*-hydroxybenzoate hydroxylase and related flavoprotein monooxygenases. *Front. Microbiol.* 9 (3050): 10.
- 84 Schlager, B., Straessle, A., and Hafen, E. (2012). Use of anionic denaturing detergents to purify insoluble proteins after overexpression. *BMC Biotechnol.* 12: 95.
- 85 Ruiz-Dueñas, F.J., Ferreira, P., Martínez, M.J. et al. (2006). *In vitro* activation, purification, and characterization of *Escherichia coli* expressed aryl-alcohol oxidase, a unique H<sub>2</sub>O<sub>2</sub>-producing enzyme. *Protein Expr. Purif.* 45 (1): 191–199.
- 86 Fürst, M.J.L.J., Kerschbaumer, B., Rinnofner, C. et al. (2019). Exploring the biocatalytic potential of a self-sufficient cytochrome P450 from *Thermothelomyces thermophila*. *Adv. Synth. Catal.* 361 (11): 2487–2496.
- 87 Fürst, M.J.L.J., Martin, C., Lončar, N. et al. (2018). Experimental protocols for generating focused mutant libraries and screening for thermostable proteins. In: *Methods in Enzymology* (ed. N. Scrutton), 608: 151–187. San Diego: Academic Press.
- 88 Fürst, M.J.L.J., Boonstra, M., Bandstra, S. et al. (2019). Stabilization of cyclohexanone monooxygenase by computational and experimental library design. *Biotechnol. Bioeng.* 116 (9): 2167–2177.
- 89 Hefti, M.H., Milder, F.J., Boeren, S.A. et al. (2003). A His-tag based immobilization method for the preparation and reconstitution of apoflavoproteins. *Biochim. Biophys. Acta* 1619 (2): 139–143.
- 90 Xu, L., Zhang, D., Mu, W. et al. (2006). Reversible resolution of flavin and pterin cofactors of His-tagged *Escherichia coli* DNA photolyase. *Biochim. Biophys. Acta* 1764 (9): 1454–1461.
- 91 Li, G., Fürst, M.J.L.J., Mansouri, H.R. et al. (2017). Manipulating the stereoselectivity of the thermostable Baeyer–Villiger monooxygenase *Tm*CHMO by directed evolution. *Org. Biomol. Chem.* 15 (46): 9824–9829.
- 92 Li, G., Garcia-Borràs, M., Fürst, M.J.L.J. et al. (2018). Overriding traditional electronic effects in biocatalytic Baeyer–Villiger reactions by directed evolution. *J. Am. Chem. Soc.* 140 (33): 10464–10472.
- 93 Fürst, M.J.L.J., Romero, E., Gómez Castellanos, J.R. et al. (2018). Side-chain pruning has limited impact on substrate preference in a promiscuous enzyme. *ACS Catal.* 8 (12): 11648–11656.
- 94 Fürst, M.J.L.J., Savino, S., Dudek, H.M. et al. (2017). Polycyclic ketone monooxygenase from the thermophilic fungus *Thermothelomyces thermophila*: a structurally distinct biocatalyst for bulky substrates. *J. Am. Chem. Soc.* 139 (2): 627–630.
- 95 Heuts, D.P.H.M., van Hellemond, E.W., Janssen, D.B. et al. (2007). Discovery, characterization, and kinetic analysis of an alditol oxidase from *Streptomyces coelicolor*. *J. Biol. Chem.* 282 (28): 20283–20291.

- 96 Huijbers, M.M.E. and van Berkel, W.J.H. (2016). A more polar N-terminal helix releases MBP-tagged *Thermus thermophilus* proline dehydrogenase from tetramer-polymer self-association. *J. Mol. Catal. B* 134: 340–346.
- 97 Huijbers, M.M.E., Martínez-Júlvez, M., Westphal, A.H. et al. (2017). Proline dehydrogenase from *Thermus thermophilus* does not discriminate between FAD and FMN as cofactor. *Sci. Rep.* 7: 43880.

