

Index

a

actinide complexes of redox non-innocent ligands
 aryloxy ligands 335–338
 bipyridyl ligands 318–320
 phosphite ligands 332
 pyridine(di-imine) ligands 329–332
 pyrrole ligands 320–322
 quinone ligands 332–335
 Schiff-base ligands 324–328
 Tmtaa ligand 323–324
 Ag(PF₆) or Ag(SbF₆) 214
 amidophenolate 11, 12, 14, 23, 256–258, 325, 332
o-amidophenolate based ligands 23
 aniline 109–113, 115, 126, 134
 anilinosalens 114, 131
 aniliny radical 110
 anilinosalens 114–115
 conjugated anilines 115
 dipyrin-anilines 115
 simple anilines 111–113
 TACN-anilines 113–114
 tripodal ligand 114
 ap crown ionic salts 256
 arachidonic acid (AA) 4
 aryloxy ligands 335–338
 aziridination reaction 41, 69, 72–73

b

BD-DFT formalism 61
 benzoquinonediimine (L_{NN}) 53, 87
 1,4-benzoquinonediimines 152–162
 [(bian)⁻Yb^{II}Br]₂ 259
 bidentate iminosemiquinone radicals 111
 bidentate *o*-diaminobenzenes
 heteroleptic complexes 120–123
 homoleptic complexes 115–120
 9,10-phenanthrenediimine 123–124
 bimetallic Yb-(^tBu-dad) adducts 263
 bipyridine (bpy) 7, 53–55, 59, 76, 78–79, 82, 86–88, 94, 96, 120, 127, 159, 260, 265, 280, 286, 291, 297
 bipyridyl ligands 318–320
 bisamidinates 265
 bis(imino)acenaphthalene (bian) system 258
 bis(imino)pyridines 14, 250
 and lanthanide complexes 271–277
 bis-iminopyrrolyl ligand system 275
 bis-mesityl-1,4-diaza-1,3-butadiene 261
 B3LYP functional 59
 boratabenzenes 265
 Born–Haber thermodynamic cycle 84–85
 branched tetradentate *o*-diaminobenzene
 macrocyclic ligands 130
 planar N4 systems 128–130

- Broken Symmetry DFT (BS-DFT) 59, 65, 68
 2-(^tBu-NOH)-py ligand 277
- c**
- [calix[4]pyrrole]⁴⁺(Me₈-calix[4]pyrrole) macrocycles 322
 catecholates redox-active ligand scaffolds 23
 Ce^{IV/III} redox couple 278
 C–H bond amination, aziridination 138–139
 2-chloro-1,3-dimethyl-4,5-dihydro-1*H*-imidazolium chloride 200
 (C₉H₇)₂Yb 265–266
 (C₉H₇)Yb^{III}(pyr-CH-N-Dipp)₂ 272
 (C₉H₇)₂Yb(THF)₂ 271–273
 classical divalent lanthanides 250
 conjugated anilines 115
 π-conjugated polymers 151, 153, 169
 copper-GFA bonds 218
 copper-radical oxidases 179–180
 [Cp*₂La^{III}(Ph₂-dad-Ph₂²⁻)]⁻ [Na(Ph₂-dad-Ph₂)]⁺ ion pair 269
 [Cp*₂Yb^{III}(^tBu₂-dad)]⁻ 264
 [Cp₂Yb^{III}(^tBu₂-dad)]⁻ compound 263
 Cp*₂Yb(R-pyr-CH-N-Dipp) compounds 274
 [Cr(bpy)₃] complex 57
 cross-coupling of phenols 238–241
 cryo X-ray crystallographic techniques 59
 [Cu(BINap-SQ-BQ)]⁺ 73
 Cu^I complexes with oxidized GFA ligands 215
 Cu^{II} complexes with neutral GFA ligands 207
 Cu^{II}(L^{SQ})₂ 69, 72–74
 cytochrome c oxidase 64, 177–178
 cytochrome P450 135, 187–188
- d**
- d,π-conjugated complexes 151
 deca-arylsamarocene complex 254
 decamethylterbocene 264
 density functional theory (DFT) 10, 28, 32, 36–39, 41, 43–44, 54–63, 68–69, 72–75, 83–84, 92–93, 98–99, 113, 126, 132, 183, 185, 317, 321, 324, 326–327, 332
 “diamido” platform 132–133
 diaminobenzene platform 133–135
 1,4-diaza-1,3-butadiene 60, 261
 scaffold 262
 diazadienes 185, 250, 261–271
 dicopper complexes 212
 2,5-dihydroxy-1,4-benzoquinone (dhbq) 254
 diimino-dipyrin acyclic ligand 320
 diiron(III)–tyrosyl-radical cofactor 4
 4,4′-dimethyl-2,2′-bipyridine (^{Me}bpy) 55
 dioxolene-type ligands 199
 dioxygen reduction coupled with
 substrate dehydrogenation
 copper-radical oxidases 179–180
 H₂ generation 184–187
 O₂ reduction 180–184
 dioxygen reduction coupled with
 substrate hydroxylation
 cytochrome P450 187–188
 O₂ reduction 188–191
 dioxygen reduction to water 177–179
 Dipp₂-dad 266–270
 complexes with dysprosium 269
 dipyrin-anilines 115
 2,6-di-*tert*-butyl-4-phenylaniline 111
- e**
- effective oxidation state (EOS) analysis 88, 98
 electron paramagnetic resonance (EPR) 25–26, 59, 317
 electron reservoirs
 energy conversion
 group transfer and radical reactivity 69–74
 noble metal reactivity 64–69
 photoredox catalysis 74–80
 redox-flow batteries (RFBs) 80–84

- electron transfer coupled to spin-flips
 nitrene radical formation and transfer
 with cobalt-TAML complexes
 41–46
 nitrene radical formation and transfer
 with [Cu(NO^{iso})₂] complex 40–41
- energy conversion
 group transfer and radical reactivity
 69–74
 noble metal reactivity 64–69
 Eu²⁺/Eu³⁺ transformation 254
 exocyclic *N*-aryl radicals 109–111, 140
- f**
 [Fe^{II}(^Fdad⁻)₂] 61–63
 [Fe^{III}(^Fdad⁻)(^Fpda²⁻)] 63
- g**
 Gd^{III} and 3,5-di-*tert*-butylcatechol 251
 Gd-semiquinonate moieties 251
 [(GFA2a)(Cu^ICl₂)₂](CuICl₂)₂ 213
 Gibbs free energy 84–85, 93, 95, 97, 99
 Green formalism 3, 10
 Grimme's D3 scheme 99
 guanidino-functionalized aromatics
 (GFAs) 200–201, 203–206
- h**
 [(HBPz₃)₂Ln(tetrox³⁻)Ln(HBPz₃)₂]⁻ 254
 heteroleptic complexes 78, 93, 111,
 120–123
 hetero-ligand compound 257
 hexanuclear palladium complex 213
 homoleptic complexes 115–120
 HYSORE measurements 72
- i**
 ideal gas–rigid rotor–harmonic oscillator
 approximation 85
 iminopyridines 271–277
 iminopyrrolyl and iminopyridine
 compounds 275
 iminoquinone-based complexes 256, 258
 iminosemiquinonato 37, 40, 257
 iminosemiquinone (SQ) 256, 258
- bidentate radicals 111
 ligands 69
- in silico* description and engineering
 of redox-active ligands
 characterizing ligand redox activity
 88–92
 computing reduction potentials 84–86
 photoredox catalysis 92–98
 quinoid and bipyridine ligands 86–88
- intramolecular electron transfer (IET)
 199
- in copper complexes
 consecutive reactions 227–228
 counterligand addition 230–232
 counterligand substitution 232–235
 RIET 229
 secondary coordination sphere
 235–237
 valence tautomerism 220–227
- intramolecular interligand electron
 transfer (IIET) 255
- ion-pairing 99
 [Ir(ppy)₂(bpy)]⁺ (d⁶L⁰) 79
- j**
 Jacob's ladder of density functional theory
 54
- l**
 lanthanide compounds 249–250, 256,
 275, 286, 299
 lanthanide-dependent redox isomerism
 259
 ligand-centered redox events 76, 88, 97,
 125–126, 132
 ligand-condensed π-density values 89
 Ln-amidophenolate complexes 257
 [Ln^{II}(bian)²⁻] framework 260
 Ln-dad systems 270
 Ln-diimine based complexes 258
- m**
 macrocyclic ligands 130, 133, 321, 325
 Mayer's effective fragment orbitals (EFOs)
 88

- [M(bpy)₃]⁰ complexes 57
 2-MePh₂-dad-Me₂ 265
 mesoionic carbene (MIC) ligand 76, 96
 metal-based two-electron redox transformations 249
 metal complexes bearing redox-active ligands
 dioxygen reduction coupled with substrate dehydrogenation
 copper-radical oxidases 179–180
 H₂ generation 184–187
 O₂ reduction 180–184
 dioxygen reduction coupled with substrate hydroxylation
 cytochrome P450 187–188
 O₂ reduction 188–191
 dioxygen reduction to water 177–179
 protons and/or electrons 175
 metal-dependent metalloenzymes 175
 metal–dithiolate synthetic complexes 3
 metal–ligand interactions 53, 250
 metrical oxidation state (MOS) 23
 [M(isq)₂Ph₂] 65
 molecular spin catalysis 69
 mono-methylated Cp co-ligand 268
- n**
- N*-aryl-amido-phenolate redox-active ligands 24
N-aryl radicals
 anilinyl radicals
 anilinosalens 114–115
 conjugated anilines 115
 dipyrrin-anilines 115
 simple anilines 111–113
 TACN-anilines 113–114
 tripodal ligand 114
 bidentate *o*-diaminobenzenes
 heteroleptic complexes 120–123
 homoleptic complexes 115–120
 9,10-phenanthrenediimine 123–124
 branched tetradentate
 o-diaminobenzene
 macrocyclic ligands 130
 planar N4 systems 128
 tripodal N4 ligands 128–130
 catalytic applications
 alcohols oxidation 135–137
 C–H bond amination, aziridination 138–139
 O₂/H₂O₂ - and H₂ - activation, H₂ production 137
 exocyclic 109
 miscellaneous transformations 139–140
 Pincer ligands 124–128
 polydentate ligands, one bidentate
 diiminosemiquinone radical
 diamido platform 132–133
 diaminobenzene platform 133–135
 Salphen 131–132
 Natural Population Analysis (NPA) 89
 4N-dimensional wavefunction 54
 N-heterocycles 250, 280–299
 N-heterocyclic carbene (NHC) 14, 76, 96
 nitrene radical formation and transfer with a [Cu(NOisq)₂] complex 40–41
 nitrene radical formation and transfer with cobalt-TAML complexes 41–46
 nitroxides 250, 277–279, 300
 N,N'-bis(4'-dimethylaminophenyl)-1,4-benzoquinonediimine 153
 N,N'-diphenyl-*p*-phenylenediamine 156
 non-innocence 6–7, 9, 10, 24, 80, 86, 88, 93, 115, 127
 N-phenyltetrahydroisoquinoline 169
 nuclear magnetic resonance (NMR) 27–28, 31, 38–39, 68–69, 134, 154, 210, 217, 232, 255, 298, 324, 328–329
- p**
- pairwise electron transfer (PET) 65, 67, 74
 PANI (emeraldine salt) 164
 Pd-catalyzed *N*-arylation reactions 53
 (PDI)FeN₂ 64, 68
 (*R,R*)-13syn 160

(PDI)Fe(N₂)₂ and (PDI)FeN₂ 68
 Pd-polyaniline-catalyzed Wacker oxidation 163
 Ph₂-dad-Me₂ adduct 264
 Ph₂-dad-Ph₂ ligand 267
 9,10-phenanthrenediimine 123–124
 phenantren-*o*-iminoquinone system 255
o-phenylenediamine 60, 185–187, 250–262
 POT 157–158
 phosphite ligands 332
 photoredox catalysis 13, 54, 64, 74–80, 92–98
 photo-redox catalysts 74–80, 92, 93, 96, 99
 photo-redox cycle 78, 80, 93, 95, 97
 pincer ligands 37, 118, 124–128, 176, 181, 184, 185, 190
 ping-pong mechanism 179, 180
 planar N₄ systems 128–130
 PMAS/Au NPs catalyst 168, 169
 PMAS-catalyzed oxidation 164
 poly(3-heptylpyrrole) 153
 poly(2-methoxyaniline-5-sulfonic acid) 163
 poly(*o*-toluidine) (POT) 152
 polyaniline 151
 CuCl₂-catalyzed oxidation reactions 163
 iron oxide NPs 166–167
 Pd NPs 165
 polydentate ligands, one bidentate
 diiminosemiquinone radical
 diamido platform 132–133
 diaminobenzene platform 133–135
 Salphen 131–132
 potential energy surface (PES) 85, 206
 prostaglandin H synthase (PGHS) 4
 proton affinity 205
 pyridine bis-iminophosphorane ligands 277
 pyridine(di-imine) ligands 329–332
 pyridine sidearms 96
 pyrrole ligands 176, 189, 320–322

q

quasi-restricted orbitals (QROs) 78–80, 98
 quinone derivatives 59
 quinone, iminoquinone and
o-phenylenediamine-based complexes 250–262
 quinone ligands 11, 13, 14, 178, 257, 332–335

r

redox-active azines (RAAs) 200, 201, 227
 redox-active π -conjugated ligands 152, 169
 redox-active guanidine ligands
 in coordination chemistry 199
 of IET in copper complexes 219–239
 survey of 206–220
 cross-coupling of phenols 238–241
 intramolecular electron transfer 219–200
 properties and reactivity 200–206
 redox-active ligand
 advantage 21
 anatomy of
 basic/minimal features 7–8
 biological systems radicals 4
 C–C bond formation 14–15
 metal complexes with dithiolene ligands 6–7
 modes of action 8–9
 non-innocent, cooperative, electro-active or redox-active 10–12
 open-shell reactivity radical formation 13–14
 reactivity at ligand 12–13
 redox cofactors 4–6
 spectroscopy and theory 9–10
 computational perspective
 electron reservoirs 63–84
 electronic structure determination through DFT and spectroscopy 55–63

- redox-active ligand (*contd.*)
- in silico description and engineering of 84–98
 - computational studies 28, 53–55, 98, 99, 290, 330
 - diazadienes 250, 262–271
 - effective magnetic moment and spin state 27
 - electron paramagnetic resonance 25–26, 59, 317
 - iminopyridines and
 - bis(imino)pyridines 271–277
 - impact 22, 75, 252, 267, 269, 274, 292
 - mechanistic studies
 - computational methods 32
 - Hammett analysis 31
 - kinetic analysis 29–30
 - kinetic isotope effect 30
 - trapping and poisoning 31–32
 - in metal complexes 249
 - metal-coordinated substrates 22
 - modes of action 22
 - N*-heterocycles 280–299
 - nitrene transfer
 - electronic structures of 33–34
 - electron transfer coupled to spin-flips 40–46
 - ligand-to-substrate single-electron transfer 37–38
 - metal-to-substrate single-electron transfer 34–37
 - spin-flip assisted reactions 39–40
 - nitroxides 277–279
 - nuclear magnetic resonance 27
 - quinone, iminoquinone and
 - o*-phenylenediamine-based complexes 250–262
 - single-crystal X-ray diffraction 23
 - UV-vis & IR absorption spectroscopy and (spectro)-electrochemistry 28
 - X-ray absorption spectroscopy 23–25
 - redox-active polyanilines
 - anti*-configuration 154, 160
 - conjugated homobimetallic palladium(II) complex 153
 - conjugated oligomeric nickel complex 157
 - conjugated tetranuclear macrocycle 157
 - cross-linked conjugated complex 153
 - hetero-bimetallic palladium-nickel complex 157
 - N,N'*-diphenyl-*p*-phenylenediamine 156
 - palladium(II) complex $L^1Pd(MeCN)$ 161
 - PD derivative and the QD derivative 155
 - redox-active ferrocenyl groups and PD bridging spacer 155
 - redox interconversion 160
 - [(*S,S*)- $L^4Pd(MeCN)$] or [(*R,R*)- $L^4Pd(MeCN)$] 157–159
 - syn* configuration 155
 - transition metals 162–169
 - vanadium 162
 - redox-flow batteries (RFBs) 64, 80–84
 - redox-induced electron transfer (RIET) 12, 200, 220, 229, 241
 - reduction potentials 75–78, 83–85, 87, 92–95, 99, 283, 287, 293
 - reliability index 88, 89
- S**
- Salphen 131–132
 - Schiff-base ligands 324–328
 - simple anilines 111–113
 - single-crystal X-ray diffraction (SC-XRD) 23, 38, 41
 - single-walled carbon nanotubes (SWCNTs) 203
 - singly occupied molecular orbitals (SOMO) 26, 34, 58, 69, 74, 80, 129, 324
 - [$Sm^{III}(Dipp_2-dad-Me_2)^2-(\mu-OMe)_2$] 270
 - Solvation Model Based on Density (SMD) 85, 94, 97, 99
 - spectator or innocent ligands 11

spectroscopic methods 55, 59, 63, 318
 spin-flip assisted reactions 39–40
 SQUID measurements 210, 218, 220
 Sterically Induced Reduction 253, 266

t

TACN-anilines 113–114
 2,2',6',2'-terpyridine (tpy) as ligands 55
 tetrakis(imino)pyracene 261
 1,2,4,5-tetrakis(tetramethylguanidino)-
 benzol 202
 tetraoxolene-based compounds 254
 tetraoxolene ligands 199, 229
 tetravalent complex 278
 theoretical square schemes 91
 $[\text{Ti}(\text{bpy})_3]^0$ 57, 58
 time-dependent density functional theory
 (TD-DFT) approaches 41, 69, 75,
 92, 123
 Tmtaa ligand 323–324
p-toluenesulfonic acid-doped polyaniline
 166, 167
 transition metal complexes 21, 23, 24,
 53, 55, 63, 64, 86, 98, 99, 116, 151,
 206
 tripodal ligand 114, 335
 2,4,6-tri-*tert*-butylaniline 111
 trivalent neodymium system 275
 tropolonate ligand 252

u

unpaired electron transfers (UET)
 66
 UV-vis & IR absorption spectroscopy and
 (spectro-)electrochemistry 28

v

valence tautomerism (VT) 12, 114, 136,
 199, 220–227, 284
 vanadium complex $[\text{V}(\text{bpy})_3]^0$ 59

x

X-ray absorption spectroscopy (XAS)
 23–25, 126, 130, 328

y

Yb-bian Cl-bridged system 259
 Yb-dad based complexes 264
 $\text{Yb}^{\text{III}}(\text{pyr-CH-N-Dipp})_3$ 272, 274

z

zero-point-energy (ZPE) 30, 85, 94
 correction 85
 zerovalent lanthanides 252
 $[\text{Zr}(\text{isq})_2\text{Ph}_2]$ 64
 $\text{Zr}^{\text{IV}}(\text{isq})_2$ system 65
 $\text{Zr}(\text{pda})_2$ ($\text{pda}^{2-} = N,N'$ -bis(neo-pentyl)-
 ortho-phenylene-diamide) 67
 $\text{ZrX}_2(\text{disq})_2$ 68

