

Index

a

AB-type monomers 63, 223
 all-optical logic based on polariton condensates 39–43
 amidinium-carboxylate salt 268, 270
 anisotropic structural fibers 328, 330
 annulation reaction 4, 103, 112, 115, 219, 221
 anthracene-decked trimeric porphyrin array 125
 antibunched photons 29
 anti conformer 305, 306
 atomic force microscopy (AFM) 70, 100–102, 104–106, 108–109, 142, 143, 157, 234, 237, 241, 250, 251, 268–271, 310–312, 331, 332

b

α -band of the absorption spectrum 21
 benzotriptycene monomer 188, 189
 benzotriptycene PIM 188
 biotechnology 313–318
 block ladderphanes 260
 Bose–Einstein condensate (BEC) 14, 34, 37–39, 44–46
 branched DNA 309–311
 bromoarenes 2, 223–224
 butadiyne-bridged porphyrin oligomers 125, 126

c

CANAL-Me-iPr 225–227
 catalytic Arene-Norbornene Annulation (CANAL) polymerization 219
 from Catellani reaction 220–222
 fluorene polymers 226
 rigid kinked ladder polymers 222–226
 Catellani reaction 220–222
 caterpillar track complexes 133
 chlorophylls 121–122
 chromium metal-string complexes 358–360
 chromophores, test for 29

cobalamin 121
 Complete Active Space Perturbation Theory (CASPT2) method 359
 conjugated hydrocarbon ladder polymers 21–22
 conjugated ladder polymers (cLPs) 97
 in-situ reaction 110
 solution-dispersed nanoparticles 108–110
 solution-processing
 end group modification 101, 102
 Lewis acids 100–101
 non-planar backbone 105–108
 protic acid 99–100
 side chain modification 102–105
 structural design 101
 conjugated polymers (CP), defined 13
 controlled spatial confinement of exciton polaritons 43–48
 conventional cryogenic single-molecule spectroscopy 27
 covalent organic framework (COF) 123, 139, 179, 242
 cross-coupling polymerization 219
 cytosine-rich four stranded DNA 308

d

daisy chaining 40–41
 dendritic multiporphyrin arrays 135, 151–152, 155
 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) 16, 126
 DNA
 hybridization and dissociation 306–307
 nano assembly 310
 nanomaterials 311–313
 non-cannonical structures 307
 branched 309–310
 cytosine-rich four stranded 308, 309
 G-quadruplex 307–308
 triple-stranded 307
 nucleobase conformation 305–306

- DNA (*contd.*)
- rod-like 1D wire 310–311
 - stacking interactions 304–305
 - structure 302
 - sugar packing 304
 - synthesis 302–304
- DNA Origami 301, 311–313, 319
- double-helix polymers 323
- concentric giant vesicle formation 331–333
 - with functional groups 325
 - liquid crystal gels
 - anisotropic materials 327–330
 - polymorphism 330–331
 - pendant oligomer 335–336
 - polymethylene compounds
 - synthesis and properties 333–335
- double-stranded DNA 248, 301, 304, 306, 310
- e**
- effective conjugation length 15, 27
 - electrocatalytic hydrogen production 166, 167
 - 16- π -electron-macrocyclic metal complex 255
 - EMACs building blocks 370
 - ethylene-bridged phenylene ladder polymers 21–24
 - ethynylhelicene oligomers 326–327, 331, 337, 340
 - exciton polaritons 34
 - condensation in planar microcavities 36
 - in microcavities 35–36
- f**
- Fabry-Pérot microcavity 40, 46
 - ferrocene linkers 250–253
 - field-effect transistors (FETs) 60, 85–88, 100, 111–112, 291
 - Förster-type intramolecular energy transfer 27
 - Frenkel excitons 34, 39–40
 - functionality transfer reaction 316–318
- g**
- gas separation, in PIMs 201–206
 - gas storage 182, 206, 210
 - Gaussian defect system 46
 - geometrically-confined monomers, pericyclic reactions 219
 - G-quadruplex DNA 307–308
 - graphene nanoribbons (GNRs)
 - device integration 83–88
 - electronic properties 59
 - magnetic properties 79–83
 - non-planarity and chirality 77–79
 - on-surface synthesis 69–77
 - pathways 60
 - solution-based synthesis 62–69
 - spin bearing 79–83
- h**
- heptachromium 359
 - hetero-double-helix oligomers 326–328, 330–332, 337, 342, 344
 - heteronuclear metal-string complexes (HMSCs) 351, 373
 - $M_A M_A M_B$ 377–380
 - M_A – M_B distances 376
 - $M_A M_B M_C$ 380–381
 - synthetic strategies 374–375
 - high-resolution atomic force microscopy (AFM) 234, 310
 - high-resolution spectroscopy of LPPP 25–26
 - high resolution-transmission electron microscopic (HR-TEM) 259
 - H_2 napany 366
 - homo-double-helix oligomer, inverse thermoresponse 337–340
 - Hopfield coefficients 36
 - hydrophilic double-helix oligomers 336–344
 - liquid water 336–337
- i**
- i-motif 308–309
 - interrupting sites 314, 315
 - inverse thermoresponse, in aqueous solvents 340
- l**
- ladderphanes 248
 - array 274
 - chemical reaction 272–275
 - cyclic 274–275
 - dielectric properties 278
 - physical properties 275–278
 - ladder polymers 59, 222–226
 - strategic considerations 234–237
 - synthetic aspects 232
 - ladder polysiloxanes (LPSs) 285, 301
 - preparation 286–290
 - hydrolysis–condensation procedures 286–288
 - supramolecular architecture-directed confined polymerization 289–290
 - structure 286
 - ladder polysilsesquioxanes (LPSQs) 285
 - applications 291
 - coating 291
 - composite materials 294–295
 - electrochromic and electrofluorochromic bifunctional materials 292–293
 - fabrication 295–296
 - LED encapsulants 291–292
 - self-healing polymeric materials 294
 - supermolecular structures 296
 - ladder-type poly(*para*-phenylenes) (LPPPs) 24
 - high-resolution spectroscopy of 25–26

- as single-photon source 27–33
 - type ladder polymer 15–19
 - Laguerre–Gaussian (LG n l) modes 44–45
 - Lewis acids 100–101
 - Lieb lattices 44, 46–48
 - light-harvesting antenna complexes (LHs) 122
 - linear multiporphyrin arrays 123
 - linear oligomeric porphyrin arrays 126
 - linear polymers 1, 5, 123, 125, 219, 227, 231–232, 275, 401
 - liquid crystal gels (LCG)
 - anisotropic materials 327–330
 - polymorphism 330–331
 - liquid water 336–337, 344
 - lower (LPB) and an upper polariton (UPB) branch 35–36
 - LPPPPy ladder polymers 18
- m**
- macrocyclic metal complexes 253–255
 - MeLBTDTTPP 16–18
 - MeLPPP 14–19, 21, 28, 34–37, 39–40, 44–45, 47, 49, 50
 - melting temperature 306
 - metal-string complex 351
 - building blocks 371
 - chromium 358–360
 - cobalt 355–358
 - conductance 387
 - nickel 352–355
 - physical properties 351
 - principle 363
 - ruthenium, rhodium, and iron 360–363
 - stereoisomers 384
 - synthesis 351
 - triiron and tetrairon 362
 - metathesis cyclopolymerization (MCP) 258–260
 - methanesulfonic acid (MSA) 99–100, 108, 110, 112
 - Michelson interferometry technique 47
 - microporosity, and stability 226, 227
 - monomers, examples 234–235
 - mononuclear metal complexes 363
 - multiporphyrin arrays
 - biological systems 122
 - constructed cages 143–147
 - function and applications
 - chemical adsorption and separation 159–162
 - diverse chemical reactions 162–170
 - host-guest chemistry and supramolecular assemblies 153–159
 - natural photosynthesis 148–153
 - linear and ladder shapes 123–128
 - ring and tube shapes 128–134
 - rotaxanes 147–148
 - spherical shapes 134–137
 - structure variations and synthetic strategies 123–148
 - two-dimensional sheet-like shapes 137–143
 - multiwalled carbon nanotubes (MWNTs) 295–296
- n**
- nanofabricated 2D Lieb lattice 46
 - nanographenes (NGs) 61
 - nature photosynthesis 148–153, 170
 - N*-bromosuccinimide (NBS) 209, 223
 - non-covalent approach 143
 - non-local density functional theory (NLDFT) 182
 - norbornadiene (NBD) 3, 221–224
 - norbornene 220, 223, 248, 250, 255, 265, 267, 275
 - nucleation 306
- o**
- octanickel metal strings 367
 - oligomers, defined 323
 - oligonaphthyridylamine ligands 367
 - oligophenylenes 61
 - Onsager theory 326
 - optoelectronic applications of aromatic ladder polymers 24–33
 - organic field effect transistors (OFETs) 100, 105, 232
 - organic light emitting diodes (OLEDs) 14, 25, 232
 - organic molecules of intrinsic microporosity (OMIMs) 182
 - organo-bridged ladder polysiloxanes (OLPSSs) 285, 286, 290
- p**
- para-phenylene building block 16, 18
 - pendant ethynylhelicene oligomers 326
 - pentachromium 359, 360
 - phenyl- C_{61} -butyric acid methyl ester (PCBM) 24
 - phosphoramidite method 302
 - photon coincidence ratio 31–33
 - physical aging 183, 205, 225
 - planar aromatic linkers 253, 256, 275
 - planarized poly(*para*-phenylene) ladder polymer (LPPP) 15–18, 20, 23–27, 31–34, 103, 104
 - poly(styrenesulfonic acid) (PSSA) 99, 100
 - polyacetylene-based Ladderphanes 258
 - charged species 260
 - synthesis 258–259
 - topochemical reaction 260
 - polybenzodioxanes 184–189, 191, 193–195, 197, 205, 209, 210

- polycyclic aromatic hydrocarbons (PAHs) 60, 61
- polymer processing 97
- polymeric backbones 247, 248, 257–258, 272
- polymerization mechanism 237–241
- polymerization-graphitization 61, 62, 67, 75
- polymers of intrinsic microporosity (PIMs)
- application
 - anion and cation exchange and energy 209–210
 - catalysis and electrochemistry 206–208
 - for pervaporation and nanofiltration 208–209
 - gas separation 201–206
 - gas storage 206
 - concept 179
 - ladder co-polymers 190–192
 - PIM-1 183–185
 - fluorinated monomers 189–190
 - modification 185–188
 - physical properties 184
 - structure 186
 - porosity 181–182
 - TB-PIMs 193–200
 - quaternisation and ring opening 198–200
 - thermal stability 182–183
- polynorbornene-based single strand polymers 257
- polynorbornene-based symmetric ladderphanes 248–257
- polypentaphene ladder polymers 19–21
- polyphosphoric acid (PPA) 99
- porous polymers 159, 162, 163, 168, 169, 182
- porphyrin rings 121
- porphyrins 121, 123, 148, 153, 154, 162, 164, 170, 253, 296
- primer on exciton polaritons in microcavities 35–36
- protic acid 99–101
- pseudo-dC derivatives 315–316
- pyrazinoquinoxaline-fused diporphyrin 125
- pyridylamino units 367
- r**
- reactive packing 234–237
- ring opening metathesis polymerization (ROMP) 219, 248–250, 257, 260, 264, 265, 267, 273, 275
- RNA modification 316–318
- room-temperature polariton condensation 46, 49
- s**
- s*- and *p*-bands 44, 47
- scanning tunneling microscopic (STM) 67, 70, 72–78, 81, 82, 87, 110–112, 129, 130, 242, 250–253, 256, 263, 265, 267, 272, 273, 278
- seeding 40, 41
- self-healing polymeric materials 293–296
- shape-persistent monomers 143, 234, 235, 401
- single-chromophore fluorescence spectra 27
- single-molecule photon antibunching 32, 33
- single-photon source, LPPP 27–33
- single-stranded precursor polymer 1, 232, 233
- solid-liquid transition 335
- Spirobisfluorene-PIM (PIM-SBF) 187, 205
- strong-coupling regime 33–35
- Suzuki-type polycondensation 15
- syn conformer 305, 306
- t**
- telomers 310
- tetraarylethylenes (TAE) 275, 277
- thermodynamic parameters 306, 311, 334, 338, 340, 341
- thermoreponse 326, 337–342, 344
- three dimensional organic linkers 255–257
- three-dimensional protein matrix 134
- tip-enhanced Raman spectroscopy (TERS) 234, 237
- Tröger's base PIMs (TB-PIMs) 193–200
- trinickel strings 352
- triple helix DNA 308, 314
- triple-stranded DNA 307
- 2D polymer 7, 231–243
- u**
- unsymmetric ladderphanes 261
 - polycyclobutene-based 267–272
 - polynorbornene-based 262–264
 - sequential polymerization 265–272
- v**
- Vinylene-bridged phenylene ladder polymers 14, 19–21
- Vitamin B12 121
- w**
- Wannier–Mott-type excitonic transitions 34
- water-soluble hetero-double-helix oligomers 342
- W-shaped nucleoside analogues (WNAs) 315
- x**
- X-ray diffraction (XRD) 104, 234, 235, 270, 288, 297, 359, 374, 384
- z**
- Z-DNA 304–306
- zero-dimensional (0D) confined structure 44
- zero-dimensional cavity 46
- zero-phonon line 27
- Ziegler–Natta polymerization 219