

Index

a

- AFM metallic state (A-type AFM) 28
- ambipolar cuprate 28–29
- anisotropic plasmons, 1T'-phase 2D-TMDs
- one-dimensional plasmon 110–112
- 1H–1T' phase transition 116–117
- 1T'-phase MoS₂/Au monolayer 112–115
- sample preparation 107
- WSe₂/Au 108–110
- antiferromagnetic (AFM) 23, 25, 28, 34, 36, 152–153, 155–158, 160

b

- Bardeen, Cooper, Schrieffer (BCS) theory 19
- Bethe–Salpeter equation 100, 137
- bond disproportionation (BD) 156, 160
- Brewster angle microscopy 3, 8, 169
- Brillouin zone 62, 65, 69, 71, 76, 91, 93, 101, 116, 133–134, 140, 151

c

- charge–density–wave (CDW) 34, 61, 106
- charge-transfer insulator 24
- chemical vapor deposition (CVD) 60, 68, 137
- colossal magnetoresistance (CMR) 19, 26–36
- condensed-matter physics 19, 20, 26, 68, 87, 141
- conduction band minimum (CBM) 62
- Cooper pairs 19, 107, 111
- Coulomb interactions 23, 28, 65, 87, 111
- crystal field splitting 23, 26, 151
- crystal structures
 - arising electronic structures
 - indirect-to-direct bandgap transition 61–62
 - 1T'-phase electronic band structure 63–64
 - spin–orbit splitting 62–63
 - excitons 64–67
 - 1H–1T' phase energetics 61
 - 1H-phase 60
 - 1T and 1T'-phase 61
- cuprates
 - crystalline structure 20
 - electronic structure 22–26

- cuprates (*contd.*)
- La_{1-x}Sr_xCuO₄ 20
 - Nd_{2-x}Ce_xCuO₄ 21–22
 - thin film on oxide substrate
37–40
 - YBa₂Cu₃O_{7-δ} 20
- Czerny–Turner scanning monochromator 8
- d**
- density functional theory (DFT)
69, 160
- deuterated triglycine sulfate (DTGS) **f**
detector 9
- dielectric models 12
- Drude–Lorentz model
13–17
 - Sellmeier and Cauchy models
17
- distinct peaks label 91
- Drude–Lorentz model
13–18
- DyScO₃ (DSO) 45–47
- e**
- electromagnetic waves 2, 3, 15
- electron–electron (e–e) interactions
69, 71, 100, 138, 141
- electron energy loss spectroscopy
(EELS) 41
- electron–hole (e–h) interactions
100, 137, 138
- electronic structures
- indirect-to-direct bandgap
transition 61–62
 - 1T'-phase electronic band
structure 63–64
 - spin–orbit splitting 62–63
- ellipsometric measurements 3,
5–7
- excitons 87
- binding energy 105
 - in LSCO/STO film 43–45
 - resonant excitons in graphene
137–138
 - and trions 67
 - 2D-TMD 65–67
 - Wannier–Mott and resonant
100
- f**
- Fermi level 48, 67, 79, 92, 107, 109,
135, 138, 140
- Fourier transform infrared (FTIR)
spectroscopy 9, 69
- Fresnel coefficients 2–3, 135
- full-width-half-maximum (FWHM)
83
- g**
- Goldschmidt tolerance factor
150
- graphene 150
- electronic band structure of
132–134
 - measurement and data analysis
135–137
 - optoelectronic properties of
134–135
 - resonant excitons in 137–138
 - spectroscopic ellipsometry study
135–137
 - substrate-induced manipulation
138–141
- graphene on STO substrate (G/STO)
139

h

- Hall effect 24, 28, 29
 Hartree–Fock theory 24, 35,
 36
 high-energy resonant exciton
 demonstrated via GW–BSE
 99–100
 high transition temperature
 superconducting cuprates
 19
 crystalline structure 20–22
 electronic structure 22–26
 Hund’s interaction 26

i

- indirect-to-direct bandgap
 transition 61–62
 IR-VASE spectroscopic ellipsometer
 9–10

j

- Jahn–Teller effect 28
 Jones matrix 4–6
 Jones vectors 4, 6

k

- Kramers–Kronig transformation
 47

l

- $\text{La}_{0.7}\text{Sr}_{0.3}\text{MnO}_3$ (LSMO) 26, 45
 $\text{La}_{1-x}\text{Sr}_x\text{CuO}_4$ 20, 21
 $\text{La}_{1-x}\text{Sr}_x\text{MnO}_3$ (LSMO) 26
 $\text{La}_{1.85}\text{Sr}_{0.15}\text{CuO}_4$ (LSCO) thin-films
 37
 $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ 23
 LaAlO_3 (LAO) 29, 170

- $(\text{LaAlO}_3)_{0.3}(\text{Sr}_2\text{AlTaO}_6)_{0.7}$ (LSAT)
 37
 La–Ba–Cu–O system 19
 LaMnO_3 26, 27, 170
 Langmuir monolayers 169–170
 local-field effect (LFE) 40
 localized transitions 90, 95
 Lorentzian–Gaussian lineshape
 93
 Lorentz model 12, 14–17
 loss function (LF) 32, 34, 38–41,
 108, 109
 lower Hubbard band (LHB) 23, 24
 LSCO/STO film
 high energy excitons 43
 low and high-energy plasmons
 40

m

- Maxwell’s equations 2
 metal–insulator transition (MIT)
 151–153, 158, 167
 metal-oxide semiconductor
 field-effect transistor
 (MOSFET) scaling 171
 mid-infrared (MIR) 32, 34, 69, 71,
 72, 76, 81, 107, 109–112,
 115–117, 135
 monolayer- MoS_2
 atomic layers 68
 1H–1T’ phase transition 72
 1T’-phase 72
 optical features 70–71
 monolayer- MoS_2/Cu 1H–1T’ phase
 transition
 optical characterization 75–76
 photoemission spectroscopic
 characterization 77–79

- monolayer-MoS₂/Cu 1H–1T' phase transition (*contd.*)
 - Raman and PL characterization 76–77
 - WSe₂/Au 80–81
- monolayer-WSe₂,
 - three-dimensional resonant exciton 87, 103
 - exciton peaks 92–94
 - optical characterization 89–92
 - sample growth 88–89
 - ϵ_2 spectra 94–95
 - Mott–Hubbard insulator 23, 26
 - Mott insulator 23, 26, 28, 142
 - Mueller matrix ellipsometry (MME) 167
 - Mueller matrix imaging
 - ellipsometer (MMIE) 167
- n**
- narrow nonbonding (NB) bands 23
- n*-butyl lithium (*n*-BuLi) 61, 68, 106
- Nd_{2-x-z}Ce_xSr_zCuO₄ 23
- Nd_{2-x}Ce_xCuO₄ 21–23
- Néel temperature 23, 25, 28
- Ni3d-orbital physics 150–151
- nickelate systems
 - crystal structure 149–150
 - electronic structures 150–151
 - metal-insulator transitions
 - magnetic phases 152–153
 - optical characterization via spectroscopic ellipsometry 153–159
 - Ni3d-orbital physics 150–151
- nickelate thin-films
 - epitaxial effects on the electronic correlations 154–155
 - long-range spin ordering 155–159
- nudged elastic band (NEB) method 72, 86
- o**
- 1H–1T' phase transition 116–117
 - first-principles 72–75
 - optical characterization 75–76
 - phase transition yield 84
 - photoemission spectroscopic characterization 77–79
 - Raman and PL characterization 76–77
- WSe₂/Au
 - optical characterization 80–81
 - photoluminescence and photoemission characterization 81–84
- 1T'-phase 2D-TMDs 61, 63, 64, 70–72, 75, 85, 87, 106–110, 115–117
- 1T'-phase electronic band structure 63–64
- 1T'-phase MoS₂/Au monolayer 112–115
- p**
- p-and s-polarized lights 2–3, 11
- photoemission spectroscopies (PES) 69, 79, 135
- photoluminescence (PL)
 - high-energy characterization 97–99
 - and photoemission WSe₂/Au 81–84

- power-dependent spectroscopic study 99
- polymethyl methacrylate (PMMA) 70, 107
- r**
- random-phase approximation 100
- rare earth nickelates 167
- reflectometry probing method 169
- resonant valence bond (RVB) theory 28
- rotating analyzer ellipsometer (RAE) 6–7
- Rydberg exciton states 65
- s**
- Sellmeier and Cauchy models 17
- single layer graphene systems
- crystal and electronic structure of 131–134
 - many-body effects 138–141
 - optoelectronic properties of 134–135
 - resonant excitons in 137–138
 - spectroscopic ellipsometry of 135–137
- Snell's law 3
- ϵ_2 spectra 38–41, 43, 45, 70, 89, 90, 94–95
- spectroscopic ellipsometry (SE)
- advantages and disadvantages 10
 - anisotropic materials 173
 - biological materials and medicines 173
 - dielectric models 12–13
 - Drude–Lorentz model 13–17
 - Sellmeier–Cauchy models 17
 - ellipsometric measurements 5–6
- features of 10
- Fresnel coefficients 2–3
- IR-VASE spectroscopic ellipsometer 9–10
- Langmuir monolayers 169–170
- metamaterial research 172
- micro devices and integrated circuits 171–172
- Mueller matrix imaging techniques 167–169
- multilayered system 11–12
- organic electronics 172
- p-and s-polarized lights 2–3
- process control analysis 173–174
- two-dimensional interfaces 170–171
- VASE spectroscopic ellipsometer 8
- YLBLCO 31–32
- sp^2 hybridization 131
- spin-orbit splitting 62–63
- spin-valley coupling 63
- $SrTiO_3$ (STO) 37, 139
- Stokes parameters 4, 7
- Stokes vector 5, 168
- strongly correlated systems
- ambipolar cuprate thin-film systems 28
 - colossal magnetoresistance manganites 26–36
 - cuprate thin film on oxide substrate 37–40
 - high-transition-temperature superconducting cuprates 19–26
 - Jahn–Teller splitting energy control 45–49
 - plasmon and high-energy exciton excitations 40–45

- superconducting phase (SC) 25, 26, 29
- t**
- three-dimensional resonant exciton band structure 101–103
high-energy photoluminescence characterization 97–99
high-energy resonant exciton demonstrated via GW-BSE 99–100
monolayer-WSe₂ 87–88, 103
exciton peaks 92–94
optical characterization 89–92
sample growth 88–89
 ϵ_2 spectra 94–95
power-dependent photoluminescence spectroscopic study 99
SOC 95–96
thickness dependence 103–105
Wannier–Mott and resonant excitons 100
trions 67, 92–94, 105
two-dimensional transition metal dichalcogenides (2D-TMDs)
anisotropic plasmon excitations in quasi-metallic 107–110
crystal structures arising electronic structures 61
1H–1T' phase energetics 61
1H-phase 60
1T and 1T'-phase 61
excitons 64–67
high-energy photoluminescence characterization 97–99
- investigative objectives 67–68
monolayer-MoS₂
atomic layers 68
1H–1T' phase transition 71–74
1T'-phase 71
optical features 70–71
monolayer-WSe₂ 87–88
trions 67
- u**
- ultraviolet–visible spectroscopy 1
upper Hubbard band (UHB) 23, 24, 36
- v**
- valence band maximum (VBM) 62, 79, 101
variable-angle spectroscopic ellipsometer (VASE) 8
- w**
- Wannier excitons 65, 75, 81
Wannier–Mott excitons 65, 87, 99, 100, 137
Weyl semimetals 68, 106
WSe₂/Au
anisotropic plasmons 1T'-phase 2D-TMDs 107
1H–1T' phase transition 79–85
- x**
- X-ray absorption spectroscopy (XAS) 28, 44, 47, 170
- y**
- Y_{0.38}La_{0.62}(Ba_{0.82}La_{0.18})₂Cu₃O_y 29–31, 36

$\text{Y}_{1-z}\text{La}_z(\text{Ba}_{1-x}\text{La}_x)_2\text{Cu}_3\text{O}_y$
(YLBLCO) 29, 31, 32

$\text{YBa}_2\text{Cu}_3\text{O}_6$ 22

$\text{YBa}_2\text{Cu}_3\text{O}_{7-\delta}$ 20

$\text{YBa}_2\text{Cu}_3\text{O}_7$ 22

z

Zaanen–Sawatzky–Allen scheme

24

Zhang–Rice singlet band 24

