

Index

a

ab initio molecular dynamics 68, 86, 355
ab initio potential energy surface (PESs) 303
 acetonitrile (ACN) 132, 134, 139, 233, 244, 257, 266, 268–270, 273–283, 285, 286, 492
 2-acetyl-1,8-dihydroxy-3,6-dimethylnaphthalene 194, 199
 adenine 41, 50, 51, 184
 adenine crystal 50
 adiabatic ionization energy 316, 325
 Aib foldamer 490, 495, 499, 500
 alcohol-water systems
 alcohol concentration and temperature 464–470
 importance of 455
 literature review of 457–462
 spectral decomposition and probes for characterization 456–464
 alleno-acetylenic cage (AAC) 511–513
 alternating least squares (ALS) method 417, 418, 428
 amide-appended corannulenes 511
 γ -aminobutyric acid 213
 aminocyclohexanecarboxylic acid (ACHC) 480, 481, 492, 502
 aminoisobutyric acid (Aib) 489, 490, 492, 494–497, 499–501
 anions of ionic liquids 124
 anomalous Ubbelohde effect 356
 aplanar proton-transfer mechanism 107
 apolar gases 448

ATR-FUV spectrometer 409, 411, 428
 attendant protium-deuterium isotopic ratio 89
 attenuated total reflection (ATR) spectroscopy technique 128
 attractive or repulsive interactions 488

b

backbone CONH(D) isotope effects 186
 Badger-Bauer correlation 380
 barrier-penetration phenomena 90
 basis set superposition error (BSSE) 377
 Beer-Lambert law 127, 445
 benzene dimer 317
 benzoic acid (BA) 219–225, 360–362
 bifunctional 2-naphthols 285–288
 binding energy of H-bonded complexes 314–331
 Birge–Spomer extrapolation method 296
 Birge–Spomer (BS) extrapolation technique 325
 bis(2-quinolylmethyl)(2-pyridylmethyl) amine (BQPA)-ligated copper complex 495
 2,6-bis(*N*-imidazolidin-2-onyl)pyridines 488
 blueshifted H-bonds 345
 bond strength 87, 89–91, 95, 96, 107, 173, 178, 184, 376, 377, 380, 383–387, 441, 452–454, 463, 479, 480, 483
 Born–Oppenheimer approximation 47, 175

- Born–Oppenheimer Molecular Dynamics spectroscopic 41
- Brønsted-type PT reactions 261
- bulk water
- OH (OD)-stretching mode 438
 - temperature-dependent infrared spectra 439–441
- C**
- campestarenes 508, 510
- canonical tunneling 92
- carboxylic acid dimers ignoring Fermi resonances
- gaseous acrylic and propynoic acids 15, 17
 - gaseous and liquid acetic acid dimers 14–16
- carboxylic acids account Fermi resonances
- combined crystalline oxindole acid dimers 31
 - crystalline adipic acid 16, 18
 - crystalline furoic acid dimer with slow mode Morse potential and Fermi resonances 28–30
 - crystalline H(D)-3-thiopheneacrylic acid 20, 24
 - crystalline polarized and unpolarized glutaric acid 17, 20, 21
 - crystalline thiopheneacetic acid 17, 20, 23
- H-bonded compounds, combined
- crystalline oxindole acid dimers 31
- liquid formic acid mixing of monomer and dimer 27, 29
- 1,2-naphthylacetic acid (2-NA) crystals 20, 21, 25
- monomer of $(\text{CH}_3)_2\text{O} \dots \text{HCl}$, electrical anharmonicity 34–36
- Morse potential 23–24, 26
- phosphinic acid dimer 31, 33, 34
- phthalic and terephthalic acid crystals 25, 27–29
- 6-carboxy-2-naphthol 264
- carboxy-substituted naphthols 265
- Car–Parrinello molecular dynamics 41, 56
- CBr_3COOH 217
- CCl_3COOH 217
- CF_3COOH 217
- chain length effect on the degree of self-association of 1-alcohols 237–240
- charge assisted hydrogen bonds (CAHBs) 124, 131–134
- chelate-ring breathing 100
- chelate-ring deformation modes 100
- chemically equivalent oxygen moieties 91
- 2-chlorobenzoic acid (2CBA) 221, 222, 224, 225
- cis*-1,2-dimethylcyclohexane 513
- classical hydrogen bond 129–131
- classical view 442
- Cl/F-H stretchings 39
- 1-Color REMPI (1-C-REMPI) spectrum 330
- condensed-phase acid dissociation constants 89
- continuum of hydrogen bonding 87–93, 97
- cooperative resonance-assisted hydrogen bonds 134–138
- corrphycene 146, 152–154, 169
- Cotton effect 510–512
- Coulomb interaction 263, 268, 285
- Coulomb potential 263
- coupled-cluster theory 93, 94, 107
- coupled reaction-diffusion reactive system 262
- 4-cyano-2,2,6,6-tetramethyl-3,5-heptanedione enol 197
- cyclic hydrogen bond chains
- in alleno-acetylenic cages 512
 - amide-appended corannulenes 511
 - G-quadruplexes 508
 - open and closed form of the AAC 512
 - 4-pyrrolidinopyridines 513
 - tautomerism in campestarenes 510

- cyclic trimer of (HCl)₃ 305
 cyclochiral 513, 514
 cyclohexane 243, 482, 487, 503
 cyclohexanol 243, 244
 cyclohexene-1,2-dicarboxylate
 monoanion 195
 cyclopropanes 482
 cytosine crystal 51–52
- d**
- delocalization of probability density 86
 density functional theory (DFT)
 calculations 40, 68, 75, 146, 173,
 175, 176, 186, 240, 241, 253, 453
 deuterium 74
 isotope effects on chemical shifts
 176–177
 isotope effects on ¹H chemical shifts
 187
 deuterium kinetic isotope effect (DKIE)
 91, 97, 103–105
 2,4-dialkoxy-meta-aminomethylbenzoic
 acid (MAMBA) scaffold 486
 2,6-dibromophenol 249–251
 dicarboxylic acid anion system 195
 2,6-dichlorophenol 249–251
 difluorobenzene 299–300, 331
 difluoromaleate monoanion 195
 2,6-difluorophenol 249–251
 2,6-dihalogenated phenols 249, 251
 2,6-dihalogenophenols 249, 257
 dihydrogen bonds 124
 diisopropylamine-formate system 189
 β-diketone systems 197
 dimensionless oscillator strength 309,
 310, 313
 dimer strong anharmonic coupling theory
 adiabatic approximation for monomer
 3, 5–6
 dimer involving damping Davydov
 coupling and Fermi resonances
 12–13
 Fermi resonances 6–8
 H-bonded centrosymmetric dimer
 8–13
- dimethyl ether (DME) 39, 40, 42, 314
 dimethylsulfoxide (DMSO) 130, 131,
 135–138, 271, 273, 486, 506
 dipalmitoylphosphatidylcholine
 (DOPC)-derived vesicles 496
 dipole moment function (DMF) 36,
 253–256, 309, 385
 2,3-dipyrrol-2-ylquinoxalines 190
 Dirac constant 69
 dispersed fluorescence (DF) spectra 102,
 103
 dispersed fluorescence (DF) spectroscopy
 98
 dissociation energy 296
 of cationic H-bonded complexes
 325–328
 of H-bonded complexes 296–297
 SEP-REMPI technique 328–331
 of the ionic complex 316
 DME-HCl complex 40, 46, 47
 DME-H(D)Cl complexes 45–47
 DME-HF complexes 39, 40, 44–48
 donor-acceptor distance 85, 86, 88–92,
 94, 95, 104–107, 145, 152
 double boundary (DB) solution 267, 284
 dynamical tunneling phenomena 92
 dynamic foldamers 489, 494
 hydrogen-bonded 488–501
 reversible hydrogen-bond directionality
 in 501–508
 dynamic view 442
- e**
- electrolyte aqueous solutions 411–412
 electron density 50–52, 89, 94, 126, 169,
 192, 193, 295, 413, 468, 501
 electronically excited HFF 102, 107
 Energy gap law 301
 energy splitting 67, 69, 97, 104
 EOM-CCSD method 414–416, 418
 equilibrium constant 271, 279, 280, 285,
 307–309, 351, 353
 equilibrium dissociation energy 294,
 333–335
 ethaline (ETH) 132

excess spectroscopy 126–129
 excited state intramolecular hydrogen
 transfer in tropolone 71, 75–79
 excited-state potential surfaces 106
 experimental binding energy 295, 305

f

far-ultraviolet (FUV) spectroscopy
 biomedical science 429
 characteristics and advantages of
 410–411
 development of 428
 of electrolyte aqueous solutions
 411–412
 Group I, II, and XIII cation 412
 interfacial water adsorbed 416–418
 lithium ion (Li⁺) 424–427
 nylons 422–424
 F-H stretching absorption band 46
 fifth, interstitial, non-tetrahedral or
 mismatched water molecules 454
 first-order saddle point 67, 87
 five unit cells 52, 53
 fluorescence-based spectroscopic probes
 98
 fluorescence excitation spectra of HFF
 isotopologs 99
 2-fluorobenzoic acid (2FBA) 220, 221
 foldamers 480
 Aib 490, 495, 499, 500
 β -Hairpin-mimetic trimeric MAMBA
 486
 dynamic (*see* dynamic foldamers)
 helical aromatic oligoamide 484
 hydrogen-bonded (*see* hydrogen-
 bonded foldamers)
 hydrogen-bonded dynamic (*see*
 hydrogen-bonded dynamic)
 reversible hydrogen-bond directionality
 in dynamic (*see* reversible
 hydrogen-bond directionality in
 dynamic)
 fragment molecular-orbital-
 multicomponent molecular-orbital
 method 175

Franck–Condon type progressions 43
 Frank–Condon (FC) distribution 317
 free energy of complexation 307
 freon cryosolvents 188
 Fukui functions 50

g

GAMESS 71
 gas-phase proton affinities 89
 gauche/anti conformations 487
 Gibbs Free energy of H-bonded complex
 formation 307–314
 G protein-coupled receptor (GPCR) 496
 G-quadruplexes 508
 ground-state potential surfaces 95
 ground-state properties of model systems
 93–97
 ground-state TrOH and HFF 94
 guanine crystal 51
 Gutmann-Beckett scale 360

h

β -hairpin-mimetic trimeric MAMBA
 foldamers 486
 halogen-substituted acetic acids
 215–219, 228
 H-bond acceptor (HBA) 273, 304, 314,
 321
 H-bond characteristics
 correlations between geometry 165,
 166
 corrrphycene 152–154
 hemiporphycene 150–152
 inverted/confused porphyrin 162–164
 isoporphycene 154–156
 neo-confused porphyrin 164–165
 parameters 167–168
 porphine 147–148
 porphycene 148–150
 porphyrin-(2.2.0.0) 156, 157
 porphyrin-(3.1.0.0) 157–158
 porphyrin-(4.0.0.0) 158–162
 tautomerization mechanisms 168–169

- H-bond dissociation energy, multiphoton ionization (MPI) techniques 316–324
- H-bond donor (HBD) 270, 273, 274, 304, 308, 311, 314, 320
- H-bonded complex difluorobenzene 299
- H-bonding interactions, microwave spectroscopy 332–335
- Heaviside step function 70
- helical aromatic oligoamide foldamers 484
- hemiporphycene 145, 146, 150–152, 157, 158, 167
- Henri-Rousseau and Blaise model 3
- 1-hexanol 240–241, 243, 244, 257
- homo-conjugated anions 356, 357, 359, 370, 371
- H₂S dimer 294
- hydrogen bond 40, 85
 - charge assisted hydrogen bonds (CAHBs) 131–134
 - classical 129–131
 - continuum of 87–93, 97
 - cooperative resonance-assisted hydrogen bonds 134–138
 - criteria of the existence of 124–125
 - definition 123, 293
 - excess spectroscopy 126–129
 - polarizability 348
 - potentials 173–175, 189, 197, 202
 - spectroscopic metrics for (*see* spectroscopic metrics for hydrogen bonding)
 - strength of 125–126
 - types 176, 385
 - by use of higher overtones 249
 - of water 235
 - weak/moderate hydrogen bonds 138–142
- hydrogen-bond donors (HBD) 270, 273, 274, 455, 479–483, 501–504, 506, 508
- hydrogen-bonded dynamic foldamers
 - Aib helix 494, 495
 - Aib nonapeptides 490
 - attractive or repulsive interactions 488
- 2,6-bis(N-imidazolidin-2-onyl)pyridine oligomers 489
- BQPA 495, 496
- Cu(II)-BQPA binding site 496
- definition 488
- determination of helical excess 490–501
- light-modulated screw-sense induction
 - in phospholipid bilayers 498
- screw-sense preference 490, 493
- screw-sense preference in
 - DOPC-derived vesicles 497
- synthetic molecular communication 489
- Tendrils in Aib foldamers 499
- hydrogen-bonded foldamers
 - accessible conformations of ethylene-bridged triureas 487, 488
 - ACHC- and ACPC-derived foldamers 480, 481
 - amide in 482
 - β-sheet-mimetic oligoureas foldamers 487
 - exploiting repulsive interactions 485
 - helical aromatic oligoamide foldamers 484
 - N,N-diarylformamide balances 480
 - π-stacking interactions 484
 - supramolecular and biological systems 479
 - VT NMR 483
 - X-ray crystallography 482
- hydrogen-bonded systems
 - adenine and thymine crystals 50, 51
 - DME-H(D)Cl and DME-HF complexes 45–48
 - guanine and cytosine crystals 51–52
 - 1-methyl-4-thiouracil 50
 - 1-methyluracil 50
 - molecular dynamics, methodology of 47–48

- hydrogen-bonded systems (*contd.*)
 (CH₃)₂O-HCl and (CH₃)₂O-HF
 complexes 42–45
 spectroscopic signature for ferroelectric
 ice 52–55
 uracil 49–50
- hydrogen-bonded water molecules 418,
 453
- hydrogen bond energy 193
 definition 375–377
 IR characterization of
 carbonyl stretching vibration 387
 low-frequency hydrogen bond
 stretching frequency 385–386
 proton donor deformational
 vibrations 385
 proton donor stretching band
 intensity 384–385
 proton donor stretching band shift
 378, 380–384
 stretching vibrations' force constants
 386–387
 NMR characterization of 377–379
- hydrogen bond geometry
 averaging of NMR parameters and
 proton tautomerism 350–353
 description of 348–350
 IR hydrogen bond correlations
 371–375
 NMR hydrogen bond correlations
 353–371
- hydrogen-bonding motifs 87, 88, 92, 97,
 100
- hydrogen rac-dimethylsuccinate 197
- 10-hydroxybenz[h]quinolines 181
- 9-hydroxy-9-fluorenicarboxylic acid
 (9-HFCA) 315
- 6-hydroxy-2-formylfulvene (HFF) 87, 93
 excited-state spectroscopy of 98–102
 ground-state spectroscopy of 102–105
- 2-hydroxynaphthaldehyde 199
- 8-hydroxypyrene 1,3,6-
 trisdimethylsulfonamide
 (HPTA) 271
- 8-hydroxypyrene 1,3,6-trisulphonate
 (HPTS) photoacid 262
- 8-hydroxyquinoline molecular assemblies
 124
- 3-hydroxyquinoline photoacid 284
- i**
- icebergs 442, 450, 453, 470
- infrared (IR) excitation
 dissociation energy of H-bonded
 complexes 296–297
 Gibbs free energy of H-bonded complex
 formation 307–314
- integrated absorbance 308, 309, 313
- intermolecular complexes 349, 360, 361,
 375, 376
- intermolecular H-bond energy 294, 295
- intermolecular hydrogen bonds (IHB)
 40, 42, 49, 86, 176, 177, 187–189,
 217, 251, 372
- internal complex salt-bridge 214
- intramolecular hydrogen bonds (IMHB)
 93, 177
 halogen-substituted acetic acids
 215–219
ortho chloro-and/or fluoro-substituted
 benzoic acids 219–225
 thiotropolone (TT) 225–228
- intramolecular hydrogen transfer in
 malonaldehyde 68, 71, 73–74, 79
- intramolecular vibrational relaxation
 (IVR) process 228, 328
- intrinsic reaction coordinates (IRC)
 67–77, 79, 227
- intrinsic reaction coordinate-vibrational
 adiabatic (IRC-VA) 72–74
- intrinsic reaction coordinate-Wentzel-
 Kramers-Brillouin (IRC-WKB)
 67–69, 71–75, 77, 79
- inverted/confused porphyrin 146, 162
- ion cluster-CH₃CN complex 139
- ion clusters 139
- ionic liquids (ILs) 124, 189, 194, 383,
 411, 428
- IR hydrogen bond correlations

- carbonyl stretching vibration 375
- proton donor deformational vibrations 374
- proton donor stretching vibration 371–374
- IR-UV double resonance spectroscopy 314–316
- isodesmic reactions 215
- isoporphycene 146, 154–156, 165, 169
- isotope D/H effect 283–285
- isotope effects
 - calculation of 175–176
 - deuterium isotope effects 187
 - deuterium isotope effects on chemical shifts 176–177
 - exchange in the solid state 192–193
 - hydrogen bond energies 193
 - hydrogen bond potentials 173–174
 - hydrogen bond types 176
 - intermolecular hydrogen bonds 187–189
 - intramolecular hydrogen bonds 177–185
 - primary deuterium isotope effects 189–191
 - proteins 185–187
 - protonation states 191
 - solid state NMR 197–202
 - solvent isotope effects and exchange rates 192
 - tautomerism 194–197
- isotope substitution 173, 175, 191, 194, 202, 287

- k**
- kinetic isotope effects (KIE) 86, 91, 97, 108, 283–287
- Kramers–Kronig transformation 418, 419, 440

- l**
- laser-induced fluorescence (LIF) spectroscopy 98–100, 102, 108
- level of theory 96, 97, 105, 133, 244, 309, 334, 355, 384, 386, 413, 425
- linear distortion parameter 44
- lithium ion (Li⁺) 424–427
- local mode perturbation theory (LMPT) model 310
- long-range isotope effects 182, 184–185, 191
- low-barrier hydrogen bonding (LBHBing) 85–109

- m**
- magnetic anisotropy effects 378
- Makri–Miller method 68, 79
- malonaldehyde 68, 71, 73–75, 78, 79
- malondialdehyde 71, 73, 74
- mass analyzed threshold ionization (MATI) technique 323, 324
- mass selected two-color multiphoton ionization technique 321, 322
- 1-methyl-4-thiouracil 40, 41, 49–50
- 1-methyluracil 40, 41, 49–50
- microheterogeneity in alcohol/alcohol and alcohol/alkane binary mixtures 241–244
- microwave spectroscopy 213, 214, 224, 317, 321, 332–335
- molar absorption coefficient 127, 128
- molecular dynamics (MD) simulations 41, 47, 48, 356, 367, 375, 453, 454
- Møller–Plesset perturbation theory 105
- Møller–Plesset second-order perturbation theory (MP2) 72
- multiphoton ionization (MPI) techniques 316–324
- multivariate curve resolution (MCR) analysis 238, 242, 243, 440, 441, 456, 463–465
- mutual diffusion coefficient 263, 268, 285

- n**
- 1-naphthol...2-butyne complex 330, 331
- natural transition orbitals (NTOs) 413
- N-(3,5-dibromosalicylidene)-methylamine 194
- 2N68dS photoacid 266, 268

- neo-confused porphyrin 146, 164–165
- n*-hexane 240–241, 243, 244, 249–251, 257
- NH stretching frequency 147, 153, 160, 166–168
- NIR spectroscopy
- association of 1-hexanol in *n*-hexane 240–241
 - chain length effect on the degree of self-association of 1-alcohols 237–240
 - characteristics 233–235
 - hydrogen bonding by use of higher overtones 249–251
 - hydrogen bonding of water 235–237
 - microheterogeneity in alcohol/alcohol and alcohol/alkane binary mixtures 241–244
 - overtones of $\nu\text{C}\equiv\text{N}$ vibration 244–245
 - pyrrole 252–256
 - weak hydrogen bond in poly(3-hydroxybutyrate) (PHB) 246–249
- nitromalonamide 197
- N-methylacetamide (NMA) 135–138, 418–422
- N-methylformamide (NMF) 135–138, 418–421
- NMR hydrogen bond correlations
- FHF, FHN and FHO bonds 365–369
 - NHN bonds 363–365
 - OHN bonds 360–363
 - OHO bonds- ^{13}C and ^{31}P NMR chemical shifts 356–360
 - OHO bonds- ^1H chemical shifts 353–356
 - vicinal H/D isotope effects 369–371
- N,N*-diarylformamides 479
- N,N'*-disubstituted (thio)ureas 482
- noncovalent interactions 85–87, 92, 93, 95, 96
- nonlinear dielectric effect (NDE) 240, 257
- normal mode (NM) model 295, 310
- Nowick's β -sheets 487
- N-(pyridoxylidene)-methylamine 194, 201, 202
- N-(pyridoxylidene)tolylamine 201, 202
- N-triflylphosphoramidate 492
- O**
- OD(H)-stretching band 438
- $(\text{CH}_3)_2\text{O-HCl}$ complex 42–45
- $(\text{CH}_3)_2\text{O-HF}$ complex 42–45
- OH photoacid 263, 270–273
- OH (OD)-stretching mode 438
- o*-hydroxy Schiff bases 194
- o*-hydroxythioacetophenones 181, 198
- one-bond deuterium isotope effects on ^{15}N chemical shifts 185, 191, 199
- one-water-bridge ESPT reaction 286
- optothermal bolometric determination 297–299
- ortho* chloro-and/or fluoro-substituted benzoic acids 219
- oscillator strength 309–311, 313, 414, 423, 424
- overtones of $\nu\text{C}\equiv\text{N}$ vibration 244–245
- P**
- pentachlorophenol-4-methylpyridine complex 202
- p*-fluorophenol 325–328
- phenol 249–251, 257, 272, 273, 320, 327, 380, 479, 480
- phosphinic acids 31, 33, 189, 358–360, 370, 371
- photoactive yellow protein 175, 370, 371
- photoionization efficiency curve 325
- phthalate monoanion 195
- piroxicam 195
- point-to-point PT reactions 261
- poly(3-hydroxybutyrate) (PHB) 246–249, 257
- porphine 146–148, 164, 168
- porphycene 145, 146, 148–150, 155, 156, 165, 167–169, 175, 363, 364
- porphyrin-(2.2.0.0) 146, 156
- porphyrin-(3.1.0.0) 146, 157–160
- porphyrin-(4.0.0.0) 146, 158, 160–162

- 8-position monosulfonated 2-naphthol
2N8S photoacid 268
- potential-barrier tunneling 86
- potential energy surface (PES) 47, 48,
67, 68, 75, 89, 93, 101, 103, 105,
108, 214, 215, 224, 227–229, 296,
303, 305, 306, 309, 310, 484, 488
- primary deuterium isotope effect 176,
188–191, 202
- proteins 134, 175, 176, 185–187, 193,
357, 362, 370, 429, 437, 479, 481,
483, 497
- proton association 263, 264
- protonation reaction 262, 264, 266, 269,
288
- proton dissociation 262–264, 268, 269,
285
- proton transfer 85–87, 89–94, 96, 97,
100, 103, 104, 106–108, 173, 176,
188, 196, 261, 274–282, 285–289,
346–349, 351–354, 356–358, 360,
363–367, 369, 375–377
- pseudo-diatomic model 333
- pyridoyl benzoyl β -diketones 198
- 4-pyridoyl derivative 198
- pyrrole 252
monomer 253
pyridine complex 253, 254
- 4-pyrrolidinopyridines (PPY) 513, 514
- q**
- quasi-lattice model 236, 256
- r**
- Reisler's measurement 293
- remote protonation 261, 264, 265
- resonance assisted hydrogen bonds
(RAHBs) 124, 134, 135, 137, 138,
176, 177, 190, 384
- resonance enhanced multi-photon
ionization (REMPI) detection
scheme 300–306, 315, 319, 330
- resonant-assisted hydrogen bonding
(RAHBing) 93, 95
- resonant-2-photon ionization (R2PI)
technique 328–330
- reverse spectroscopic problem 345–346
- reversible geminate proton recombination
GEM reactions 262
- reversible hydrogen-bond directionality in
dynamic foldamers
dual hydrogen-bond donor/acceptor
monomers 502
- ethylene-bridged oligoureas foldamers
505, 506
- in ethylene-bridged oligoureas 507
- Eyring-Polanyi equation 505
- hydrogen-bond acceptor 501
- hydrogen bond-donating/accepting
ability 504
- oligoureas 503, 504
- screw-sense preference 503
- reversible proton dissociation-
recombination reaction 262
- rotameric formamides 479
- S**
- SAC-CI calculations 421
- Sapporo-double zeta polarization (DZP)
basis function 72
- Schiff-base macrocycles 508
- secondary intrinsic isotope effects 177
- self-modeling curve resolution (SMCR)
236
- semiclassical tunneling approach
69–71
- SEP-REMPI technique 328–331
- short strong hydrogen bonds (SSHBs)
126, 377
- single-boundary (SB) GEM model 268
- solid state NMR 192, 197–202, 362
- solvatomers 109, 350, 356, 375
- solvent isotope effects and exchange rates
192
- SO₃-substituted 2-naphthol 266
- spectral density 13–14, 33
- spectral markers for proton transfer and
H-bond length 346–348

- spectroscopic metrics for hydrogen bonding
- continuum of hydrogen bonding 87–92
 - excited-state properties of model systems 105–108
 - excited-state spectroscopy of 6-hydroxy-2-formylfulvene 98–102
 - ground-state properties of model systems 93–97
 - ground-state spectroscopy of 6-hydroxy-2-formylfulvene 102–105
 - relationship to tunnelling 92–93
 - spectroscopic signature
 - ferroelectric ice 52–55
 - of low-barrier hydrogen bonding in neutral species 85–109
 - stimulated emission pumping (SEP) 328, 329
 - stretching force constant 332–335
 - strongly hydrogen-bonded (SHB) water species 235–236
 - sulfonamides 482
 - sulfonate substituted naphthols 265
- t**
- tautomerism 194
 - averaging of NMR parameters and proton 350–353
 - in campesterenes 510
 - isotope effects 194–197
 - tautomerization mechanisms 168–169
 - t-butyl alcohol 314
 - tetraethylammonium hydrogen
 - R-(+)-methylsuccinate 197
 - tetraethylammonium (TEA) hydrogen succinate 197
 - 2,3,5,6-tetrafluoroterephthalic acid 175
 - tetramethylammonium (TMA) hydrogen meso-dimethylsuccinate 197
 - 2,2,2-t-fluoroethanol 314
 - thiodibenzoylmethane 198
 - thiophenoxyketenimines 183
 - thiotropolone (TT) 225–228
 - β -thioxoketones 198
 - thymine 41, 50, 51, 184
 - time correlated single photon counting 277
 - time-dependent density functional theory (TDDFT) 75, 413
 - time-independent Schrodinger equation 251
 - time-of-flight (TOF) mass spectrometer 323
 - time resolved fluorescence (TRF) 262
 - decay signal 268
 - measurements 277
 - time-resolved Stokes shift 271
 - TOF-MD-WKB method 69–71, 75, 76, 79
 - trajectory on-the-fly molecular dynamics (TOF-MD) 67–79
 - trans*-2-aminocyclopentanecarboxylic acid (ACPC) 480, 481, 502
 - trans*-1,2-dimethylcyclohexane 513
 - transient absorption spectroscopy 262
 - trihalogeno-substituted acetic acids 217, 221
 - tropolone (TroH) 69, 71, 75–79, 93, 97, 105, 106, 109
 - two-bond deuterium isotope effects on ^{13}C chemical shifts 178–183, 203
 - two-color photodissociation experiments 319
 - two-color REMPI (2C-REMPI)
 - measurements 319
 - two-colour two-photon resonant ionization techniques 321
 - two-dimensional correlation spectroscopy (2DCOS) 238, 239
 - two-dimensional hydrogen bond potentials 173
- u**
- Ubbelohde effect 89, 175, 356, 366
 - umbrella inversion of ammonia 71–73
 - uracil 40, 49, 50, 184

V

- vacuum ultraviolet (VUV) region 409
- valence bond orders 350, 355, 361, 365
- variable-temperature NMR (VT NMR) 483, 491, 505
- velocity map imaging (VMI) 299–307
- vibrational local mode (LM) theory 309
- vibrationally-induced selective conformational isomerization 216
- vibrational pre-dissociation 296
- vibrationless zero-point level 91
- vibronically resolved fluorescence techniques 108
- vibronic transitions 101
- vicinal H/D isotope effect 369

W

- water...amine complexes 311
- water near fully hydrophobic solutes
 - experimental procedure 443–446
 - clathrate formations 448, 449
 - temperature and pressure on the OD stretching band 446–448
- Monte Carlo and MD simulations 453–454
- pure hydrophobic solutes in water solution 449–453
- transmission high pressure cell 443

water spectroscopy

- bulk water 438–441
- IR spectroscopy, water hydrogen bonding, alcohol-water systems 455–470
- water near fully hydrophobic solutes 442–454
- weak hydrogen bond in poly(3-hydroxybutyrate) (PHB) 246–249
- weakly hydrogen-bonded (WHB) water species 235
- weak/moderate hydrogen bonds 138–142
- weak organic Brønsted acids 261
- Wentzel–Kramers–Brillouin (WKB) 67–69, 227

X

- XH spectroscopic method 233

Z

- zero electron kinetic energy (ZEKE) spectroscopy 315, 323, 325, 326
- zero-point averaged rotational constants 333
- zero point energy (ZPE) 5, 7, 72–75, 79, 86, 90, 91, 96, 108, 147, 148, 174, 227, 294–296, 376, 377
- zero-point vibrational energy 72

