

Index

a

adamantane-thiol-protected butterfly-like
 NC [Cu₁₈H₃(S-Adm)₁₂(PPh₃)₄Cl₂]
 (Cu₁₈) 73

Ag₁₂-BPHF@CB[6] 48

Ag₁₂-bpy 46–48, 54–57

Ag₁₄-CH₃CN 26, 27, 49

Ag₁₅Cu₆ cluster 81

Ag₁₄(2e) series 26, 34, 49

Ag₂₃(8e) series 43

Ag₂₉(8e) series 30, 32, 41–42

Ag₃₀(6e) series 40

Ag₄₄(18e) series 32, 39

Ag₇₀(20e) series 30

Ag₇₈(36e) series 27

aggregation-induced emission (AIE) 1,
 56, 69, 70, 80

Ag(I)₂₀, Ag(I)₂₂ series 43

Ag(I)₁₀ series 50

Ag(I)₅ series 43

Ag(I)₆ series 37

Ag(I)₁₂ series 27, 34, 46

Ag(I)₂₀ series 27, 31

Ag₇₈-R/S-BDPP 29

Ag₁₄-R/S-1-PPA 27, 28

Al-Cu intermetalloid cluster
 [(Cp*AlCu)₆H₄] 84

aliphatic dithiolate bridged
 ligands 187, 190–191

alkaline exchange membrane fuel cells
 (AEMFCs) 259

alkynyl 3–5, 45, 65, 68, 77–82, 114, 249

alkynyl ligands modified Cu
 clusters 77–82

all-alkynyl-protected Cu(I) clusters 78

amphiphilic POM assembly
 structures 97

anionic giant POM clusters 94

anionic monolacunary
 [SiW₁₁O₃₉]⁸⁻ clusters 114

anion template method 289

aqueous medium 12, 273–275

aromatic dithiolate bridged ligands 187,
 191–193, 206

asymmetric catalysis, POM-based
 nanostructures for 107–110

atomically precise ultrasmall quantum
 dots (QDs) 157, 205, 217,
 219–221, 227–237

(AuAg)_{34n} 16, 17

Au₂₅(SBu)₁₈⁰ 9, 10, 17–18

b

Ball-and-stick model 168, 286, 288, 290

BDT 24, 41, 197

bidentate ligands 10, 227

c

cadmium chalcogenide supertetrahedral
 clusters (CCSCs)

C_n-type clusters 161–162

inorganic open frameworks 163–169

organic open frameworks

- cadmium chalcogenide supertetrahedral clusters (CCSCs) (*Continued*)
 N-donor ligands 169–175
 other organic ligands 175–176
 photodegradation of organic dyes 178–179
 photoluminescent properties 176–178
Pn-type clusters 160–161
Tn-type clusters 158–160
 carboxylate-based clusters 275
 carboxylate ligands 31, 103, 118, 198, 220–222, 235, 253, 258, 259, 261, 273, 275–280, 283, 288
 carboxyl group-stabilized polyoxozincate cluster 104–105
 catalytic properties 1, 5, 54, 57–58, 65, 67, 68, 108, 109, 116, 183, 192, 198, 262, 287
 cation exchange process 224
 cationic organo-Cu cluster, [Cu₄(PCP)₃]⁺ 74
 [Cd₃₂S₁₄(SPh)₃₆(DMF)₄] cluster 162
 chemically/physically induced methods 242, 245–247
 Chini clusters 241, 242, 244, 245, 248, 255, 256, 258
 chiral amine ligands 27
 chirality 1, 3, 27, 29, 37, 39, 40, 50, 68, 81, 107, 108, 144
 chiral POM-based enantiomers 108, 109
 chiral POM-based materials 107
 chiral pyrrolidine based metal framework 108–109
 chiral thiolate ligands 3
 chloroform-*n*-butanol solvent system 96
 chromophore ligands (CLs) 283
 cluster-assembled materials, feature of 98
 cluster–fullerene complexes 24
 cluster functionalization 7
 cluster-surface strain-promoted alkyne-azide cycloaddition (CS-SPAAC) 7
Cn-type clusters 157, 161–162
 co-immobilization, of Keggin-type POM and Rh catalytic complex 119
 coinage-metal clusters 23
 copper clusters
 luminescence and catalytic properties 65
 copper (Cu) chalcogenide NCs 65
 53-core mixed-valence copper(0)/copper(I) cluster (ZNF/Cu₅₃) 81
 covalently bridged oligomers 10–14
 covalent superatomic crystals 142–145
 croconaine (CR) dye, self-assembly of 120
 C1-type CCSC [Me₄N]₂[S₄Cd₁₇(SPh)₂₈] 161, 162
 C2-type CCSC [Cd₃₂S₁₄(SPh)₃₆(DMF)₄] 162
 Cu alkyne (I) clusters 77
 Cu alkyne(I) clusters 78
 Cu and Cu-based NPs, synthesis of 66, 67
 cubic chalcogenide cluster [Cd₈(SPhF-3)₁₄(DMF)₆(NO₃)](NO₃) (CMM-5) 166
 cuboidal [3Fe4S] clusters 185
 cuboidal [4Fe4S] clusters 185
 Cu₁₄-8CH₃CN cluster 71–72
 Cu₆(C₇H₄ONS)₆ cluster 69
 Cu hydride complexes 74
 Cu(0) nanoclusters 67
 Cu₂₅ NC 74
 Cu(I) alkyne clusters 78
 Cu(I) alkynyl NCs R/S-Cu₁₄ 80
 CuNCs
 assembly of 84–85
 composition and structure 65
 modification and assembly 68–85
 alkynyl ligands modified Cu clusters 77–82
 phosphine ligands modified Cu clusters 74–77
 short organic ligands with carboxyl groups and pyridyl 82

- thiolates ligands modified Cu
 - clusters 68–73
 - synthesis and properties 66–67
- Cu thiolate cluster-assembled material
 - (Cu₁₂-bpy) 84
- d**
 - Dawson-type POM cluster 100
 - density functional theory (DFT) 24, 77, 150, 219, 227, 283
 - deracemization method 30
 - 3,7-diacetyl-1,3,7-triaza-5-phosphabicyclo[3.3.1]-nonane (DAPTA) 188
 - diorganophosphino ligand 252
 - diphenylphosphino-2-pyridine (dppy) 6
 - direct carbonylation method 242–244
 - discrete bridged and fused oligomers of superatoms 137–138
 - discrete silver clusters
 - coordinated S-containing ligands 32–38
 - labile solvent molecules 26–27
 - modification by supramolecular interactions 24–26
 - weakly coordinated non-S auxiliary ligands 27–32
 - DODA (dimethyl dioctadecyl ammonium) 95
 - doping method 218, 223–228
 - drug delivery system (DDS) 14
- e**
 - electrocatalysis, POM-based
 - nanostructures for 110–116
 - electrochemistry, POM-based
 - nanostructures for 110–116
 - electron spray ionization mass spectrometry (ESI MS) 4, 24, 26, 77, 249, 274
- f**
 - [FeFe]-hydrogenase 186, 187, 191, 195, 202, 206
 - Fe-S clusters
 - Cps 195–198
 - cuboidal [3Fe4S] clusters 185
 - cuboidal [4Fe4S] clusters 185
 - [2Fe2S] clusters in photocatalytic H₂ production 202–206
 - first coordination sphere
 - NHC and chelated N-based ligands 189–190
 - phosphine ligands 187–188
 - photosensitive ligands 193–195
 - heterogeneous supports 198–202
 - oxygen 185
 - polynuclear Fe–S complexes 195–196
 - proteinaceous environment 185
 - rhombic [2Fe2S] clusters 185
 - second coordination sphere
 - aliphatic dithiolate bridged ligands 190–191
 - aromatic dithiolate bridged ligands 191–193
 - photosensitive ligands 193–195
 - Fourier-transform infrared spectroscopy (FTIR) 98, 222
 - fullerene-based superatomic materials 148
- g**
 - gadolinium polyoxotungstate 121
 - Glutathione (GSH) 2, 120
 - gold cluster-assembled materials
 - (GCAM) 1, 8–13
 - covalently bridged oligomers and networks 10–14
 - metal–metal bonds 9–10
 - gold cluster assembly (GCA) 11–15
 - biomedical application 14–15
 - magnetism 17–18
 - semiconductivity 16–17
 - gold NCs
 - ligand exchange 2–6
 - post-assembly surface modification 7–8
 - surface locking through coordination 6–7

h

- hexanuclear Cu hydride complexes
 - 74
- high-nuclearity lanthanide clusters
 - anion template method 289
 - anion templates slow release
 - 289–290
 - assembly building blocks 292–293
 - ligand-controlled hydrolysis
 - approach 288–289
 - multiple anion templates, mixed
 - templating anions 290–291
- high-resolution electrospray ionization
 - mass spectrometry
 - (HRESI MS) 24
- homochiral POM-based homochiral
 - metal-organic frameworks 109
- HPLC 3, 242, 275

i

- indium phosphide magic-sized clusters
 - doping method 223–228
 - influence in the synthesis of, InP
 - QDs 230–233
 - InP nanostructures 235–236
 - InP QDs 228–230
 - ligands method 220–223
 - low temperature method 219–220
 - luminescent property of 236–237
 - research 218
 - synthesis mechanism from InP MSCs
 - to InP QDs 233–234
- inorganic open frameworks 157, 163–169
- intra-dye molecular charge transfer
 - (ICT) 283

j

- Janus POM-POSS co-clusters, assembly
 - of 101–102

k

- Keplerate giant clusters 95

l

- lanthanide oxo clusters
 - (LnOCs) 287–293
- Lewis acid and base theory 68
- ligand-controlled hydrolysis
 - approach 288–289, 294
- ligand exchange
 - alkynyl 3–5
 - N-heterocyclic carbenes (NHCs) 5–6
 - thiolate/selenolate 2–4
- ligand-to-metal charge transfer
 - (LMCT) 7, 283
- limit of detection (LOD) 54, 120
- Lindqvist-POSS hybrid clusters 102–107
- L- or D-pyrrolidine-type chiral ionic
 - liquid 108
- lowest energy excitonic transition
 - (LEET) 222, 223, 229
- low temperature method 218–220
- luminescent switching 42, 54–56

m

- magic-sized clusters (MSCs) 217–238
- mass spectrometry (MS) 4, 24, 32, 72,
 - 77, 133, 230, 249, 274
- matrix-assisted laser desorption
 - ionization (MALDI) 32, 230, 249
- matrix-assisted laser desorption
 - ionization mass spectra
 - (MALDI-MS) 249
- $M_6E_8(L)_6$ superatoms
 - assembly
 - covalent superatomic
 - crystals 142–145
 - discrete bridged and fused oligomers
 - of superatoms 137–138
 - supermolecule assembly 138–142
 - electrochemical properties, single-
 - electron currents and electronic
 - transport 145–148
 - gas-phase synthesis 132–133
 - modification

- functionalized 134–135
 - site-differentiated
 - superatoms 135–137
 - properties of 133–134
 - solution-phase synthesis 133
 - synthetic strategies 132
 - thermal transport 148–150
 - Menshutkin S_N2 reaction 8
 - $[(MeCN)_4LCu_4]_6$
 - molecular structure 70
 - metal carbonyl clusters (MCCs) 84, 242, 244, 245
 - metal chalcogenide molecular clusters 131
 - metal–metal bonds 8–10, 16, 56, 85
 - metal nanoclusters (MNCs) 30, 65, 68, 241
 - metal-organic frameworks (MOFs) 12, 14, 39, 43, 44, 54, 95, 102, 105, 106, 109, 110, 113, 114, 193, 198, 206, 271, 272, 294
 - metal oxo clusters (MOCs) 271
 - lanthanide oxo clusters (LnOCs) 287–293
 - high-nuclearity lanthanide clusters 287–291
 - titanium oxo clusters (TiOCs) 278–287
 - structural diversity of 278–282
 - zirconium oxo clusters (ZrOCs)
 - in aqueous medium 273–275
 - in organic medium 275–278
 - mixed-anion template 290
 - mixed templating anions 290–291
 - molecular docking 24, 26
 - molecular photosensitizers (PS) 193
 - molybdenum Mo-based POM cluster, self-assembly of 120
 - monodispersed POM cluster-based assemblies 96
 - multidentate ligands 10, 48, 251
- n**
- nanofibrils 17
 - nano- or sub-nano sized cadmium chalcogenide supertetrahedral clusters (CCSCs) 157–163, 165, 166, 168–170, 173, 175, 176, 178–180
 - N-donor ligands 82, 169–175, 253, 278, 281–282
 - N-heterocyclic carbene (NHC) ligands 5–6, 187, 189
 - Ni₆-substituted POTs 102, 103
 - noncovalent supramolecular interactions 24
 - non-covalently modified POM clusters 122
 - nuclear magnetic resonance (NMR) spectroscopy 24, 71, 222, 275, 278
- o**
- octadecylphosphonate acid (ODPA) 222, 223, 227
 - organic bi/multidentate linkers 45–53
 - organic medium, Zr 275–278
- p**
- pair distribution function (PDF) 274
 - palladium clusters
 - modification of 260–262
 - parabromobenzenethiol (p-BBT) 3
 - Pd carbonyl clusters 242–247
 - $[Pd_n(CO)_x(PR_3)_y]$ clusters 246
 - $[PdS_2C_2(COOMe)_2]_6$ cluster 254
 - phenylacetate ligands 221
 - phosphine ligands modified Cu clusters 74–77
 - phosphonic acids 51, 222, 232, 282, 289
 - photocatalysts, for hydrogen evolution reactions 118

- photocatalytics, POM-based
 nanostructures for 117–119
- photodegradation of organic
 dyes 178–179
- photoluminescence quantum yield
 (PLQY) 30, 32, 34, 37, 46, 74
- physically induced methods
 242, 245
- platinum
 Chini clusters 245, 248, 255
 ligand-tailoring and
 assembly 255–260
- platinum and palladium clusters
 Pt/Pd carbonyl clusters
 (PCCs) 242–247
 Pt/Pd-clusters protected by organic
 ligands 247–255
- Pn-type clusters 157, 160–161
- polyacrylamide gel electrophoresis
 (PAGE) 10
- polyethylene glycol grafted
 polyoxometalate, amphiphilic
 property of 114, 155
- polynuclear Fe–S complexes 187,
 195–196, 206
- polyoxometalates (POMs) 44, 45, 93–95,
 102, 104, 106, 107, 113, 116, 117,
 119–122, 255, 271
 based MOF materials 102–107
 linkers 44–46
 supramolecular nano-building
 blocks 96
- polyoxopalladates (POPs) 254, 255
- POM-Ag cluster coassembly 116
- POM-based homochiral metal-organic
 frameworks 109, 110
- POM-based nanostructures
 for asymmetric catalysis 107–110
 for biological applications 119–122
 for electrochemistry and
 electrocatalysis 110–116
 for photocatalytics 117–119
- POM-based organic framework 111–113
 crystals 111
 for heterogeneous catalytic
 reactions 117
- POM-based supramolecular
 composite 121
- POM-based supramolecular gel 98, 100
- POMOF/SWNT nanocomposites
 (PMNTs) 105
- POMs-based organic–inorganic hybrids,
 classification of 93
- porous CPs 198
- Prato reaction 202
- propeller-like chiral trinuclear Cu(I)
 clusters 74
- Pt carbonyl clusters 243, 244, 248,
 256, 257
- [Pt₁₂(CO)₂₄]²⁻-cluster 248
- [Pt₂Cu₃₄(PET)₂₂Cl₄]²⁻NC 70–71
- Pt/Pd carbonyl clusters (PCCs) 242
 synthesis of 242–247
- Pt/Pd-clusters protected by
 organic ligands
 synthesis 247–255
- [PW₁₂O₄₀]³⁻-based Cu-trz MOFs 105
- q**
 quantum yield (QY) 6, 30, 70, 74, 229,
 233, 237
- r**
 ratiometric luminescent temperature
 sensing 56–57
- redox induced methods 242, 244–245
- rhombic [2Fe2S] clusters 185
- [Ru(bpy)₃]²⁺-derived dicarboxylate
 ligand 118
- s**
 sensing oxygen 54–56
 silver clusters

- atomically precise extended structures 39–53
 - coordination-assembly of 42–53
 - organic bi/multi-dentate linkers 45–53
 - POMs linkers 44–46
 - supramolecular-assembly of 39–42
 - supramolecular chemistry of 58
 - single-cluster nanoring 100
 - single-cluster nanowires 100
 - single-crystal X-ray diffraction (SCXRD) 23, 27, 54, 77, 131, 134, 148, 149, 219, 272, 273
 - single POM cluster nanostructures 99–100
 - site-differentiated superatoms 135–137
 - small-angle X-ray scattering (SAXS) 197, 273–274
 - solvent-mediated precipitating synthesis 77
 - structure-property relationship 58, 59, 134, 272, 278
 - superatomic crystals 131, 139, 142–150
 - superatoms 58, 131–145, 147, 149, 151
 - supermolecular building block (SBB) 271
 - supramolecular interactions, modification 24–26
 - surface-protecting ligands 65
 - surfactant-encapsulated POM clusters 94–101
 - surfactant-encapsulated Wells–Dawson POM 98
 - synergic effect 57
- t**
- T2-CCSC-based three-dimensional (3D) open framework $[\text{Cd}_4(\text{SPh})_8]_n$ 163
 - tetrahedral quadridentate linker tetrakis(4-pyridyloxymethylene) methane (TPOM) 172, 173
 - 5,10,15,20-tetrakis[*meso*-(4-carboxyphenyl)] porphyrinatozinc (TCPP) 261
 - 9,18,27,36-tetrakis[*meso*-(4-carboxyphenyl)]-tetrabenzoporphyrinatozinc (TCPBP) 261
 - tetramercaptotetrathiacalix-[4]arene ligand 69
 - tetranuclear Pt-cluster carboxylates 253
 - thermal-treatment methods 245
 - thermochromism 57, 84
 - thiolate/selenolate 2–4
 - thiolates ligands modified Cu clusters 68–73
 - three-dimensional POM-based metal–organic frameworks 110
 - time-dependent density functional theory (TDDFT) methods 219
 - titanium oxo clusters (TiOCs) 278–287
 - carboxylate ligands-stabilization 279–280
 - ligand modification 283
 - metallic doping 283–287
 - N-donor ligands participation 281–282
 - phosphonate-stabilization 280–281
 - structural diversity of 278–282
 - T n -type clusters 158–160
 - transition metal chalcogenide clusters 131–151
 - transition-metal-substituted polyoxotungstates 102
 - 1,3,5-triaza-7-phosphaadamantane 258, 259
 - trioctylphosphine (TOP) 220, 229–232, 237
 - triphenylphosphine (TPP) 6, 24, 30, 31, 35, 39, 41, 42, 194, 195, 202, 259, 260

V

visible light-induced 1D nanowire
superstructures 116

VOCs 54–56

V-shaped POM-2POSS cluster 102

W

white light-emitting diode
(WLED) 37, 38, 66

X

X-ray absorption fine structure (XAFS) 3

X-ray crystallography 3, 70, 189, 248

Z

zirconium oxo clusters (ZrOCs)

in aqueous medium 273–275

in organic medium 275–278

{Zn- ϵ -Keggin}-based POMOFs 105

Zr oxo clusters (ZrOCs) 272–278

