

1

Synthesis of Diverse Nitrogen Heterocycles Explored in Denitrogenative Transformations

Monalisa Akter and Pazhamalai Anbarasan

Department of Chemistry, Indian Institute of Technology Madras, Chennai, Tamil Nadu, India

1.1 Introduction

Nitrogen heterocycles are extensively present in numerous drug molecules which manifest potent therapeutic activities such as antibacterial, anticancer, antiallergic, potassium-channel activator, antiplatelet, glucosidase, and HIV-1 reverse transcriptase inhibitory activities (Figure 1.1) [1].

Beside living in the core of biologically active molecules, N-heterocycles can act as ligands as well as directing groups in various transition metal catalysis. Importantly, polynitrogen heterocycles can participate in various fruitful transformations that pave the way to access structurally complex molecules of biological importance (Figure 1.2). In this context, *N*-sulfonyl-1,2,3-triazole and its analogs have been exclusively exploited as safe-to-handle diazo surrogates in various denitrogenative transformations. These methods yield a diverse range of structural motifs to facilitate structural modification and total synthesis of natural products and drug molecules [2–4]. Over a period of time, denitrogenative transformations of some related heterocycles such as 5-iodotriazoles [5], F-containing triazoles [6–8], tetrazoles [9, 10], pyridotetrazoles [11], and aminoindazoles [12] have also been explored and established as efficient synthetic tools.

Consequently, easily accessible, atom-economical, and widely compatible synthetic methodologies to access these diverse N-heterocycles are highly desirable. This chapter briefly showcases the development of such primitive to advanced synthetic methodologies to serve the aforementioned purposes.

1.2 Synthesis of 1,2,3-Triazoles

1.2.1 Synthesis of *NH*-Triazoles

NH-triazole is one of the polynitrogen heterocycles that undergoes denitrogenative transformation and their recent development emphasized its importance as a

2 | 1 Synthesis of Diverse Nitrogen Heterocycles Explored in Denitrogenative Transformations




Figure 1.1 Representative drug molecules containing N-heterocycles.






Figure 1.2 Representative drug molecules and natural products synthesized through denitrogenative transformations.

building block and their elegant synthesis. Initially, *NH*-triazoles were prepared *via* the deprotection of various *N*-protected triazoles. In this context, various organic azides with removable protecting groups such as benzyl [13], tropylium [14], trimethylsilyl [15, 16], tosyl [17, 18], (trimethylsilyl)ethoxymethyl (SEM) [19], and *p*-methoxybenzyl [20] azides have been explored along with sodium azide [21, 22]. Later, Sharpless and co-workers [23] introduced three more organic azides, azidomethyl pivalate, azidomethyl morpholine-4-carboxylate, and azidomethyl *N,N*-diethylcarbamate, which deliver a base-labile *N*-protected triazole (Scheme 1.1).

Sharpless [23]

Scheme 1.1 Synthesis of *NH*-triazoles via *N*-protected triazoles. Source: Adapted from Sharpless [23].



In 1989, Banert [24] demonstrated an efficient strategy to synthesize *NH*-triazoles from propargyl azides under mild conditions (Scheme 1.2). Mechanistically, propargyl azide **1.2b** is obtained by treatment of propargyl halide **1.2a** with sodium azide, which undergoes a [3, 3]-sigmatropic rearrangement to generate the reactive allenyl azides **1.2c** that readily cyclizes to triazafulvene intermediate **1.2d**. Finally, the intermediate **1.2d** is trapped by various nucleophiles to afford corresponding triazole **1.2e**.



Scheme 1.2 Synthesis of *NH*-triazoles by using Banert cascade. Source: Adapted from Banert [24].

Despite its high reliability and wide substrate scope, its synthetic utility was hardly explored. In 2005, Sharpless and co-workers [25] exclusively studied this pathway to access diverse *NH*-triazoles and expanded the scope of nucleophiles involved in the process (Scheme 1.3). Recently, Topczewski and co-workers [26] exploited silver(I) fluoride as nucleophile, which facilitated access to α -fluorinated *NH*-1,2,3-triazoles **1.3e** in excellent yields.


In 2016, Dehaen and co-workers [27] disclosed the synthesis of various mono-, di-, and tri-substituted triazoles **1.4c** from the reaction of enolizable ketones **1.4a**, NH₄OAc, and nitrophenyl azide **1.4b** under mild acidic condition (Scheme 1.4). Simultaneously, a β -cyclodextrin-mediated multicomponent synthesis of *NH*-triazoles **1.4e** from propynals **1.4d**, trimethylsilyl azide, and malononitrile in water was reported by Medvedeva and co-workers [28]. Besides, the use of amine in place of malononitrile under microwave irradiation furnished the imine-substituted

Scheme 1.3 Synthesis of diverse *NH*-triazoles.Scheme 1.4 Synthesis of *NH*-triazoles through three-component reactions.


triazole with a shorter reaction time [29]. Subsequently, Guan and co-workers [30–33] accomplished the synthesis of various 4-aryl-*NH*-1,2,3-triazoles **1.4g** through three-component reaction of aldehydes **1.4f**, nitromethane, and NaN₃. Later, Negrón-Silva and co-workers [34] developed a heterogeneous catalytic system consisting of Al-MCM-41 and sulfated zirconia to accomplish the same synthesis.

In 2019, Shu and Wu reported a molecular iodine-mediated cascade [4 + 1] cyclization of *N*-tosylhydrazones **1.5a** and sodium azide in presence of MsOH to access 4-aryl-*NH*-1,2,3-triazoles **1.5b** (Scheme 1.5) [35]. Subsequently, the group of Gao and Shu achieved the synthesis of 4-aryl-*NH*-1,2,3-triazoles **1.5d** via an iodine-mediated condensation-cyclization of α -azido ketones **1.5c** with *p*-toluenesulfonyl hydrazide


Shu and Wu [35]




Gao and Shu [36]



Wu [37]



Matsugi [38]




Scheme 1.5 Synthesis of 4-aryl-NH-1,2,3-triazoles.


[36]. Recently, Wu and co-workers demonstrated the synthesis of *NH*-triazole under azide-free conditions *via* an iodine-mediated [2 + 2 + 1] cyclization of methyl ketones **1.5e**, *p*-toluenesulfonyl hydrazide, and 1-aminopyridinium iodide **1.5f** [37]. Solvent-free synthesis of 4-aryl-*NH*-1,2,3-triazoles **1.5i** has been demonstrated by Matsugi and co-workers [38] from benzyl ketones **1.5h** exploiting diphenyl phosphorazidate in the presence of 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU).

Catalyst-free synthesis of 4-acyl-*NH*-1,2,3-triazoles **1.6b** was reported by Wen and Wan, which involves water-mediated cycloaddition reactions of enaminones **1.6a** and tosyl azide (Scheme 1.6) [39, 40]. Instead of enaminones, Gribanov et al. [41] employed alkyl nitriles **1.6c** and azide **1.6d** in the presence of KO^tBu

Wen and Wan [39]



Gribanov [41]


Scheme 1.6 Synthesis of various *NH*-triazoles.

for the synthesis of 5-amino-1,2,3-triazoles **1.6e**, which on subsequent Dimroth rearrangement affords **1.6f** at elevated temperature under solvent-free conditions in one pot.

1.2.2 Synthesis of *N*-Sulfonyl-1,2,3-triazoles

For years, a large number of *N*-sulfonyl-1,2,3-triazoles have been extensively exploited as diazo surrogate in numerous denitrogenative transformations. In general, sulfonylation of *NH*-1,2,3-triazoles **1.7a** with sulfonyl chlorides could furnish the corresponding *N*-sulfonyl-1,2,3-triazoles **1.7b** (Scheme 1.7). But the major drawback of this strategy is the formation of a mixture of regioisomeric products **1.7b** and **1.7c**, which significantly reduces its efficiency and applicability [42].


Beryozkina and Fan [42]



Scheme 1.7 Synthesis of *N*-sulfonyl-1,2,3-triazoles from *NH*-triazoles. Source: Adapted from Beryozkina and Fan [42].

On the other hand, 1,2,3-triazoles **1.8d** were readily achieved through the copper-catalyzed azide-alkyne cycloaddition (CuAAC) as reported by Sharpless and co-workers in 2002 (Scheme 1.8) [43–45]. This reaction appeared to be the most effective click reaction over the traditional Huisgen cycloaddition due to its remarkably high regioselectivity and yields. Various 1,4-disubstituted triazoles **1.8d** could be synthesized from terminal alkynes **1.8a** and azides **1.8b** (Scheme 1.8). However, the use of sulfonyl azides led to the formation of various secondary products **1.8g** instead of the desired triazoles **1.8d** via the generation of ketenimine


Sharpless [43]



Scheme 1.8 Synthesis of *N*-sulfonyl-1,2,3-triazoles through CuAAC. Source: Adapted from Sharpless [43].

intermediate **1.8f** [46, 47]. The formation of ketenimine was due to the poor stability of the copper-triazole species **1.8c**.

To increase the stability of sulfonyl substituted **1.8c** and for the synthesis of sulfonyltriazoles, in 2007, for the first time, Chang, Fokin and co-workers utilized stoichiometric amount of base, 2,6-dimethylpyridine in the CuI-catalyzed selective synthesis of 4-substituted *N*-sulfonyl-1,2,3-triazoles from the corresponding terminal alkynes and sulfonyl azides (Scheme 1.9) [48, 49]. Subsequently, Fu and co-workers [50] reported an inexpensive catalytic system by exploiting the thioanisole as ligand in combination with CuBr in water medium at room temperature. It was emphasized that the addition of sulfur-containing ligands could inhibit cleavage of N1—N2 bond and stabilize the 5-cuprated triazole species. Two years later, Pérez and co-workers [51] demonstrated the synthesis of *N*-sulfonyl-1,2,3-triazoles using well-defined copper complex, [Tpm^{*},^{Br}Cu(NCCH₃)]BF₄. On the other hand, Raushel and Fokin [52] employed Cu(I)-thiophene-2-carboxylate (CuTC) complex, in the absence of external ligand, under mild conditions for more efficient and general synthesis of *N*-sulfonyltriazoles (Scheme 1.9).



Reaction conditions:

Chang and Fokin [48]: CuI, 2,6-dimethylpyridine, CHCl₃, 0 °C

Fu [50]: CuBr, PhSMe, H₂O, rt, 16 h

Pérez [51]: [Tpm^{*},^{Br}Cu(NCCH₃)]BF₄, CHCl₃, 40 °C

Raushel and Fokin [52]: CuTC, toluene or H₂O, rt, 15 h


Hu [53]: Cu(OAc)₂·H₂O, 2-aminophenol, CH₃CN, rt, 20–50 min

Cazin [54]: Cu(I)-NHC complex, toluene, rt, 15 h

Scheme 1.9 Cu-catalyzed synthesis of *N*-sulfonyl-1,2,3-triazoles.

In 2011, Hu and co-workers [53] reported 2-aminophenol as a suitable ligand in copper-catalyzed highly selective synthesis of *N*-sulfonyltriazoles. Herein, it was proposed that 2-aminophenol plays a dual role as a reductant and ligand. Importantly, the first triazole possessing two electron-withdrawing groups was synthesized successfully by using this strategy. In 2018, Cazin and co-workers [54] employed Cu(I)-NHC complexes for the selective synthesis of 4-substituted 1,2,3-triazoles through click reaction.

But the limitation of all these above-mentioned methods remains in the selective synthesis of 1,4-disubstituted triazoles. Croatt and co-workers [55] documented the synthesis of 5-substituted *N*-sulfonyl-1,2,3-triazoles **1.10c** from corresponding terminal alkynes **1.10a** and sulfonyl azides **1.10b** using a strong base, such as




Scheme 1.10 Synthesis of multisubstituted *N*-sulfonyl-1,2,3-triazoles.

n-BuLi (Scheme 1.10). Interestingly, the synthesis of various 1,4,5-trisubstituted triazoles was also achieved by trapping the 4-lithio-1,5-disubstituted triazole in the reaction mixture with suitable electrophiles. In 2008, Ramachary and co-workers reported the first proline-catalyzed synthesis of 4,5-disubstituted triazoles **1.10f** from Hagemann's ester **1.10d** and tosyl azides **1.10e** [56]. Later, Anbarasan and co-workers [57] accomplished a general proline-mediated synthesis of a diverse range of 4,5-disubstituted 1,2,3-triazoles **1.10i** with excellent regioselectivity from substituted 1,3-dicarbonyl compounds **1.10g** and sulfonyl azides **1.10h** through enamine-azide cycloaddition.


In 2016, Zhang and co-worker [58] achieved the synthesis of 4-substituted 1,2,3-triazoles **1.11c** from (*Z*)-arylviny bromides **1.11a** and sulfonyl azides through the generation of arylacetylene (Scheme 1.11). This strategy involves KOH-promoted HBr elimination from the vinylbromide **1.11a** followed by a copper-catalyzed [3 + 2] cycloaddition of the resulted alkyne **1.11b** with azide. Subsequently, Ma and co-workers [59] reported a regioselective synthesis of 5-sulfamido-*N*-sulfonyl-1,2,3-triazoles **1.11f** in high yields through a Cu-catalyzed cycloaddition of terminal alkynes **1.11d** and sulfonyl azides **1.11e** using 1.2 equivalents of LiO^tBu. Most recently, Kim and co-workers [60] reported the first continuous flow synthesis of *N*-sulfonyl-1,2,3-triazoles.

1.2.3 Synthesis of *N*-Trifluoromethyl-1,2,3-triazoles

N-trifluoromethyl-1,2,3-triazoles also have been explored in the denitrogenative transformations owing to the unique physicochemical properties of F-containing organic moieties. But, unlike the synthesis of other *N*-sulfonyl-1,2,3-triazoles, the




synthesis of *N*-fluoroalkyl-1,2,3-triazoles was not achieved following conventional alkyne-azide cycloaddition strategy. In 1977, Holton and co-workers [61] studied various 1,3-dipolar cycloaddition across the C=N bond of polyfluoroazaolefins with diazomethane (Scheme 1.12). A mixture of products was obtained in the reaction of perfluoro-*N*-(trifluoromethyl)butanimidoyl fluoride **1.12a** with excess diazomethane, in which, the 1-(trifluoromethyl)-5-(heptafluoropropyl)-1,2,3-triazole **1.12c** was present as a major product. Recently, Xu, Guan, and Wang demonstrated the synthesis of *N*-trifluoromethyl-1,2,3-triazoles **1.12f** via a similar [3 + 2] cyclization of DMAP-stabilized *N*-CF₃ nitrilium salts **1.12d** with diazo compounds **1.12e** [62].



In 2017, Beier and co-workers [63] developed an efficient strategy to synthesize perfluoroalkyl azides (Scheme 1.13). Subsequently, the reactivity of perfluoroalkyl azide was explored in azide-alkyne cycloaddition reaction to access *N*-perfluoroalkyl triazoles.

Additionally, a one-pot synthesis of such triazoles has also been demonstrated directly starting from trifluoromethylsilane in good yield and regioselectivity.




Method A: [CuSO₄·5H₂O] (10 mol%), sodium L-ascorbate (10 mol%), H₂O

Method B: [CuMeSal] (1–5 mol%), THF

Scheme 1.13 Synthesis of *N*-perfluoroalkyl-1,2,3-triazoles. Source: Adapted from Beier [63].


1.2.4 Synthesis of 5-Iodo-1,2,3-triazoles

Similar to electron withdrawing group-substituted triazoles, 5-iodo-1,2,3-triazoles having a tethered nucleophile are shown to undergo denitrogenative transformations. These triazoles were also synthesized through azide-alkyne cycloaddition. In this context, in 2005, Wu et al. [64] developed a strategy to access 5-iodo-1,2,3-triazoles **1.14c** in one-pot *via* Cu(I)-catalyzed reaction of organo-azides **1.14a** with terminal alkynes **1.14b** in presence of iodine as an electrophilic trapping reagent (Scheme 1.14). It was proposed that the reaction involves the formation of an intermediate of Cu(I) salt of triazole. Interestingly, yield was increased drastically when ICl was used instead of I₂. Subsequently, Zhang and Zhang [65] reported a CuI-NBS-mediated multicomponent reaction of azides **1.14d** and alkynes **1.14e**, which also facilitated access to 5-iodo-1,4-disubstituted-1,2,3-triazoles **1.14f** [66, 67]. To improve the reactivity and chemoselectivity, ICl was *in situ* generated from the combination of CuI and NCS and used as an effective iodonium source. Recently, these strategies were further modified to accomplish the synthesis in aqueous medium [68, 69].



Scheme 1.14 Synthesis of 5-iodo-1,2,3-triazoles.

In 2009, Hein, Fokin, and co-workers [70] reported the synthesis of 5-iodotriazoles from the CuI-catalyzed cycloaddition of 1-iodoalkynes **1.15b** and organic azides **1.15a** (Scheme 1.15). Various ligands such as TBTA or TTTA were screened to furnish the desired products in good yields with low catalyst loadings and shorter reaction times. Zhu and co-workers [71] reported an assisting ligand-free strategy to access various 5-iodo-triazoles **1.15f** (Scheme 1.15). Subsequently, the effect of ligand [72] as well as mechanism [73] involved in the reaction were




Scheme 1.15 Cu-catalyzed synthesis of 5-iodo-1,2,3-triazoles.

established. Besides, several other groups have also accomplished the synthesis of 5-iodotriazoles [74–80].

1.2.5 Synthesis of Pyridotriazoles

Pyridotriazoles being analogous to triazoles are highly compatible in denitrogenative transformations, but only selected strategies are known for their synthesis. Traditionally, pyridotriazoles **1.16b** have been synthesized from the corresponding hydrazones **1.16a** of pyridine-2-carbaldehydes or 2-pyridylketones *via* oxidative cyclization (Scheme 1.16) [81, 82]. Although some of the hydrazones are just refluxed in methanol under air to afford the corresponding pyridotriazoles, suitable oxidizing agents such as nickel peroxide, air and a cupric salt, potassium ferrocyanide and bicarbonate, manganese dioxide or (diacetoxyiodo)benzene are




Scheme 1.16 Synthesis of pyridotriazoles. Source: Deady and Devine [83]/with permission of Elsevier.

necessary for most of the hydrazones. In the case of sensitive substrates, pyridotriazoles **1.16d** could be obtained from the reaction of tosyl hydrazones **1.16c** of pyridine-2-carbaldehydes or 2-pyridylketones with base, usually morpholine (Scheme 1.16). In 2006, Deady and Devine demonstrated an alternative strategy to access pyridotriazoles **1.16f** from an aminonaphthyridinone **1.16e** [83]. Compound **1.16e** could form a diazonium salt upon treatment with aqueous sodium nitrate and fluoroboric acid. Then triazolopyrido aldehyde **1.16f** is obtained as a white solid upon treatment of the salt with sodium hydroxide in water.

1.2.6 Synthesis of Triazoloindoles

Due to the growing interest in the denitrogenative transformation of triazoles, in 2014, Lu and Wang accomplished the synthesis of diverse triazoloindoles **1.17b** from alkynes **1.17a** and sulfonylazides *via* a copper-catalyzed tandem process (Scheme 1.17), and demonstrated their reactivity [84]. This strategy involves a sequence of amine group-stabilized CuAAC and C–N coupling reactions. Further study on the triazoloindoles revealed its predominant existence as open-chain form, 3-diazoindolin-2-imines **1.17c**. Subsequently, the same group developed an efficient catalyst-free synthesis of 3-diazoindolin-2-imines **1.17f** from indole and sulfonyl azide [85]. Indoles bearing alkyl, allyl, and benzyl substituents at N1-position and electron-donating substituents in the aryl ring underwent the reaction smoothly and effectively. However, a significant loss in yields was witnessed in the presence of electron-withdrawing groups in aryl ring.

Lu and Wang [84]




Lu and Wang [85]



Scheme 1.17 Synthesis of triazoloindoles. Source: Adapted from Lu and Wang [84, 85].

1.2.7 Synthesis of Benzotriazoles

Denitrogenative transformations of benzotriazoles have also witnessed significant development in past years. Particularly, benzotriazoles containing aryl, acyl, alkenyl, and alkyl substituents at N1-position have been explored well. In general, benzotriazole derivatives are obtained through the selective protection of readily available benzotriazoles (BtH) as shown in Scheme 1.18 [86].




Scheme 1.18 Synthesis of benzotriazole derivatives.


1.3 Synthesis of 1,2,3-Thiadiazoles

1,2,3-Thiadiazoles were also demonstrated to undergo denitrogenative transformation through the generation of thiavinyl carbenes. Their synthesis involves an interesting cyclization with the introduction of sulfur. In 1955, Hard and Mori reported an effective synthesis of 1,2,3-thiadiazoles from acylhydrazones and thionyl chloride (Scheme 1.19) [87]. This method is widely applicable for the synthesis of a variety of substituted thiadiazoles. Alternatively, thiadiazoles could be accessed *via* the reaction of diazo compound with Lawesson's reagent [88].

Hard and Mori [87]



By using Lawesson's reagent




Scheme 1.19 Synthesis of 1,2,3-thiadiazoles. Source: Hurd and Mori [87]/American Chemical Society.

1.4 Synthesis of Tetrazoles

1.4.1 Synthesis of 1*H*-Tetrazoles




1*H*-tetrazole as a privileged scaffold has received significant attention from researchers in development of drugs as well as in the denitrogenative transformations. Initially, 1*H*-tetrazoles have been exclusively synthesized *via* the

[3 + 2]-cycloaddition reaction of nitriles with various azides such as alkylsilyl azides, alkyltin azides, or sodium azide [89–93]. Major drawbacks of these primitive strategies were use of expensive and toxic metals, water sensitivity, or the possible generation of volatile and explosive hydrazoic acid [94–101]. Consequently, an efficient catalytic synthetic route was highly desired. In 2008, Yamamoto and co-workers [102] developed an efficient Cu(I)-catalyzed synthesis of 5-substituted 1*H*-tetrazoles **1.20b** in good yields via the [3 + 2] cycloaddition of nitriles **1.20a** and trimethylsilyl azide (Scheme 1.20). Recently, Gholizadeh and co-workers [103] achieved the synthesis of tetrazole from nitrile using ionic liquid as a catalyst.

Yamamoto [102]

Scheme 1.20 Synthesis of 1*H*-tetrazoles. Source: Yamamoto [102]/with permission of Elsevier.


In 2015, Abdollahi-Alibeik et al. [104] reported a Cu-MCM-41 nanoparticles-catalyzed novel three-component reaction of aldehydes **1.21a**, hydroxylamine, and sodium azide for the synthesis of various tetrazoles **1.21b** (Scheme 1.21). Subsequently, polymer-supported palladium nanoparticles were exploited for the synthesis of tetrazole by Abadi and co-workers [105]. Roy, Islam, and Sen [106] developed a Ni(OH)₂ nanoparticles-catalyzed synthesis of 5-substituted 1*H*-tetrazoles **1.21f** from aldoximes **1.21e** and sodium azide in water under mild reaction conditions. Following this development, a remarkable advancement in the synthesis of 1*H*-tetrazole has recently been observed [9, 10, 107–112].

Abdollahi-Alibeik [104]**Abadi [105]****Roy, Islam and Sen [106]**

Scheme 1.21 Metal nanoparticles-catalyzed synthesis of 1*H*-tetrazoles.


1.4.2 Synthesis of Pyridotetrazoles

Pyridotetrazoles analogous to tetrazoles also undergo various denitrogenative transformations facilitating access to structurally complex molecules. In 1983, Iyengar and co-workers [113] observed the unprecedented formation of tetrazolo[1,5-*a*]pyridine [114] upon reaction of pyridine-*N*-oxide with arenesulfonyl azide (Scheme 1.22). Later, Keith [115, 116] optimized the reaction conditions and explored the substrate scope extensively. Further, the reaction was modified by Tzschucke and co-workers [117] to achieve the synthesis from pyridine-*N*-oxides, *p*-toluenesulfonyl chloride, and sodium azide.



Scheme 1.22 Synthesis of pyridotetrazoles.


In 1997, Noravyan and co-workers [118] synthesized tetrazolo[1,5-*a*]pyrano[3,4-*c*]pyridines **1.23b** from chloropyridines **1.23a** and sodium azide (Scheme 1.23). Later, Cuny and co-workers [119] employed trimethylsilyl azide and tetrabutylammonium fluoride hydrate for the synthesis of various pyridotetrazoles **1.23d** from 2-halopyridines **1.23c**.




Scheme 1.23 Synthesis of pyridotetrazoles from 2-halopyridines.

Recently, Chattopadhyay and co-workers [120, 121] synthesized various pyridotetrazoles **1.24c** in two steps starting from corresponding 2-halopyridines **1.24a** (Scheme 1.24), and explored their reactivity in denitrogenative transformations. Subsequently, Ghasemzadeh and co-workers [122] reported a three-component reaction of 1*H*-tetrazole-5-amine **1.24d**, benzaldehydes **1.24e**, and 3-cyanoacetyl indole **1.24f** in presence of hexamethylenetetramine-based ionic liquid/MIL-101(Cr) metal-organic framework as a recyclable catalyst, which facilitated access to diverse tetrazolo[1,5-*a*]pyrimidine-6-carbonitriles in excellent yields.

Chattopadhyay [120]



Ghasemzadeh [122]




Scheme 1.24 Synthesis of diverse pyridotetrazoles.


1.4.3 Synthesis of Tetrazolo[1,5-*a*]quinolines


Tetrazolo[1,5-*a*]quinolines are largely found in natural products exhibiting biological activities. In 1997, Smalley and co-workers synthesized tetrazolo[1,5-*a*]quinolines **1.25c** by the reaction of 2-azidocarboxaldehyde **1.25a** with acetonitrile in the presence of piperidine or sodium ethoxide as base in ethanol (Scheme 1.25) [123]. The product formation could be rationalized through a Knoevenagel condensation of the acetonitrile with aldehyde **1.25a** followed by an intramolecular 1,3-dipolar cycloaddition reaction of azide with pendant cyano moiety present in **1.25b** intermediate. Later, Bekhit and co-workers reported the synthesis of tetrazolo[1,5-*a*]quinoline-4-carboxaldehyde **1.25e** from corresponding 2-chloroquinoline-3-carboxaldehyde **1.25d** by treating with sodium azide in DMSO/AcOH. [124–126] Recently, Chakroborty et al. modified the reaction conditions to attain a green and more efficient synthesis [127].

Smalley [123]



Bekhit [124]

Scheme 1.25 Synthesis of tetrazolo[1,5-*a*]quinolines.



Scheme 1.26 Synthesis of various tetrazolo[1,5-*a*]quinolines.


In 2019, Kuhakarn and co-workers [128] accomplished the synthesis of tetrazolo[1,5-*a*]quinolines **1.26b** from the reaction of *o*-alkynylisocyanobenzenes **1.26a** with sodium azide under metal- and base-free conditions (Scheme 1.26). This strategy involves nucleophilic addition of azide to isocyanide followed by 6-endo cyclization. On the other hand, Wu and co-workers [129] synthesized various tetrazolo[1,5-*a*]quinolines **1.26e** through *tert*-butylnitrite-mediated radical cyclization of 1*H*-tetrazol-5-amines **1.26c** with alkynes **1.26d**.

1.5 Synthesis of 3-Aminoindazoles



Interestingly, 3-aminoindazoles also has emerged as an efficient synthetic tool compatible with denitrogenative transformations. In 2002, Semple and co-workers [130–132] synthesized various 3-aminoindazoles from halobenzonitriles to access important non-covalent thrombin inhibitors. Later, the groups of Ma [133], Devkate [134], and Zhang and Hu [135] accomplished the synthesis of diverse 3-aminoindazoles **1.27c** from reaction of halobenzonitriles **1.27a** with hydrazines **1.27b** under the modified reaction conditions (Scheme 1.27). In 2010, Fabis and co-workers [136] reported a two-step synthesis of substituted 3-aminoindazoles **1.27g** from 2-bromobenzonitriles **1.27d** and hydrazones **1.27e**. This strategy involves *N*-arylation of hydrazone **1.27e** with *o*-bromobenzonitriles **1.27d** followed by deprotection of the hydrazone intermediate **1.27f** in methanol.

In 2011, Callot and co-workers [137] demonstrated a Buchwald–Hartwig C–N coupling reaction of various 3-haloindazoles **1.28a** to access 3-aminoindazoles **1.28b** in good to moderate yields (Scheme 1.28). Subsequently, Charrette and co-workers [138, 139] prepared 3-aminoindazoles from tertiary amides *via* the generation of aminohydrazones **1.28c** followed by intramolecular C–H amination.

Recently, Olmos and co-workers [140] reported the synthesis of 3-aminoindazoles **1.28g** from 3-(2-bromoaryl)-1,2,4-oxadiazol-5(4*H*)-ones **1.28e** and amines **1.28f** which involves an intramolecular N–N bond formation.




Scheme 1.27 Synthesis of 3-aminoindazoles from 2-halobenzonitriles.



Scheme 1.28 Synthesis of various substituted 3-aminoindazoles.

1.6 Synthesis of Benzotriazinones


Another unique N-heterocycle, benzotriazinone, has been explored in several denitrogenative transformations. In 2015, Xu and co-workers [141–146] prepared several benzotriazinones **1.29b** from anthranilamides **1.29a** *via* diazotization, and cyclization reactions in one-pot (Scheme 1.29). They further derivatized these benzotriazinones by attaching spirocyclic indoline-2-one moieties, and studied



Scheme 1.29 Synthesis of benzotriazinones.

their nematocidal activities. Alternatively, the groups of Foroumadi [147], and Zhang and Li [148] prepared the benzotriazinones **1.29f** from isatoic anhydride **1.29c** and alkyl amines **1.29d** following the reaction sequence shown below. At first, isatoic anhydride **1.29c** and alkyl amines **1.29d** were stirred in water at room temperature to furnish 2-amino-*N*-(prop-2-yn-1-yl)benzamide **1.29e** as a white solid. Next, exposure of **1.29e** to acidic solution of sodium nitrite resulted in the formation of benzotriazinones **1.29f** via intramolecular nitrogen–nitrogen bond formation.


In 2017, Rocha-Alonzo and co-workers [149] developed a novel strategy to access 1,2,3-benzotriazinones **1.30b** from the reaction of 2-(*o*-aminophenyl)oxazolines **1.30a** with isoamyl nitrite in methanol (Scheme 1.30). Recently, Rayes and co-workers [150] reported the synthesis from anthranilhydrazide as well as methyl anthranilate.



Scheme 1.30 Synthesis of various *N*-substituted benzotriazinones.

In 2010, Murakami and co-workers reported a Ni-catalyzed annulation of 1,2,3,4-benzothiazine-1,1(2*H*)-dioxides **1.31d** with allenes (Scheme 1.31) [151]. Herein, benzothiazinone **1.31d** was synthesized from *ortho*-nitrobenzenesulfonyl chloride **1.31a**. Coupling of **1.31a** with methylamine followed by reduction of

Murakami [151]



Scheme 1.31 Synthesis of benzothiazinones. Source: Murakami [151]/John Wiley & Sons.

1.31b using Zn gives *ortho*-amino-*N*-methylbenzenesulfonamide **1.31c**. Finally, HONO-mediated ring closing of **1.31c** provides **1.31d** in 82% yield.

1.7 Summary and Outlook

Owing to the remarkable footprints left by various nitrogen heterocycles in the advancement of drugs, and total synthesis of natural products of biological importance, the development of methodologies to access such N-heterocycles has caught significant attention of synthetic chemists in past years. Wherein, atom and cost economy, green approach, and sustainability of the strategy were utmost priorities among other parameters. This chapter highlights the important modifications encountered over the years to accomplish the synthesis of diverse N-heterocycles, for example, triazoles, tetrazoles, 3-aminobenzotriazoles, benzotriazinones, etc. with better yields and selectivity, specifically, which undergo various denitrogenative transformations. Delightfully, many synthetic routes have been developed in a metal-free manner and using milder conditions, some of which are even sustained in a water medium. In recent times, various metal nanoparticles have also been exploited in the advancement of synthesis of these N-heterocycles. Nonetheless, the synthesis of triazoloindoles, pyridotriazoles, thiadiazoles, and benzotriazinones has not received much attention. Hence, further development of synthetic methodologies to attain better accessibility is desirable due to the growing demand for these heterocycles as synthetic tools.

References

- 1 Anbarasan, P., Yadagiri, D., and Rajasekar, S. (2014). Recent advances in transition-metal-catalyzed denitrogenative transformations of 1,2,3-triazoles and related compounds. *Synthesis* 46: 3004–3023.

- 2 Akter, M., Rupa, K., and Anbarasan, P. (2022). 1,2,3-Triazole and its analogues: new surrogates for diazo compounds. *Chem. Rev.* 122: 13108–13205.
- 3 Kadhim, M.M., Mahmood, E.A., Heravi, M.R.P. et al. (2022). The synthesis of biologically active 1-sulfonyl-1,2,3-triazoles from sulfonyl azides and alkynes: a focus review. *J. Sulfur Chem.* 44: 377–391.
- 4 Bahadorikhalili, S., Divar, M., Damghani, T. et al. (2021). *N*-Sulfonyl ketenimine as a versatile intermediate for the synthesis of heteroatom containing compounds. *J. Org. Chem.* 939: 121773–121812.
- 5 Danilkina, N.A., Govdi, A.I., and Balova, I.A. (2020). 5-Iodo-1*H*-1,2,3-triazoles as versatile building blocks. *Synthesis* 52: 1874–1896.
- 6 Kumar, A.S., Kommu, N., Ghule, V.D., and Sahoo, A.K. (2014). Synthesis of trifluoromethyl-substituted *N*-aryl-poly-1,2,3-triazole derivatives. *J. Mater. Chem. A* 2: 7917–7926.
- 7 Niedermann, K., Früh, N., Senn, R. et al. (2012). Direct electrophilic *N*-trifluoromethylation of azoles by a hypervalent iodine reagent. *Angew. Chem. Int. Ed.* 51: 6511–6515.
- 8 Usachev, B.I. (2018). Chemistry of fluoroalkyl-substituted 1,2,3-triazoles. *J. Fluor. Chem.* 210: 6–45.
- 9 Neochoritis, C.G., Zhao, T., and Dömling, A. (2019). Tetrazoles via multicomponent reactions. *Chem. Rev.* 119: 1970–2042.
- 10 Verma, A., Kaur, B., Venugopal, S. et al. (2022). Tetrazole: a privileged scaffold for the discovery of anticancer agents. *Chem. Biol. Drug Des.* 100: 419–442.
- 11 Das, S.K., Roy, S., and Chattopadhyay, B. (2023). Transition-metal-catalyzed denitrogenative annulation to access high-valued N-heterocycles. *Angew. Chem. Int. Ed.* 62: e202210912.
- 12 Guo, Y. and Gao, Q. (2022). Recent advances in 3-aminoindazoles as versatile synthons for the synthesis of nitrogen heterocycles. *Org. Biomol. Chem.* 20: 7138–7150.
- 13 Wiley, R.H., Hussung, K.F., and Moffat, J. (1956). The preparation of 1,2,3-triazole. *J. Org. Chem.* 21: 190–192.
- 14 Looker, J.J. (1965). Preparation of 1,2,3-triazoles from 7-azido-1,3,5-cycloheptatriene. A displacement from nitrogen. *J. Org. Chem.* 30: 638–639.
- 15 Birkofer, L., Ritter, A., and Richter, P. (1963). Thermolyse silylierter tetrazole. *Chem. Ber.* 96: 2750–2757.
- 16 Birkofer, L. and Wegner, P. (1966). 1.3-Cycloadditionen mit trimethylsilyl-azid; über isomere *N*-acetyl-1.2.3-triazole. *Chem. Ber.* 99: 2512–2517.
- 17 Huisgen, R., Knorr, R., Möbius, L., and Szeimies, G. (1965). Einige Beobachtungen zur Addition organischer Azide an CC-Dreifachbindungen. *Chem. Ber.* 98: 4014–4021.
- 18 Harvey, G.R. (1966). A new synthesis of 1,2,3-triazoles and diazo esters from phosphorus ylids and azides. *J. Org. Chem.* 31: 1587–1590.
- 19 Holzer, W. and Ruso, K. (1992). Functionalisation of 1,2,3-triazole via lithiation of 1-[2-(trimethylsilyl)ethoxy]methyl-1*H*-1,2,3-triazole. *J. Heterocyclic Chem.* 29: 1203–1207.

- 20 Buckle, B.D.R. and Rockell, C.J.M. (1982). The 4-methoxybenzyl group, a versatile N-protecting group for the synthesis of N-unsubstituted *v*-triazoles. *J. Chem. Soc. Perkin Trans. 1*: 627–630.
- 21 Tanaka, Y., Velen, S.R., and Miller, S.I. (1973). Syntheses and properties of *H*-1,2,3-triazoles. *Tetrahedron* 29: 3271–3283.
- 22 Jankovič, D., Virant, M., and Gazvoda, M. (2022). Copper-catalyzed azide-alkyne cycloaddition of hydrazoic acid formed in situ from sodium azide affords 4-monosubstituted-1,2,3-triazoles. *J. Org. Chem.* 87: 4018–4028.
- 23 Loren, J.C., Krasinski, A., Fokin, V.V., and Sharpless, K.B. (2005). *NH*-1,2,3-triazoles from azidomethyl pivalate and carbamates: base-labile N-protecting groups. *Synlett* 18: 2847–2850.
- 24 Banert, K. (1989). Synthesis of 1,2,3-triazoles from propargyl azides by rearrangement of the azido group. Indication of short-lived allenyl azides and triazafulvenes. *Chem. Ber.* 122: 911–918.
- 25 Loren, J.C. and Sharpless, K.B. (2005). The Banert cascade: a synthetic sequence to polyfunctional *NH*-1,2,3-triazoles. *Synthesis* 9: 1514–1520.
- 26 Alexander, J.R., Kevorkian, P.V., and Topczewski, J.J. (2021). Intercepting the Banert cascade with nucleophilic fluorine: direct access to α -fluorinated *NH*-1,2,3-triazoles. *Chem. Commun.* 57: 5024–5027.
- 27 Thomas, J., Jana, S., Liekens, S., and Dehaen, W. (2016). A single-step acid catalyzed reaction for rapid assembly of *NH*-1,2,3-triazoles. *Chem. Commun.* 52: 9236–9239.
- 28 Medvedeva, A.S., Demina, M.M., Vu, T.D. et al. (2016). β -Cyclodextrin-catalyzed three-component synthesis of 4,5-disubstituted 1,2,3-(*NH*)-triazoles from propynals, trimethylsilyl azide and malononitrile in water. *Mendeleev Commun.* 26: 326–328.
- 29 Medvedeva, A.S., Demina, M.M., Kon'kova, T.V. et al. (2017). Microwave assisted solvent- and catalyst-free three-component synthesis of *NH*-1,2,3-triazoloimines. *Tetrahedron* 73: 3979–3985.
- 30 Hui, R., Zhao, M., Chen, M. et al. (2017). One-pot synthesis of 4-aryl-*NH*-1,2,3-triazoles through three-component reaction of aldehydes, nitroalkanes and NaN_3 . *Chin. J. Chem.* 35: 1808–1812.
- 31 Quan, X.-J., Ren, Z.-H., Wang, Y.-Y., and Guan, Z.-H. (2014). *p*-Toluenesulfonic acid mediated 1,3-dipolar cycloaddition of nitroolefins with NaN_3 for synthesis of 4-Aryl-*NH*-1,2,3-triazoles. *Org. Lett.* 16: 5728–5731.
- 32 Efimov, I.V. (2019). Recent methods for the synthesis of *NH*-1,2,3-triazoles. *Chem. Heterocycl. Compd.* 55: 28–30.
- 33 Dutta, B., Hazarika, P.K., Saika, P. et al. (2023). [DDQM][TFSI]: a room temperature ionic liquid as an active electrode material for supercapacitor devices and a catalyst for rapid synthesis of 4-aryl-*NH*-1,2,3-triazoles under microwave irradiation. *New J. Chem.* <https://doi.org/10.1039/D3NJ00862B>.
- 34 Vergara-Arenas, B.I., Lomas-Romero, L., Ángeles-Beltrán, D. et al. (2017). Multicomponent synthesis of 4-aryl-*NH*-1,2,3-triazoles in the presence of Al-MCM-41 and sulfated zirconia. *Tetrahedron Lett.* 58: 2690–2694.

- 35 Shu, W.-M., Zhang, X.-F., Zhang, X.-X. et al. (2019). Metal-free cascade [4 + 1] cyclization access to 4-aryl-NH-1,2,3-triazoles from *N*-tosylhydrazones and sodium azide. *J. Org. Chem.* 84: 14919–14925.
- 36 Ren, M.-T., Li, M., Wang, A.-J. et al. (2020). Iodine-mediated condensation-cyclization of α -azido ketones with *p*-toluenesulfonyl hydrazide for synthesis of 4-aryl-NH-1,2,3-triazoles. *Eur. J. Org. Chem.* 2020: 2233–2236.
- 37 Huang, C., Geng, X., Zhao, P. et al. (2021). Direct synthesis of 4-aryl-1,2,3-triazoles via I₂-promoted cyclization under metal- and azide-free conditions. *J. Org. Chem.* 86: 13664–13672.
- 38 Ishihara, K., Ishihara, K., Kato, Y. et al. (2022). A solvent-free synthesis of 4-aryl-NH-1,2,3-triazoles from ketones utilizing diphenyl phosphorazidate. *Synth. Commun.* 52: 1326–1335.
- 39 Yang, L., Wu, Y., Yang, Y. et al. (2018). Catalyst-free synthesis of 4-acyl-NH-1,2,3-triazoles by water-mediated cycloaddition reactions of enamines and tosyl azide. *Beilstein J. Org. Chem.* 14: 2348–2353.
- 40 Opsomer, T. and Dehaen, W. (2021). Metal-free syntheses of *N*-functionalized and NH-1,2,3-triazoles: an update on recent developments. *Chem. Commun.* 57: 1568–1590.
- 41 Gribanov, P.S., Atoian, E.M., Philippova, A.N. et al. (2021). One-pot synthesis of 5-amino-1,2,3-triazole derivatives via dipolar azide–nitrile cycloaddition and Dimroth rearrangement under solvent-free conditions. *Eur. J. Org. Chem.* 2021: 1378–1384.
- 42 Beryozkina, T.V., Efimov, I.V., Fabian, W.M.F. et al. (2015). Reactivity of 1,2,3-triazoles towards sulfonyl chlorides. A novel approach to 1- and 2-sulfonyl-4-azolyl-1,2,3-triazoles. *Tetrahedron* 71: 6189–6195.
- 43 Rostovtsev, V.V., Green, L.G., Fokin, V.V., and Sharpless, K.B. (2002). A step-wise Huisgen cycloaddition process: copper(I)-catalyzed regioselective “ligation” of azides and terminal alkynes. *Angew. Chem. Int. Ed.* 41: 2596–2599.
- 44 Tornøe, C.W., Christensen, C., and Meldal, M. (2002). Peptidotriazoles on solid phase: [1,2,3]-triazoles by regiospecific copper(I)-catalyzed 1,3-dipolar cycloadditions of terminal alkynes to azides. *J. Org. Chem.* 67: 3057–3064.
- 45 De Nino, A., Maiuolo, L., Costanzo, P. et al. (2021). Recent progress in catalytic synthesis of 1,2,3-triazoles. *Catalysts* 11: 1120–1167.
- 46 Meldal, M. and Tornøe, C.W. (2008). Cu-Catalyzed azide-alkyne cycloaddition. *Chem. Rev.* 108: 2952–3015.
- 47 Hein, J.E. and Fokin, V.V. (2010). Copper-catalyzed azide-alkyne cycloaddition (CuAAC) and beyond: new reactivity of copper(I) acetylides. *Chem. Soc. Rev.* 39: 1302–1315.
- 48 Yoo, E.J., Ahlquist, M., Kim, S.H. et al. (2007). Copper-catalyzed synthesis of *N*-sulfonyl-1,2,3-triazoles: controlling selectivity. *Angew. Chem. Int. Ed.* 46: 1730–1733.
- 49 Yoo, E.J., Ahlquist, M., Bae, I. et al. (2008). Mechanistic studies on the Cu-catalyzed three-component reactions of sulfonyl azides, 1-alkynes and amines, alcohols, or water: dichotomy via a common pathway. *J. Org. Chem.* 73: 5520–5528.

- 50 Wang, F., Fu, H., Jiang, Y., and Zhao, Y. (2008). Copper-catalyzed cycloaddition of sulfonyl azides with alkynes to synthesize *N*-sulfonyltriazoles “on water” at room temperature. *Adv. Synth. Catal.* 350: 1830–1834.
- 51 Cano, I., Nicasio, M.C., and Pérez, P.J. (2010). Copper(I) complexes as catalysts for the synthesis of *N*-sulfonyl-1,2,3-triazoles from *N*-sulfonylazides and alkynes. *Org. Biomol. Chem.* 8: 536–538.
- 52 Raushel, J. and Fokin, V.V. (2010). Efficient synthesis of 1-sulfonyl-1,2,3-triazoles. *Org. Lett.* 12: 4952–4955.
- 53 Liu, Y., Wang, X., Xu, J. et al. (2011). Highly controlling selectivity of copper(I)-catalyzed azide/alkyne cycloaddition (CuAAC) between sulfonyl azides and normal alkynes or propynoates. *Tetrahedron* 67: 6294–6299.
- 54 Lazreg, F. and Cazin, C.S.J. (2018). Copper(I)-NHC complexes as efficient catalysts for the synthesis of 1,4-disubstituted 1,2,3-sulfonyltriazoles in air. *Organometallics* 37: 679–683.
- 55 Meza-Avina, M.E., Patel, M.K., Lee, C.B. et al. (2011). Selective formation of 1,5-substituted sulfonyl triazoles using acetylides and sulfonyl azides. *Org. Lett.* 13: 2984–2987.
- 56 Ramachary, D.B., Ramakumar, K., and Narayana, V.V. (2008). Amino acid-catalyzed cascade [3 + 2]-cycloaddition/hydrolysis reactions based on the push-pull dienamine platform: Synthesis of highly functionalized *NH*-1,2,3-triazoles. *Chem. Eur. J.* 14: 9143–9147.
- 57 Rajasekar, S. and Anbarasan, P. (2019). A general proline-catalyzed synthesis of 4,5-disubstituted *N*-sulfonyl-1,2,3-triazoles from 1,3-dicarbonyl compounds and sulfonyl azide. *Chem. Asian J.* 14: 4563–4567.
- 58 Zhang, W. and Xu, W. (2016). Synthesis of 1-arylsulfonyl-1,2,3-triazoles from (*Z*)-arylvinyl bromides by sequential elimination–cycloaddition reaction. *Chem. Heterocycl. Compd.* 52: 192–195.
- 59 Zheng, X., Wan, Y., Ling, F., and Ma, C. (2017). Copper-catalyzed tandem reaction of terminal alkynes and sulfonyl azides for the assembly of substituted aminotriazoles. *Org. Lett.* 19: 3859–3862.
- 60 Kwon, Y.-J., Lee, S.-G., and Kim, W.-S. (2023). Continuous flow synthesis of *N*-sulfonyl-1,2,3-triazoles for tandem relay Cu/Rh dual catalysis. *J. Org. Chem.* 88: 1200–1214.
- 61 Coe, P.L. and Holton, A.G. (1977). Some reactions of diazomethane with polyfluoroazaolefins. *J. Fluor. Chem.* 10: 553–564.
- 62 Zhang, R.Z., Zhang, R.X., Wang, S. et al. (2022). An *N*-trifluoromethylation/cyclization strategy for accessing diverse *N*-trifluoromethyl azoles from nitriles and 1,3-dipoles. *Angew. Chem. Int. Ed.* 61: e202110749.
- 63 Blastik, Z.E., Voltrová, S., Matoušek, V. et al. (2017). Azidoperfluoroalkanes: synthesis and application in copper(I)-catalyzed azide–alkyne cycloaddition. *Angew. Chem. Int. Ed.* 56: 346–349.
- 64 Wu, Y.-M., Deng, J., Li, Y., and Chen, Q.-Y. (2005). Regiospecific synthesis of 1,4,5-trisubstituted-1,2,3-triazole via one-pot reaction promoted by copper(I) salt. *Synthesis* 8: 1314–1318.

- 65 Li, L., Zhang, G., Zhu, A., and Zhang, L. (2008). A convenient preparation of 5-iodo-1,4-disubstituted-1,2,3-triazole: multicomponent one-pot reaction of azide and alkyne mediated by CuI-NBS. *J. Org. Chem.* 73: 3630–3633.
- 66 Li, L., Li, Y., Li, R. et al. (2011). A new synthetic protocol for one-pot preparations of 5-halo-1,4-disubstituted-1,2,3-triazoles. *Aust. J. Chem.* 64: 1383–1389.
- 67 Bogdan, A.R. and James, K. (2011). Synthesis of 5-iodo-1,2,3-triazole-containing macrocycles using copper flow reactor technology. *Org. Lett.* 13: 4060–4063.
- 68 Li, L., Ding, S., Yang, Y. et al. (2017). Multicomponent aqueous synthesis of iodo-1,2,3-triazoles: single-step models for dual modification of free peptide and radioactive iodo labeling. *Chem. Eur. J.* 23: 1166–1172.
- 69 Li, L., Xing, X., Zhang, C. et al. (2018). Novel synthesis of 5-iodo-1,2,3-triazoles using an aqueous iodination system under air. *Tetrahedron Lett.* 59: 3563–3566.
- 70 Hein, J.E., Tripp, J.C., Krasnova, L.B. et al. (2009). Copper(I)-catalyzed cycloaddition of organic azides and 1-iodoalkynes. *Angew. Chem. Int. Ed.* 48: 8018–8021.
- 71 Brotherton, W.S., Clark, R.J., and Zhu, L. (2012). Synthesis of 5-iodo-1,4-disubstituted-1,2,3-triazoles mediated by in situ generated copper(I) catalyst and electrophilic triiodide ion. *J. Org. Chem.* 77: 6443–6455.
- 72 Barsoum, D.N., Brassard, C.J., Deeb, J.H.A. et al. (2013). Synthesis of 5-iodo-1,2,3-triazoles from organic azides and terminal alkynes: ligand acceleration effect, substrate scope, and mechanistic insights. *Synthesis* 45: 2372–2386.
- 73 Barsoum, D.N., Okashah, N., Zhang, X., and Zhu, L. (2015). Mechanism of copper(I)-catalyzed 5-iodo-1,2,3-triazole formation from azide and terminal alkyne. *J. Org. Chem.* 80: 9542–9551.
- 74 Wang, D., Chen, S., and Chen, B. (2014). “Green” synthesis of 1,4-disubstituted 5-iodo-1,2,3-triazoles under neat conditions, and an efficient approach of construction of 1,4,5-trisubstituted 1,2,3-triazoles in one pot. *Tetrahedron Lett.* 55: 7026–7028.
- 75 Lal, S., Rzepa, H.S., and Díez-González, S. (2014). Catalytic and computational studies of N-heterocyclic carbene or phosphine-containing copper(I) complexes for the synthesis of 5-iodo-1,2,3-triazoles. *ACS Catal.* 4: 2274–2287.
- 76 Zhang, J., Chen, W., Wang, B. et al. (2015). One-pot three-component synthesis of 1,4,5-trisubstituted 5-iodo-1,2,3-triazoles from 1-copper(I) alkyne, azide and molecular iodine. *RSC Adv.* 5: 14561–14566.
- 77 Griбанov, P.S., Topchiy, M.A., Karsakova, I.V. et al. (2017). General method for the synthesis of 1,4-disubstituted 5-halo-1,2,3-triazoles. *Eur. J. Org. Chem.* 2017: 5225–5230.
- 78 Govdi, A.I., Danilkina, N.A., Ponomarev, A.V., and Balova, I.A. (2019). 1-Iodobuta-1,3-diyne in copper-catalyzed azide-alkyne cycloaddition: a one-step route to 4-ethynyl-5-iodo-1,2,3-triazoles. *J. Org. Chem.* 84: 1925–1940.
- 79 Gharpure, S.J., Naveen, S., Chavan, R.S., and Padmaja (2020). Regioselective synthesis of halotriazoles and their utility in metal catalyzed coupling reactions. *Eur. J. Org. Chem.* 2020: 6870–6886.
- 80 Arenas, J.L. and Crousse, B. (2021). An overview of 4- and 5-halo-1,2,3-triazoles from cycloaddition reactions. *Eur. J. Org. Chem.* 2021: 2665–2679.

- 81 Jones, G. (2002). The chemistry of the triazolopyridines: an update. *Adv. Heterocycl. Chem.* 83: 1–70.
- 82 Jones, G. and Abarca, B. (2010). The chemistry of the [1,2,3]triazolo[1,5-*a*]pyridines: an update. *Adv. Heterocycl. Chem.* 100: 195–252.
- 83 Deady, L.W. and Devine, S.M. (2006). Novel annulated products from aminonaphthyridinones. *Tetrahedron* 62: 2313–2320.
- 84 Xing, Y., Sheng, G., Wang, J. et al. (2014). Preparation of triazoloindoles via tandem copper catalysis and their utility as α -imino rhodium carbene precursors. *Org. Lett.* 16: 1244–1247.
- 85 Sheng, G., Huang, K., Chi, Z. et al. (2014). Preparation of 3-diazoindolin-2-imines via cascade reaction between indoles and sulfonylazides and their extensions to 2,3-diaminoindoles and imidazo[4,5-*b*]indoles. *Org. Lett.* 16: 5096–5099.
- 86 Katritzky, A.R., Lan, X., Yang, J.Z., and Denisko, O.V. (1998). Properties and synthetic utility of N-substituted benzotriazoles. *Chem. Rev.* 98: 409–548.
- 87 Hurd, C.D. and Mori, R.I. (1955). On acylhydrazones and 1,2,3-thiadiazoles. *J. Am. Chem. Soc.* 77: 5359–5364.
- 88 Kurandina, D. and Gevorgyan, V. (2016). Rhodium thiavinyl carbenes from 1,2,3-thiadiazoles enable modular synthesis of multisubstituted thiophenes. *Org. Lett.* 18: 1804–1807.
- 89 Duncia, J.V., Pierce, M.E., and Santella, J.B. III, (1991). Three synthetic routes to a sterically hindered tetrazole. A new one-step mild conversion of an amide into a tetrazole. *J. Org. Chem.* 56: 2395–2400.
- 90 Huff, B.E. and Staszak, M.A. (1993). A new method for the preparation of tetrazoles from nitriles using trimethylsilylazide/trimethylaluminum. *Tetrahedron Lett.* 34: 8011–8014.
- 91 Wittenberger, S.J. and Donner, B.G. (1993). Dialkyltin oxide mediated addition of trimethylsilyl azide to nitriles. A novel preparation of 5-substituted tetrazoles. *J. Org. Chem.* 58: 4139–4141.
- 92 Koguro, K., Oga, T., Mitsui, S., and Orita, R. (1998). Novel synthesis of 5-substituted tetrazoles from nitriles. *Synthesis* 1998: 910–914.
- 93 Curran, D.P., Hadida, S., and Kim, S.-Y. (1999). Tris(2-perfluorohexylethyl)tin azide: A new reagent for preparation of 5-substituted tetrazoles from nitriles with purification by fluorous/organic liquid-liquid extraction. *Tetrahedron* 55: 8997–9006.
- 94 Demko, Z.P. and Sharpless, K.B. (2001). Preparation of 5-substituted 1*H*-tetrazoles from nitriles in water. *J. Org. Chem.* 66: 7945–7950.
- 95 Himo, F., Demko, Z.P., Noodleman, L., and Sharpless, K.B. (2002). Mechanisms of tetrazole formation by addition of azide to nitriles. *J. Am. Chem. Soc.* 124: 12210–12216.
- 96 Demko, Z.P. and Sharpless, K.B. (2002). An expedient route to the tetrazole analogues of α -amino acids. *Org. Lett.* 4: 2525–2527.
- 97 Himo, F., Demko, Z.P., Noodleman, L., and Sharpless, K.B. (2003). Why is tetrazole formation by addition of azide to organic nitriles catalyzed by zinc(II) salts? *J. Am. Chem. Soc.* 125: 9983–9987.

- 98 Amantini, D., Beleggia, R., Fringuelli, F. et al. (2004). TBAF-Catalyzed synthesis of 5-substituted 1*H*-tetrazoles under solventless conditions. *J. Org. Chem.* 69: 2896–2898.
- 99 Ek, F., Manner, S., Wistrand, L.-G., and Frejd, T. (2004). Synthesis of fused tetrazole derivatives via a tandem cycloaddition and N-allylation reaction and parallel synthesis of fused tetrazole amines. *J. Org. Chem.* 69: 1346–1352.
- 100 Kantam, M.L., Kumar, K.B.S., and Sridhar, C. (2005). Nanocrystalline ZnO as an efficient heterogeneous catalyst for the synthesis of 5-substituted 1*H*-tetrazoles. *Adv. Synth. Catal.* 347: 1212–1214.
- 101 Kantam, M.L., Balasubrahmanyam, V., and Kumar, K.B.S. (2006). Zinc hydroxyapatite-catalyzed efficient synthesis of 5-substituted 1*H*-tetrazoles. *Synth. Commun.* 36: 1809–1814.
- 102 Jin, T., Kitahara, F., Kamijo, S., and Yamamoto, Y. (2008). Copper-catalyzed synthesis of 5-substituted 1*H*-tetrazoles via the [3 + 2] cycloaddition of nitriles and trimethylsilyl azide. *Tetrahedron Lett.* 49: 2824–2827.
- 103 Aali, E., Gholizadeh, M., and Noroozi-Shad, N. (2022). 1-Disulfo-[2,2-bipyridine]-1,1-dium chloride ionic liquid as an efficient catalyst for the green synthesis of 5-substituted 1*H*-tetrazoles. *J. Mol. Struct.* 1247: 131289–131295.
- 104 Abdollahi-Alibeik, M. and Moaddeli, A. (2015). Multi-component one-pot reaction of aldehyde, hydroxylamine and sodium azide catalyzed by Cu-MCM-41 nanoparticles: a novel method for the synthesis of 5-substituted 1*H*-tetrazole derivatives. *New J. Chem.* 39: 2116–2122.
- 105 Abadi, S.S.A.D.M. and Zarchi, M.A.K. (2017). A novel route for the synthesis of 5-substituted 1-*H* tetrazoles in the presence of polymer-supported palladium nanoparticles. *New J. Chem.* 41: 10397–10406.
- 106 Halder, M., Islam, M.M., Singh, P. et al. (2018). Sustainable generation of Ni(OH)₂ nanoparticles for the green synthesis of 5-substituted 1*H*-tetrazoles: a competent turn on fluorescence sensing of H₂O₂. *ACS Omega* 3: 8169–8180.
- 107 Swami, S., Sahu, S.N., and Shrivastava, R. (2021). Nanomaterial catalyzed green synthesis of tetrazoles and its derivatives: a review on recent advancements. *RSC Adv.* 11: 39058–39086.
- 108 Ghorbani-Choghamarani, A., Shiri, L., and Azadi, G. (2016). The first report on the eco-friendly synthesis of 5-substituted 1*H*-tetrazoles in PEG catalyzed by Cu(II) immobilized on Fe³O₄@SiO₂@l-arginine as a novel, recyclable and non-corrosive catalyst. *RSC Adv.* 6: 32653–32660.
- 109 Movaheditabar, P., Javaherian, M., and Nobakht, V. (2022). Igneous rock powder as a heterogeneous multi-oxide nano-catalyst for the synthesis of 5-substituted-1*H*-tetrazoles in polyethylene glycol. *J. Iran. Chem. Soc.* 19: 1805–1816.
- 110 Mashhoori, M.-S. and Sandaroos, R. (2022). New ecofriendly heterogeneous nano-catalyst for the synthesis of 1-substituted and 5-substituted 1*H*-tetrazole derivatives. *Sci. Rep.* 12: 15364–15379.

- 111 Sajjadi, M., Nasrollahzadeh, M., Ghafuri, H. et al. (2022). Modified chitosan-zeolite supported Pd nanoparticles: a reusable catalyst for the synthesis of 5-substituted-1*H*-tetrazoles from aryl halides. *Int. J. Biol. Macromol.* 209: 1573–1585.
- 112 Bharathi, M., Indira, S., Vinoth, G. et al. (2022). A green one-pot three-component syntheses of tetrazole under microwave irradiation using Ni Schiff base complex of mixed ligands on MCM-41 as an efficient heterogeneous catalyst. *J. Porous. Mater.* 30: 691–703.
- 113 Reddy, K.S., Iyengar, D.S., and Bhalerao, U.T. (1983). The reaction of pyridine N-oxide and its benzoanalogues with arenesulfonyl azides: novel synthesis of tetrazoloazines. *Chem. Lett.* 1983: 1745–1748.
- 114 Boyer, J.H. and Schoen, W. (1956). 2,3- Ψ -Dinitrosopyridines. *J. Am. Chem. Soc.* 78: 423–425.
- 115 Keith, J.M. (2006). One-step conversion of pyridine N-oxides to tetrazolo [1,5-*a*]pyridines. *J. Org. Chem.* 71: 9540–9543.
- 116 Tamba, S., Nitani, M., Seo, T. et al. (2022). Tetrazolo[1,5-*a*]pyridine-containing π -conjugated systems: synthesis, properties, and semiconducting characteristics. *Org. Lett.* 24: 3792–3796.
- 117 Liu, S., Lentz, D., and Tzschucke, C.C. (2014). Conversion of pyridine N-oxides to tetrazolopyridines. *J. Org. Chem.* 79: 3249–3254.
- 118 Paronikyan, E.G., Sirakanyan, S.N., and Noravyan, A.S. (1997). Synthesis of derivatives of tetrazolo[1,5-*a*]- and oxazolo[4,5-*b*]pyrano(thiopyrano) [3,4-*c*]pyridines. *Chem. Heterocycl. Compd.* 33: 1101–1103.
- 119 Laha, J.K. and Cuny, G.D. (2008). Synthesis of tetrazolo[1,5-*a*]pyridines utilizing trimethylsilyl azide and tetrabutylammonium fluoride hydrate. *Synthesis* 2008: 4002–4006.
- 120 Roy, S., Khatua, H., Das, S.K., and Chattopadhyay, B. (2019). Iron(II)-based metalloradical activation: switch from traditional click chemistry to denitrogenative annulation. *Angew. Chem. Int. Ed.* 58: 11439–11443.
- 121 Khatua, H., Das, S.K., Roy, S., and Chattopadhyay, B. (2021). Dual reactivity of 1,2,3,4-tetrazole: manganese-catalyzed click reaction and denitrogenative annulation. *Angew. Chem. Int. Ed.* 60: 304–312.
- 122 Abdollahi-Basir, M.H., Mirhosseini-Eshkevari, B., Zamani, F., and Ghasemzadeh, M.A. (2021). Synthesis of tetrazolo[1,5-*a*]pyrimidine-6-carbonitriles using HMTA-BAIL@MIL-101(Cr) as a superior heterogeneous catalyst. *Sci. Rep.* 11: 5109–5120.
- 123 Porter, T.C., Smalley, R.K., Teguche, M., and Purwono, B. (1997). Tetrazolo[1,5-*a*]quinolines and 1,2,3-triazolo[1,5-*a*]quinazolines by the action of cyanocarbanions on 2-azidoarylcarbonyl compounds. *Synthesis* 1997: 773–777.
- 124 Bekhit, A.A., El-Sayed, O.A., Aboulmagd, E., and Park, J.Y. (2004). Tetrazolo[1,5-*a*]quinoline as a potential promising new scaffold for the synthesis of novel anti-inflammatory and antibacterial agents. *Eur. J. Med. Chem.* 39: 249–255.

- 125 Meth-Cohn, O., Narine, B., and Tarnowski, B. (1981). A versatile new synthesis of quinolines and related fused pyridines. Part 5. The synthesis of 2-chloroquinoline-3-carbaldehydes. *J. C. S. Perkin I. Perkin I*: 1520–1530.
- 126 Kategaonkar, A.H., Labade, V.B., Shinde, P.V. et al. (2010). Synthesis and antimicrobial activity of tetrazolo[1,5-*a*]-quinoline-4-carbonitrile derivatives. *Monatsh. Chem.* 141: 787–791.
- 127 Chakroborty, S., Ramírez-Lopez, S.C., and Unnamatla, M.V.B. (2022). Synthesis of alkoxy-alkyl- tetrazolo[1,5-*a*] quinoline & tetrazolo[1,5-*a*] quinoline-4-carbaldehyde derivatives under green conditions. *Mater. Today Proc.* 57: 2423–2427.
- 128 Khaikate, O., Soorukram, D., Leowanawat, P. et al. (2019). Azide-triggered bicyclization of *o*-alkynylisocyanobenzenes: synthesis of tetrazolo[1,5-*a*]quinolines. *Eur. J. Org. Chem.* 2019: 7050–7057.
- 129 Liu, T., Ji, Y.-G., and Wu, L. (2019). *tert*-Butyl nitrite-mediated radical cyclization of tetrazole amines and alkynes toward tetrazolo[1,5-*a*]quinolines. *Org. Biomol. Chem.* 17: 2619–2623.
- 130 Cui, J.J., Araldi, G.L., Reiner, J.E. et al. (2002). Non-covalent thrombin inhibitors featuring P3-heterocycles with P1-bicyclic arginine surrogates. *Bioorg. Med. Chem. Lett.* 12: 2925–2930.
- 131 Halley, F. and Sava, X. (1997). Synthesis of 5-cyanoindazole and 1-methyl and 1-aryl-5-cyanoindazoles. *Synth. Commun.* 27: 1199–1207.
- 132 Orsini, P., Menichincheri, M., Vanotti, E., and Panzeri, A. (2009). Highly efficient synthesis of 5-benzyl-3-aminoindazoles. *Tetrahedron Lett.* 50: 3098–3100.
- 133 Xu, L., Peng, Y., Pan, Q. et al. (2013). Assembly of substituted 3-aminoindazoles from 2-bromobenzonitrile via a CuBr-catalyzed coupling/condensation cascade process. *J. Org. Chem.* 78: 3400–3404.
- 134 Devkate, C.G., Patil, A.M., Warad, K.D. et al. (2017). Ceric(IV) ammonium nitrate catalyzed highly efficient synthesis of 3-aminoindazole and their antibacterial screening. *Int. J. Res. Pharm. Chem.* 7: 513–517.
- 135 Zhang, C., Zhao, H., Li, Z. et al. (2020). Rapid access to 3-aminoindazoles from nitriles with hydrazines: a strategy to overcome the basicity barrier imparted by hydrazines. *Chem. Commun.* 56: 9521–9524.
- 136 Lefebvre, V., Cailly, T., Fabis, F., and Rault, S. (2010). Two-step synthesis of substituted 3-aminoindazoles from 2-bromobenzonitriles. *J. Org. Chem.* 75: 2730–2732.
- 137 Lohou, E., Collot, V., Stiebing, S., and Rault, S. (2011). Direct access to 3-aminoindazoles by Buchwald–Hartwig C–N coupling reaction. *Synthesis* 16: 2651–2663.
- 138 Cyr, P., Régnier, S., Bechara, W.S., and Charette, A.B. (2015). Rapid access to 3-aminoindazoles from tertiary amides. *Org. Lett.* 17: 3386–3389.
- 139 Sajadi, M.S., Darehkordi, A., and Hosseini, S.M.S. (2021). Synthesis of *N*-aryl-3*H*-indazol-3-imine and *N*-aryl-1*H*-indazol-3-amine via Na₂WO₄/H₂O₂ mediated by intramolecular N–N coupling. *Tetrahedron* 84: 132023–132028.

- 140** Zhu, W.F., Krämer, A., Knapp, S. et al. (2022). Cascade synthesis of kinase-privileged 3-aminoindazoles via intramolecular N—N bond formation. *J. Org. Chem.* 87: 3856–3862.
- 141** Wang, G.-L., Chen, X., Chang, Y.-N. et al. (2015). Synthesis of 1,2,3-benzotriazin-4-one derivatives containing spirocyclic indoline-2-one moieties and their nematocidal evaluation. *Chin. Chem. Lett.* 26: 1502–1506.
- 142** Fiorino, F., Caliendo, G., Perissutti, E. et al. (2005). Synthesis by microwave irradiation and antiarrhoeal activity of benzotriazinone and saccharine derivatives. *Arch. Pharm. Chem. Life Sci.* 338: 548–555.
- 143** Chen, X., Jia, H., Li, Z., and Xu, X. (2019). Synthesis and nematocidal evaluation of 1,2,3-benzotriazin-4-one derivatives containing piperazine as linker against *Meloidogyne incognita*. *Chin. Chem. Lett.* 30: 1207–1213.
- 144** Chang, Y., Zhang, J., Chen, X. et al. (2017). Synthesis and nematocidal activities of 1,2,3-benzotriazin-4-one derivatives containing thiourea and acylthiourea against *Meloidogyne incognita*. *Bioorg. Med. Chem. Lett.* 27: 2641–2644.
- 145** Zhang, R., Guo, W., Wang, G. et al. (2020). Synthesis and nematocidal activities of 1,2,3-benzotriazin-4-one derivatives containing benzo[d][1,2,3]thiadiazole against *Meloidogyne incognita*. *Bioorg. Med. Chem. Lett.* 30: 127369–127372.
- 146** Caliendo, G., Fiorino, F., Grieco, P. et al. (1999). Preparation and local anaesthetic activity of benzotriazinone and benzoyltriazole derivatives. *Eur. J. Med. Chem.* 34: 1043–1051.
- 147** Moghimi, S., Goli-Garmroodi, F., Pilali, H. et al. (2016). Synthesis and anti-acetylcholinesterase activity of benzotriazinone-triazole systems. *J. Chem. Sci.* 128: 1445–1449.
- 148** Liu, C., Li, X., Zhang, J. et al. (2023). DNA-compatible benzotriazinone formation through aryl diazonium intermediates. *J. Org. Chem.* 88: 6565–6572.
- 149** Rocha-Alonzo, F., Chávez, D., Ochoa-Terán, A. et al. (2017). A novel synthesis of 1,2,3-benzotriazinones from 2-(o-aminophenyl)oxazolines. *J. Chemother.* 2017: <https://doi.org/10.1155/2017/2384780>.
- 150** El Rayes, S.M., Ali, I.A.I., Fathalla, W., and Mahmoud, M.A.A. (2020). Synthesis and biological activities of some new benzotriazinone derivatives based on molecular docking; promising HepG2 liver carcinoma inhibitors. *ACS Omega* 5: 6781–6791.
- 151** Miura, T., Yamauchi, M., Kosaka, A., and Murakami, M. (2010). Nickel-catalyzed regio- and enantioselective annulation reactions of 1,2,3,4-benzothiazine-1,1(2H)-dioxides with allenes. *Angew. Chem. Int. Ed.* 49: 4955–4957.