

## Index

Note: Page numbers in *italics* and **bold** indicates figures and tables respectively.

- a**
- acetylcholine probes 250–252, 251
  - acrylate recognition mechanism 90–93, 91–93
  - active molecules and enzymes
    - development of biomedical research 305–306
    - double enzyme 309–312, 311
    - drug discovery 305
    - real-time monitoring and imaging 305
    - reveal complex biological processes 305
  - ROS and enzymes 307–309, 308, 310
  - RSS and enzymes 306–307, 307
  - specificity and sensitivity of detection 305
  - activity-based sensing (ABS) 210, 210
  - adenosine triphosphate (ATP)
    - metal complexes-based fluorescent probes 343–344, 344
    - rhodamine-based irreversible fluorescent probes 342–343, 342, 343
    - rhodamine-based reversible fluorescent probes 339–342, 339–342
  - aggregation-induced emission (AIE) 35–36, 35, 72, 99
  - AIE-active red-emissive photosensitizer 77
  - $\alpha$ -ketoamide linkage strategy 48–49
  - aldehyde groups 138–140, 139
  - aldehyde-mediated cyclization 104
  - aldehyde recognition
    - mechanism 93–94, 94
  - allylamine 206–207, 206, 207
  - allyl ether quenching/recognition 198–199, 198, 199
  - allyl formate quenching/recognition 196–198, 196, 197
  - $\alpha$ ,  $\beta$ -unsaturated ketene 177–178, 178
  - $\alpha$ ,  $\beta$ -unsaturated ketone 238, 238
  - $\alpha$ -ketoamide groups 177–178, 178
  - anisotropy 27
  - a-PeT (acceptor-based PeT/reducing PeT) 29
  - Arabidopsis 194, 207, 210, 283, 331
  - aromatic substitution–rearrangement cascade reaction strategy 101
  - ATP-dual-responsive fluorescent probes
    - ROS and ATP 313–315, 314, 315
    - RSS and ATP 312–313
  - azide group reduction 122–124, 123

*Fluorescent Probes for Bioactive Species: Design, Synthesis, and Biomedical Applications*, First Edition. Edited by Caixia Yin. © 2026 Wiley-VCH GmbH. All rights reserved, including rights for text and data mining and training of artificial intelligence technologies or similar technologies. Published 2026 by Wiley-VCH GmbH.

**b**

- Baeyer–Villiger oxidative rearrangement reactions
- a-ketoamide linkage strategy 48–49
  - direct linkage strategy (a-ketocarbonyl) 47–48, 47
  - oxygenium bond strategy 48
- benzothiazoline 65–66, 66
- bicarbonitrile isophorone-based NADH probe 326, 326
- bioactive species fluorescent probes
- adenosine triphosphate (ATP) 338–344
  - lactic acid probes 331–332, 332
  - lipid peroxidation probes 332–338
  - NAD(P)H 319–331
  - pyruvic acid probes 331–332
- bioethanol identification methods 84
- biological functions
- reactive carbonyl substances (RCS) 193–195
  - reactive nitrogen species (RNS) 163–164
- BODIPY (Boron-Dipyrrromethene) dyes 13, 37, 206, 334–337, 334–336
- borate ester 173–174, 174
- boric acid 175–176, 175
- boric acid oxidation reactions
- direct linkage strategy 41–43, 42, 43
  - ester linkage strategy 44–45, 45
  - ethernet linkage strategy 43–44, 44
  - pyridine linkage strategy 46
- bovine serum albumin (BSA) 100
- c**
- Caenorhabditis elegans* 48
- caffeic acid 66–68, 66, 67
- carbon monoxide probes
- allyl ether quenching/ recognition 198–199, 198, 199
  - allyl formate quenching/ recognition 196–198, 196, 197
  - heavy atom quenching effect 199–201, 200
- nitro as the reaction site 202–203, 202, 203
  - Ru(II) vinyl complexes 203–205, 204
  - spirolactam ring opening 201–202, 201
- C=C unsaturated double bonds 57–58, 57, 58
- cellular signaling 352
- chemical messengers 243
- chemiluminescence (CL) signals 72
- Coelenterazine 72
- coumarin 96
- cyanide-mediated ATP depletion 344
- cyanine-based probe 323, 324
- cyclic disulfide-based fluorescent probes 236–237, 236
- cyclization reaction mechanism 102–104, 102, 103
- cyclohexanone-based fluorescent probe 322, 323
- cysteine (Cys) probes
- acrylate recognition mechanism 90–93, 91–93
  - aldehyde recognition mechanism 93–94, 94
- bioethanol identification methods 84
- chemical structure 96
- detection of 84
- metal complex recognition mechanism 89–90, 90
- nucleophilic substitution recognition mechanism 85–87, 85–87
- physiological level of 84
- role of 84
- Se–N bond cleavage recognition mechanism 87–88, 88, 89
- d**
- detecting mechanism
- H<sub>2</sub>O<sub>2</sub> and biological thiols 291–292, 292, 293
  - H<sub>2</sub>O<sub>2</sub> and H<sub>2</sub>S 289–290, 290
  - H<sub>2</sub>O<sub>2</sub> and SO<sub>2</sub> 290–291, 291

- HClO and biological thiols 284–286, 284–287
- HClO and H<sub>2</sub>S 279–283, 280–283
- HClO and sulfur dioxide 287–289, 288, 289
- hydroxyl radicals and RSS 295–296
- O<sub>2</sub><sup>•-</sup> and biological thiols 294
- O<sub>2</sub><sup>•-</sup> and H<sub>2</sub>S<sub>n</sub> 293, 294
- DIAKI pathogenesis 304
- dibenzoxazine-based fluorescent probe 321–322, 322
- dihydropyridine-based fluorescent probes 169, 169
- dimethylthiocarbamoyl chloride 56–57, 56
- dinitrile coumarin NAD(P)H probe 325, 325
- diphenylphosphine-based lipid peroxidation probes 337–338, 337
- diphenyl phosphinic acid 68–70, 68, 69
- diphenyl phosphoryl groups 176–177
- direct linkage strategy (a-ketocarbonyl) 41–43, 42, 43, 47–48, 47
- diselenide bond-based fluorescent probes 237–238, 237
- disulfide bonds 125–127, 126
- donor–acceptor–donor (D–A–D) 14
- dopamine (DA) probes coordination–substitution reaction 249–250, 250 nucleophilic substitution ring 250 phenylboric acid with catechol 249
- d-PeT (donor-based PeT/oxidizing PeT) 29
- D- $\pi$ -A naphthalene-BODIPY platform 294
- dual-response fluorescent probes, RSS and RNS active molecules and enzymes 304–312 active species, ATP 312–315 detecting H<sub>2</sub>O<sub>2</sub> and biological thiols 291–292, 292, 293 detecting H<sub>2</sub>O<sub>2</sub> and H<sub>2</sub>S 289–290, 290 detecting H<sub>2</sub>O<sub>2</sub> and SO<sub>2</sub> 290–291, 291 detecting HClO and biological thiols 284–286, 284–287 detecting HClO and H<sub>2</sub>S 279–283, 280–283 detecting HClO and sulfur dioxide 287–289, 288, 289 detecting hydroxyl radicals and RSS 295–296 detection efficiency 296 detection of O<sub>2</sub><sup>•-</sup> and biological thiols 294 detection of O<sub>2</sub><sup>•-</sup> and H<sub>2</sub>S<sub>n</sub> 293, 294 diagnosis and treatment 296–297 life science research 297 NO and thiols 297–298, 297, 299 ONOO<sup>-</sup> (oxidative cleavage) and SO<sub>2</sub> 298–299, 300 reactive oxygen species (ROS) 273–278 reactive sulfur species (RSS) 261–273 reveal the interaction 296 thiol and ONOO<sup>-</sup> 299–304, 302, 303 dual-salt fluorescent probe 327, 328
- e**
- electrophilic unsaturated C=C bonds 183–184, 183
- elimination reaction mechanism 104–106, 104–106
- energy gap (EG) law 7
- enhanced and quenched (turn on/off) turn-off probes (quenching type) 24 turn-on probes (enhanced type) 23–24
- Escherichia coli* 74
- ester bond breakage 132–133, 132
- ester linkage strategy 44–45, 45

ether bonds 178–179, 179  
 ethernet linkage strategy 43–44, 44  
 ethylene fluorescent probes  
   activity-based sensing  
     (ABS) 210, 210  
   olefins 209–210, 209  
   vinyl  $\pi$  and transition metals 209  
 excited-state intramolecular proton  
 transfer (ESIPT) 31–33, 32

**f**

fluorescence  
   discovery of 1–4  
   emerging trends in 15  
   energy gap (EG) law 7  
   Kasha's rule 6  
   Kasha–Vavilov rule 6  
   mechanism of 4–6  
   molecular identification 349  
   Nobel prize winner 16  
   quantum yield 7–8  
 fluorescence resonance energy transfer  
 (FRET) 86, 232  
 fluorescent dyes  
   chemical structures 13  
   classification of 8–9  
   development of 11–14  
   inorganic pigments 9–10  
   nitrobenz-2-oxa-1,3-diazole  
     (NBD) 263  
   protein dyes 11  
 fluorescent probes  
   excitation modes 19–23, 20  
   formaldehyde (FA) 207–209, 208  
   low-molecular-weight fluorescent  
     indicators 36–39  
   optical modulation  
     mechanism 27–36  
   oxidoreductase enzymes 351  
   reactive oxygen species 41–78  
   role of 350  
   single-photon excitation 20  
   two-photon excitation 20–22  
   upconversion luminescence 22–23

formaldehyde (FA)  
   allylamine 206–207, 206, 207  
   BODIPY dye 206  
   fluorescent probe 207–209, 208  
   free amino group ( $-\text{NH}_2$ )  
     205–206  
   hydrazine ( $-\text{NH}-\text{NH}_2$ ) 206  
 Förster resonance energy transfer  
 (FRET) 33–35, 33, 34  
 free amino group ( $-\text{NH}_2$ )  
   205–206

**g**

gamma-glutamyltranspeptidase  
 (GGT) 308  
 glutamic acid 252–253, 252  
 glutathione (GSH) probes  
   fluorescent probes 121  
   nucleophilic addition reaction  
     mechanism 109–113, 109–112  
   overview 108  
   reduction reaction mechanism  
     119–120, 120  
   substitution reaction mechanism  
     113–119, 113–115, 117, 119  
 green fluorescent protein (GFP) 4

**h**

heavy atom quenching effect  
 199–201, 200  
 hepatic ischemia-reperfusion injury  
 (HIRI) 313  
 histamine 254–255, 255  
 homocysteine (Hcy) probes  
   chemical structure 96  
   cyclization reaction mechanism  
     102–104, 102, 103  
   elimination reaction mechanism  
     104–106, 104–106  
   metal complex reaction  
     mechanism 106–107, 107  
 Michael addition reaction  
   mechanism 96–98, 97  
 overview 95–96

- rearrangement reaction
    - mechanism 98–102, 98–101
  - HUMO levels 29
  - hydrazine (–NH – NH<sub>2</sub>) 206
  - hydrazine/phenylhydrazine-modified
    - spirolactam 53–54, 53, 54
  - hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>)
    - aryl sulfonyl esters 49, 49
    - Baeyer–Villiger oxidative
      - rearrangement reactions 46–49
    - boric acid oxidation reactions 41–46
    - tandem Pinnick/Dakin reaction 49
  - hydrogen sulfide (H<sub>2</sub>S) probes
    - azide group reduction 122–124, 123
    - 2,4-dinitrophenyl ether bond
      - 127–128, 128
    - 2,4-dinitrosulfonic Acid
      - 128–130, 129
    - disulfide bonds 125–127, 126
    - ester bond breakage 132–133, 132
    - Michael addition reaction
      - 133–134, 134
    - NBD cleavage 130–132, 131
    - nitro reduction, amine group
      - 124–125, 125
    - overview 121–122
    - response mechanisms 134–136
  - 4-hydroxybenzyloxy moiety-based
    - fluorescent probes 232–233
  - hydroxyl radical fluorescent probes
    - 60–64, 61–63
  - 5-hydroxytryptamine
    - aldehyde amine condensation
      - reaction 253
    - substitution–cyclization strategy
      - identification 253–254, 253, 254
  - hypobromous acid (HBrO) 308
  - hypochlorous acid
    - C=C unsaturated double bonds
      - 57–58, 57, 58
    - dimethylthiocarbamoyl
      - chloride 56–57, 56
    - nitrogen-containing functional
      - groups 52–56
    - oxidation and breakage of amide
      - bonds 58–60, 59, 60
    - oxygen-containing functional
      - groups 57
    - single-electron oxidation process
      - 50–52, 50, 51
- i**
- imidazolpyrazinone 71–73, 72
  - indocyanine green (ICG) 12
  - intramolecular charge transfer
    - (ICT) 29–31
  - intramolecular electron transfer
    - (IET) 15
- k**
- Kasha's rule 6
  - Kasha–Vavilov rule 6
- l**
- lactic acid probes 331–332, 332
  - L-carnitine treatment 304
  - levulinate groups 137–138, 137
  - light absorption 9
  - lipid peroxidation probes
    - BODIPY 334–337, 334–336
    - design and reaction
      - mechanism 333, 333
    - diphenylphosphine 337–338, 337
    - polyunsaturated fatty acids
      - (PUFAs) 332, 333
  - low-molecular-weight fluorescent
    - indicators
      - chemical covalent bond formation
        - and cleavage 36–37, 37
      - coordination reaction 38, 38
      - protonated/deprotonated
        - mechanism 38–39, 39
  - LUMO levels 29
- m**
- malonaldehyde probes 211–212,
    - 211, 212
  - metal complexes-based fluorescent
    - probes 343–344, 344

- metal complex reaction
  - mechanism 106–107, 107
- metal complex recognition
  - mechanism 89–90, 90
- metastable intermediate 22
- Michael addition 271–273, 272, 273
- Michael addition reaction 96–98, 97, 109, 133–134, 134
- mitochondria 334
- mitochondrial dysfunction 315
- molecular recognition 1
- monoamine oxidases (MAOs)
  - MAO-A fluorescent probes 216–218, 216–218
  - MAO-B fluorescent probes 218–220, 219, 220
  - overview 215–216
- monoamine probes 215–220
- multispectral optoacoustic tomography (MSOT) imaging 35
  
- n**
- NAD(P)H 353
  - cyclohexanone-based fluorescent probe 322, 323
  - dibenzoxazine-based fluorescent probe 321–322, 322
  - pyridine-based fluorescent probe 322–323, 323
  - quinoline-based fluorescent probes 323–330, 324–331
  - quinone-based fluorescent probe 319–320
  - resazurin-based fluorescent probe 320–321
- naphthylimides-based NAD(P)H probe 326, 326
- NBD cleavage 130–132, 131
- neurotransmitters 352
  - acetylcholine probes 250–252, 251
  - autonomic nervous system regulation 244
  - biological function of 243–245
  - DA probes 249–250
  - emotion and behavior 244
  - glutamic acid 252–253, 252
  - histamine 254–255, 255
  - hormone secretion and regulation 245
  - 5-hydroxytryptamine 253–254
  - learning and memory 244
  - motor control 244
  - NE probes 245–249
  - pain perception and modulation 245
  - pH-responsive FFNs 256
  - vesicular monoamine transporter 2 (VMAT2) 255
- neutrophil elastase (NE) probes
  - aldo-amine condensation reaction 245–246, 245
  - dual-site cooperative reaction 246–247, 246
  - recognition strategy 247–249, 247, 249
- nitric oxide (NO) probes
  - design strategies 164, 165
  - dihydropyridine-based fluorescent probes 169, 169
  - o-diaminoaromatic fluorescent probes 165–168, 166
  - ortho-phenylamine cyclization 171–172, 172
  - oxidation of secondary amines 170–171, 170
  - oxidative amination 168–169, 168
- nitro as the reaction site 202–203, 202, 203
- nitrobenz-2-oxa-1,3-diazole (NBD) 263
- nitrobenzene fluorescent probes 225–229, 226, 228, 229
- nitro-covalently linked fluorescent probes 230–232, 231
- nitrofuran fluorescent probes 230, 231
- nitrogen-containing functional groups 52–56
  - cleavage of p-methoxyaniline 54, 55

- hydrazine/phenylhydrazine-modified
    - spirolactam 53–54, 53, 54
  - oxidation of hydroxamic
    - acid 52–53, 52
  - oxidation of pyrrole
    - groups 54–56, 55
  - nitroimidazole fluorescent probes
    - 229–230, 230
  - nitroreductase probes
    - (NTRs) 225–232
  - nitrobenzene fluorescent probes
    - 225–229, 226, 228, 229
  - nitro-covalently linked fluorescent
    - probes 230–232, 231
  - nitrofurans fluorescent
    - probes 230, 231
  - nitroimidazole fluorescent
    - probes 229–230, 230
  - nitro reduction, amine group
    - 124–125, 125
  - nitroxy (HNO)
    - non-organelle targeting type
      - 184–186, 185
    - organelle targeting type 186–187, 187
    - ratiometric type 187–188, 188
    - redox-related RNS 350
  - N-dearylation 181–183, 182
  - N-methyl-p-hydroxyphenylamine 33
  - N-nitrosation 181, 182
  - nucleophilic addition reaction
    - mechanism 109–113, 109–112
  - nucleophilic substitution
    - cyclization reaction mechanism
      - 102–104, 102, 103
    - elimination reaction mechanism
      - 104–106, 104–106
    - rearrangement reaction
      - mechanism 98–102, 98–101
  - nucleophilic substitution
    - rearrangement
      - double response probes 262–268, 262–267
      - multiplexed RSS-responsive
        - probes 268–271, 268–271
    - nucleophilic substitution recognition
      - mechanism 85–87, 85–87
- O**
- o-diaminoaromatic fluorescent
    - probes 165–168, 166
  - O-GlcNAcase (OGA) 312
  - olefins 209–210, 209
  - one-photon microscopy (OPM) 119
  - o-phenylenediamine (OPD) 266
  - optical modulation mechanism
    - aggregation-induced emission (AIE)
      - 35–36, 35
    - excited-state intramolecular proton
      - transfer (ESIPT) 31–33, 32
    - Förster resonance energy transfer
      - (FRET) 33–35, 33, 34
    - intramolecular charge transfer
      - (ICT) 29–31
    - photon-induced electron transfer
      - (PeT) 28–29, 28, 29
  - organic fluorescent dyes 11–14
  - ortho-phenylamine cyclization
    - 171–172, 172
  - Otto Loewi's acetylcholine 352
  - oxidation
    - amination 168–169, 168
    - and breakage of amide bonds
      - 58–60, 59, 60
    - dearomatization 179–180, 180
    - hydrolysis of hydrazides 180, 181
    - of hydroxamic acid 52–53, 52
    - of pyrrole groups 54–56, 55
    - of secondary amines 170–171, 170
  - oxidative furan ring-opening 76–77
  - oxidoreductase enzymes 351
    - monoamine probes 215–220
    - nitroreductase probes (NTRs) 225–232
    - quinine oxidoreductase
      - probes 220–224
    - in redox modulation 215
    - thioredoxin reductase (TrxR)
      - enzymes 236–238
    - tyrosinase probes 232–235

oxygen-containing functional groups 57  
oxygenium bond strategy 48

**p**

peroxynitrite (ONOO<sup>-</sup>) probes  
   $\alpha$ -ketoamide groups 177–178, 178  
   $\alpha$ ,  $\beta$ -unsaturated ketene 177–178, 178  
  borate ester 173–174, 174  
  boric acid 175–176, 175  
  diphenyl phosphoryl groups 176–177  
  electrophilic unsaturated C=C bonds 183–184, 183  
  ether bonds 178–179, 179  
  N-dearylation 181–183, 182  
  N-nitrosation 181, 182  
  overview 173  
  oxidative dearomatization 179–180, 180  
  oxidative hydrolysis of hydrazides 180, 181  
photon-induced electron transfer (PeT) 28–29, 28, 29, 112, 331  
polysulfide probes  
  H<sub>2</sub>S<sub>2</sub>-mediated addition and cleavage reactions 148–149, 149  
  H<sub>2</sub>S<sub>2</sub>-mediated aromatic nucleophilic substitution 153–154, 154  
  H<sub>2</sub>S<sub>2</sub>-mediated departure of 2-(benzoylthio)benzoate 145–147, 146  
  H<sub>2</sub>S<sub>2</sub>-mediated departure of p-nitrofluorobenzene 143–145, 144  
  H<sub>2</sub>S<sub>2</sub>-mediated dithiole formation reaction 154–155, 155  
  H<sub>2</sub>S<sub>2</sub>-mediated electrophilic reactions 150–151, 150, 151  
  H<sub>2</sub>S<sub>2</sub>-mediated nucleophilic addition 147–148, 147, 148  
  H<sub>2</sub>S<sub>2</sub>-mediated reduction 151–153, 152, 153

polyunsaturated fatty acids (PUFAs) 332, 333  
Protect-Deprotect strategy 247  
protein O-GlcNAcylation 312  
pyridine-based fluorescent probe 322–323, 323  
pyridine linkage strategy 46  
pyruvic acid probes 331–332

**q**

quenching effects 15  
quinine oxidoreductase probes 220–224  
  bioluminescent probes 224, 224  
  chemiluminescent probes 223–224, 223  
  NQO1 fluorescent probes 221, 221  
  open fluorescent probes 221–222, 222  
  ratiometric fluorescent probes 222–223  
quinoline-based fluorescent probes 323–330, 324–331  
quinone-based fluorescent probe 319–320

**r**

ratiometric probes 24–25, 25  
reactive carbonyl substances (RCS) 351  
  biological functions of 193–195  
  carbon monoxide 194  
  cytotoxic mechanisms 194  
  ethylene 195  
  formaldehyde (FA) 195  
  signaling molecules 193  
reactive nitrogen species (RNS) 350  
  biological functions of 163–164  
  nitric oxide (NO) 164–173  
  nitroxyl (HNO) 184–189  
  peroxynitrite (ONOO<sup>-</sup>) probes 173–184  
reactive oxygen/nitrogen species (RONS) family 60

- reactive oxygen species (ROS)  
 biomedical research 274  
 detection sensitivity 274  
 HClO and H<sub>2</sub>O<sub>2</sub> 276  
 HClO and ONOO<sup>-</sup> 276–277,  
 277, 278  
 multi-target detection 274  
 peroxyxynitrite (ONOO<sup>-</sup>)  
 274–276, 275  
 reduce false positive results 274  
 superoxide anion (O<sub>2</sub><sup>-</sup>)  
 274–276, 275
- reactive sulfur species (RSS)  
 biological functions of 83  
 cysteine (Cys) probes 84–95  
 double response probes 262–268,  
 262–267  
 glutathione (GSH) probes  
 108–121  
 homocysteine (Hcy) probes  
 95–108  
 hydrogen sulfide (H<sub>2</sub>S)  
 probes 121–136  
 Michael addition 271–273, 272, 273  
 multiplexed RSS-responsive  
 probes 268–271, 268–271  
 polysulfide probes 143–155  
 sulfur dioxide (SO<sub>2</sub>) probes 136–143
- reduction reaction mechanism  
 119–120, 120
- reporting mode, fluorescence  
 anisotropy 27  
 enhanced and quenched  
 (turn on/off) 23–24  
 lifetime-based sensing 25–27, 26  
 ratiometric probes 24–25, 25
- resazurin-based fluorescent  
 probe 320–321
- reversible photoacoustic probe 294
- rhodamine-based irreversible  
 fluorescent probes 342–343,  
 342, 343
- rhodamine-based NADH probe 324,  
 324, 325
- rhodamine-based reversible fluorescent  
 probes 339–342, 339–342
- Ru(II) vinyl complexes 203–205, 204
- S**
- Schizophrenia 70
- Se–N bond cleavage recognition  
 mechanism 87–88, 88, 89
- single-electron oxidation process  
 50–52, 50, 51
- single-photon excitation 20
- singlet oxygen (<sup>1</sup>O<sub>2</sub>) fluorescence probe  
 [4+2] cycloaddition peroxide  
 bonds 74–76  
 double bond to form an aldehyde  
 group 77–78  
 overview 73–74  
 oxidative furan ring-opening 76–77
- spirolactam ring opening  
 201–202, 201
- substitution reaction mechanism  
 113–119, 113–115, 117, 119
- sulfur dioxide (SO<sub>2</sub>) probes  
 aldehyde groups 138–140, 139  
 double bond reaction site  
 140–142, 141  
 levulinate groups 137–138, 137  
 overview 136–137
- sulfur-substituted hemianthocyanine  
 fluorescent probe 127
- superoxide anion (O<sub>2</sub><sup>-</sup>)  
 benzothiazoline 65–66, 66  
 caffeic acid 66–68, 66, 67  
 diphenyl phosphinic acid 68–70,  
 68, 69  
 imidazolpyrazinone 71–73, 72  
 overview 64–65  
 trifluoromethanesulfonate 70–71,  
 70, 71
- surface-enhanced Raman spectroscopy  
 (SERS)-assisted fluorescence 266
- t**
- tetraphenylethylene (TPE) 230
- thiol dual-responsive probes 266

- thioredoxin reductase (TrxR)
    - enzymes 236–238
  - thioredoxin reductase (TrxR) probes
    - $\alpha$ ,  $\beta$ -unsaturated ketone 238, 238
    - diselenide bond-based fluorescent probes 237–238, 237
    - five-membered cyclic disulfide-based fluorescent probes 236–237, 236
  - trifluoromethanesulfonate 70–71, 70, 71
  - turn-off probes (quenching type) 24
  - turn-on probes (enhanced type) 23–24
  - two-photon excitation 20–22
  - two-photon-excited fluorescence (TPEF) 119
  - tyrosinase probes 232–235
    - dopamine-based fluorescent probes 235, 235
    - 4-hydroxybenzyloxy moiety-based fluorescent probes 232–233
    - meta-hydroxybenzyloxy moiety-based fluorescent probes 233–235, 233, 234
- V**
- vinyl  $\pi$  and transition metals 209







